

Kansas Agricultural Experiment Station Research Reports

Volume 0
Issue 1 *Cattleman's Day (1993-2014)*

Article 357

2001

Performance of beef heifers limit-fed growing diets containing alfalfa hay and wet corn gluten feed

Sean P. Montgomery

J.J. Sindt

T.B. Farran

See next page for additional authors

Follow this and additional works at: <https://newprairiepress.org/kaesrr>



Part of the [Other Animal Sciences Commons](#)

Recommended Citation

Montgomery, Sean P.; Sindt, J.J.; Farran, T.B.; Pike, N.J.; Coetzer, C.M.; LaBrune, H.J.; Trater, A.M.; Hunter, R.D.; Stock, R.A.; and Drouillard, James S. (2001) "Performance of beef heifers limit-fed growing diets containing alfalfa hay and wet corn gluten feed," *Kansas Agricultural Experiment Station Research Reports*: Vol. 0: Iss. 1. <https://doi.org/10.4148/2378-5977.1760>

This report is brought to you for free and open access by New Prairie Press. It has been accepted for inclusion in Kansas Agricultural Experiment Station Research Reports by an authorized administrator of New Prairie Press. Copyright 2001 Kansas State University Agricultural Experiment Station and Cooperative Extension Service. Contents of this publication may be freely reproduced for educational purposes. All other rights reserved. Brand names appearing in this publication are for product identification purposes only. No endorsement is intended, nor is criticism implied of similar products not mentioned. K-State Research and Extension is an equal opportunity provider and employer.



Performance of beef heifers limit-fed growing diets containing alfalfa hay and wet corn gluten feed

Abstract

Three hundred thirty-nine crossbred beef heifers were used in a 99-day growing study to identify optimum combinations of alfalfa hay and wet corn gluten feed (WCGF) in limit-fed growing diets. Diets contained 10, 20, or 30% ground alfalfa hay, and 0, 40, or 68% Sweet Bran® WCGF (dry basis) in a 3 x 3 factorial arrangement of treatments. An interaction occurred ($P < 0.05$) between level of alfalfa hay and level of WCGF for both average daily gain and feed efficiency. Increasing the levels of alfalfa hay or WCGF reduced cattle performance, with the exception of the 30% alfalfa hay and 40% WCGF diet, which supported average daily gains similar ($P > 0.10$) to diets containing 20 or 30% alfalfa hay and no WCGF. Feed efficiencies for the 30% alfalfa hay and 40% WCGF diet were better ($P < 0.05$) than the diet containing 30% alfalfa hay and no WCGF. Dry matter intake as measured two hours after feeding increased linearly ($P < 0.01$) with increasing levels of alfalfa hay, and decreased linearly ($P < 0.01$) with increasing levels of WCGF. This study suggests that including WCGF at 40% of the diet (dry basis) can effectively replace steam-flaked corn in limit-fed diets containing 20 or 30% alfalfa hay.

Keywords

Cattlemen's Day, 2001; Kansas Agricultural Experiment Station contribution; no. 01-318-S; Report of progress (Kansas State University. Agricultural Experiment Station and Cooperative Extension Service); 873; Beef; Wet corn gluten feed; Alfalfa hay; Limit feeding; Growing

Creative Commons License



This work is licensed under a [Creative Commons Attribution 4.0 License](https://creativecommons.org/licenses/by/4.0/).

Authors

Sean P. Montgomery, J.J. Sindt, T.B. Farran, N.J. Pike, C.M. Coetzer, H.J. LaBrune, A.M. Trater, R.D. Hunter, R.A. Stock, and James S. Drouillard

PERFORMANCE OF BEEF HEIFERS LIMIT-FED GROWING DIETS CONTAINING ALFALFA HAY AND WET CORN GLUTEN FEED

S. P. Montgomery, J. S. Drouillard, J. J. Sindt, T. B. Farran,
N. J. Pike, C. M. Coetzer, H. J. LaBrune,
A. M. Trater, R. D. Hunter, and R. A. Stock¹

Summary

Three hundred thirty-nine crossbred beef heifers were used in a 99-day growing study to identify optimum combinations of alfalfa hay and wet corn gluten feed (WCGF) in limit-fed growing diets. Diets contained 10, 20, or 30% ground alfalfa hay, and 0, 40, or 68% *Sweet Bran*[®] WCGF (dry basis) in a 3 × 3 factorial arrangement of treatments. An interaction occurred ($P < 0.05$) between level of alfalfa hay and level of WCGF for both average daily gain and feed efficiency. Increasing the levels of alfalfa hay or WCGF reduced cattle performance, with the exception of the 30% alfalfa hay and 40% WCGF diet, which supported average daily gains similar ($P > 0.10$) to diets containing 20 or 30% alfalfa hay and no WCGF. Feed efficiencies for the 30% alfalfa hay and 40% WCGF diet were better ($P < 0.05$) than the diet containing 30% alfalfa hay and no WCGF. Dry matter intake as measured two hours after feeding increased linearly ($P < 0.01$) with increasing levels of alfalfa hay, and decreased linearly ($P < 0.01$) with increasing levels of WCGF. This study suggests that including WCGF at 40% of the diet (dry basis) can effectively replace steam-flaked corn in limit-fed diets containing 20 or 30% alfalfa hay.

(Key Words: Wet Corn Gluten Feed, Alfalfa Hay, Limit Feeding, Growing.)

Introduction

Wet corn gluten feed, a by-product of corn wet milling, consists mainly of corn

bran and corn steep liquor. Because of high levels of corn bran, WCGF can supply additional energy for maintenance and growth without the reduction in fiber digestion often seen when substantial quantities of high-starch ingredients (grain) are fed to cattle consuming high-roughage diets. This study was conducted to identify optimum combinations of alfalfa hay and WCGF in limit-fed growing diets containing steam-flaked corn.

Experimental Procedures

Three hundred thirty-nine crossbred beef heifers averaging 609 lb were used in a randomized complete block experiment. To minimize differences in gastrointestinal tract fill, heifers had *ad libitum* access to a common diet for 15 days preceding the growing study. Heifers were then blocked by weight and allotted to pens containing four to seven head per pen, with six pens per treatment. Treatments (Table 1) consisted of diets containing 10, 20, or 30% ground alfalfa hay, and 0, 40, or 68% Sweet Bran WCGF (dry basis) in a 3 × 3 factorial arrangement. All diets provided 30 grams of Rumensin[®] per ton of dry matter and were fed once daily at 1.6% of body weight (dry basis) for 84 days. On days 8, 22, 37, 51, 64, and 79, unconsumed feed was removed from the feed bunks 2 hours after feeding, immediately weighed, and returned to the bunk in order to measure rate of feed consumption. Prior to obtaining final weights, heifers had *ad libitum* access to a common diet for 15 days.

¹Cargill Corn Milling, Blair, NE.

Results and Discussion

Performance data are shown in Figures 1 and 2. There was an interaction between alfalfa hay and WCGF level in the diet. Increasing the level of alfalfa hay and WCGF reduced growing performance of heifers, with the exception of the diet containing 30% alfalfa hay and 40% WCGF. Heifers fed that diet had average daily gains similar ($P > 0.10$) to those consuming diets containing 20 or 30% alfalfa hay and no WCGF. They also had improved feed efficiencies compared to heifers fed the diet containing 30% alfalfa hay and no WCGF ($P < 0.05$). Dry matter

intake was not different among treatments ($P > 0.05$) ($11.4 \pm .4$ lb/day), so these results indicate a positive associative effect of WCGF on the digestibility of alfalfa hay in the diet containing 30% alfalfa hay and 40% WCGF. Increasing the level of alfalfa hay in the diet increased ($P < 0.01$) dry matter intake as measured 2 hours after feeding (Figure 3), whereas increasing the level of WCGF decreased ($P < 0.01$) dry matter intake. We conclude that the value of WCGF relative to steam-flaked corn increases in diets containing greater amounts of roughage when WCGF is included at 40% of the diet dry matter.

Table 1. Experimental Diets (% of Dry Matter)

Ingredients	% Alfalfa Hay	Treatments								
		10	10	10	20	20	20	30	30	30
	% WCGF	0	40	68	0	40	68	0	40	68
Steam-flaked corn		73.4	48.6	20.0	65.3	39.2	10.0	57.1	29.5	-
Alfalfa hay		9.8	9.9	10.0	19.6	19.9	20.1	29.5	30.0	30.3
Wet corn gluten feed		-	38.9	67.1	-	39.0	67.3	-	39.1	67.6
Soybean meal		9.0	-	-	7.6	-	-	6.2	-	-
Cane molasses		4.9	-	-	4.9	-	-	4.9	-	-
Urea		1.0	0.3	-	1.0	0.1	-	1.0	-	-
Limestone		1.2	1.6	2.3	0.8	1.2	2.0	0.5	0.9	1.6
Calcium phosphate		0.2	-	-	0.3	-	-	0.3	-	-
Sodium chloride		0.4	0.4	0.4	0.4	0.4	0.4	0.4	0.4	0.4
Potassium chloride		-	0.2	0.1	-	0.1	0.1	-	-	-
Vitamin/trace mineral premix ^a		0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1
Crude protein, analyzed		16.1	16.5	20.1	16.1	16.8	20.8	16.1	17.0	21.4

^aFormulated to provide (total diet dry matter): 1,200 IU/lb vitamin A, 0.1 ppm cobalt, 10 ppm copper, 0.6 ppm iodine, 60 ppm manganese, 0.2 ppm selenium, 60 ppm zinc and 30 g/ton of Rumensin.




Figure 1. Average Daily Gain of Heifers Limit-Fed Growing Diets Containing 10, 20, or 30% Alfalfa Hay and 0, 40, or 68% Wet Corn Gluten Feed. Interaction between alfalfa hay and WCGF level in diet ($P < 0.05$).




Figure 2. Feed:Gain of Heifers Limit-Fed Growing Diets Containing 10, 20, or 30% Alfalfa Hay and 0, 40, or 68% Wet Corn Gluten Feed. Interaction between alfalfa hay and WCGF level in diet ($P < 0.05$).




Figure 3. Dry Matter Intake as Measured Two Hours After Feeding. Linear effect of alfalfa hay ($P < 0.01$) and linear effect of WCGF ($P < 0.01$).