New markers and linkage data

D. D. Perkins
Stanford University

M. R. Smith
Stanford University

D. R. Galeazzi
Stanford University

Follow this and additional works at: <https://newprairiepress.org/fgr>

This work is licensed under a [Creative Commons Attribution-Share Alike 4.0 License](https://creativecommons.org/licenses/by-sa/4.0/).

Recommended Citation

Perkins, D. D., M.R. Smith, and D.R. Galeazzi (1973) "New markers and linkage data," *Fungal Genetics Reports*: Vol. 20, Article 50. <https://doi.org/10.4148/1941-4765.1858>

This Linkage Data is brought to you for free and open access by New Prairie Press. It has been accepted for inclusion in Fungal Genetics Reports by an authorized administrator of New Prairie Press. For more information, please contact cads@k-state.edu.

New markers and linkage data

Abstract

New markers and linkage data

New markers and linkage data.

have not been completed. Strain T51M171 is probably a point mutant at a new locus, with poor ascospore ripening and viability, but chromosome aberration has not been excluded. The requirements of T51M152(t), T42M59(t), T51M147(t) have been identified and confirmed by H. Inoue and T. Ishikawa (personal communication). Locus assignments for 1003 at act-1, T51M147(t) at asp, and 76P.5 at rib-1 are based on map location rather than direct allelism tests.

Linkage data are given in Tables 2 and 3. [following pages] There concern not only markers in Table 1, but also give new information on other markers and intervals. The usual conventions are followed: the top number of each pair of complementary classes represents progeny of the genotype that contains the wild-type allele of the leftmost marker scored. Regions are numbered from left to right. Where dashes replace numbers, the corresponding classes of segregants were not scored. Interval lengths have usually been rounded to the nearest digit, because it was felt that showing tenths of units conveyed a false impression of precision and predictability.

pi (B101) and col-10 (R2438) are distinctly different in appearance, and are capable of recombination to give true wild-type morphology. It is therefore suggested that separate names be retained, although Srb and Basl (1969 *Neurospora* Newsl. 15: 22) have reported that pi and col-10 do not show complementation with respect to ascus morphology. Genes thr-2 and thr-3 have never recombined in our experience. They were designated separate loci on the basis of phenotypic differences and complementation (Teas 1947 Ph.D. Thesis, Calif. Inst. Technol.; Emerson 1950 Cold Spring Harbor Symp. Quant. Biol. 14:40). Mutant T42M59(t) is designated thr-3 because it resembles thr-3 allele 44104(t) phenotypically and does not resemble thr-2. The mutant spco-15 (R2537), reported to be on III by Garnjobst and Tatum (1967 *Genetics* 57:579), shows no linkage in 111 when extracted from the original strain spco-15; inos A (FGSC #1799). Possibly the III linkage was due to a reparable aberration.

Recently published linkage data: The following is an incomplete list of linkages that have been published from this and other laboratories since the 1969 tabulation of Perkins et al. in *Genetics* 40: 271.

Linkage Group I:

aza-1 (10-20) mt; aza-2 near mt (Jho 1971 *Mol. Gen. Genet.* 114: 156)
leu-3 (12) mt (1.6) upr-1 (7) org-1 (Tuveson 1972 *Mutat Res.* 15:411)
mt sru-2 sru-3 al-2 (Seale 1972 *Genetics* 70:385).
m+ ret-3 arg-3 (Catcheside et al. 1971 *Austral. J. Biol. Sci.* 24: 107)
cr (1 1) uvs-6 (6) al-2 (Schroeder et al. 1972 *Neurospora* Newsl. 19: 17)
nit-2 (17) mig tre (20) al-2 (Sussman et al. 1971 *J. Bacteriol.* 108:59)
ty-2 right of al-2 (Ho 1972 *Neurospora* Newsl. 19: 15)
or-1 (13) arom-8.(4) R (Turner et al. 1969 *Can. J. Genet Cytol* 11:622)

Linkage Group II:

worn-1 (2) ff-1 (4) mo (P240at) (Ton 1972 M. S. Thesis)

Linkage Group III:

prol-1 ff-5 me-8 hist-7 (Tan 1972 M. S. Thesis)

Linkage Group IV:

cys-10 centromere un-8 (5) col-4 (Perkins et al. 1972 *Neurospora* Newsl. 19:24)
pdx-1 (2) mtr (1) col-4 (0.2) org-2 (Boone et al. 1970 *Neurospora* Newsl. 16:12)
cys-10 (25) pyr-1 (3) pdx-1 (2) col-4 (1) arg-2 (16) col-1 (25) uvs-2 (D. A. Smith 1968 M. S. Thesis)

Linkage Group V:

sp (5) am-1 (0.05) gul-1 (3) hist-1 (Smyth 1970 Ph. D. Thesis, Austral. National Univ.)
al-3 near arg-8 and inos (Wang et al. 1971 *Neurospora* Newsl. 18:16)
inos (15) pab-2 (9) pyr-6 (Caroline 1969 *J. Bacteriol.* 100: 1371)

Linkage Group VI:

ad-8 (8) ssu-7 (14) ylo-1 (Seale 1972 *Genetics* 70: 385)

Linkage Group VII:

nit-3 (31) me-7 (3.5) wc (2.5) aga (27) arg-10 (Morgan 1969 *John Inner Inst., Ann. Rept.* 60:46)
thi-3 ars-1 (<1) me-7 me-9 wc (Murray 1970 *Genet. Res.* 15: 109)
nit-3 ssu-4 me-7 ssu-1 Su-1 nt (Seale 1972 *Genetics* 70:385)

We propose henceforth to limit the publication of routine mapping data from this laboratory to the Newsletter, which seems ideally suited for the purpose, unless such data are logically presented in the context of a journal paper devoted to a broader subject.

We are grateful to R. J. Lloyd and R. E. Padilla for technical assistance and to M. Flavin, T. Ishikawa, D. Newmeyer, M. L. Pall and B.C. Turner for stocks and unpublished information. - - - Department of Biological Sciences, Stanford University, Stanford, California, 94305.

Table 1. Mutants previously unassigned to linkage groups.

Locus symbol and name	Isolation No.	Linkage group and arm	Approximate location or evidence of allelism	Phenotype	Origin (strain, mutagen)	Reference to origin
<u>atr-1</u> : aminotriazole-1	RC2(r)	IL	0/39 recombination x <u>suc</u> /	Resistant 3-AT 1 mg/ml Abnormal morphology	74A, diethyl sulfate	M. L. Pall, unpubl.
<u>acr-7</u> : acriflavine-7	P1676(r)	VIR	Near <u>ws-1</u>	Resistant to acriflavine 0.5 mg/ml	Mixed, spontaneous?	
<u>bg</u> : brown ascospore	AR62	IR	R. of <u>un-1</u> (9%)	Autonomous, viable	OR23-1A, UV	
<u>un-9</u> : unknown-9	<u>T54M56</u> (t)	VR	R. of <u>pyr-6</u> (2%)	Irreparable temperature-sensitive (34° vs 25°)	74A, UV	Inoue & Ishikawa, 1970 un(42)
<u>un-10</u>	<u>T42M25</u> (t)	VIIR	R. of <u>wc</u> (7%)	"	74A, UV	" un(11)
<u>un-11</u>	<u>T42M30</u> (t)	VR	0/47 x <u>al-3</u>	"	74A, UV	" un(10)
<u>un-12</u>	<u>T51M118</u> (t)	IVR	0/73 x <u>col-4</u>	"	74A, X-rays	" un(17)
<u>un-13</u>	<u>T42M24</u> (t)	VI	2/106 x <u>ylo-1</u>	"	74A, UV	" un(9)
<u>un(T42M56)</u>	<u>T42M56</u> (t)	VI	0/81 x <u>ylo-1</u>	"	74A, UV	" un(12)
<u>un(T42M69)</u>	<u>T42M69</u> (t)	IL	0/86 x <u>mt</u>	"	74A, UV	" un(16)
<u>un(T51M154)</u>	<u>T51M154</u> (t)	VIL	0/44 x <u>lys-5</u>	"	74A, X-rays	" un(22)
<u>un(T51M171)</u>	<u>T51M171</u> (t)	IIIR	R. of <u>tyr-1</u> (5%)	"	74A, X-rays	" un(25)
<u>un(T54M55)</u>	<u>T54M55</u> (t)	III	Linked <u>acr-2</u> (8%)	"	74A, UV	" un(36)
RECURRENCES:						
<u>acr-2</u>	<u>1001</u> (r)	IIIL	0/46 wt x KH2		SL, UV	Howe & Terry 1962
<u>acr-2</u>	<u>1002</u> (r)	IIIL	0/38 wt x KH2		SL, UV	"
<u>act-1</u> (probable)	<u>1003</u> (r)	IR	Linked <u>aur</u> (10%), M. Flavin		SL, UV	"
<u>asp</u>	<u>T51M147</u> (t)	VR	Between <u>al-3</u> , <u>hist-6</u>	Temperature-sensitive	74A, X-rays	Inoue et al. 1970 un(19)
<u>fl</u>	P4499	IIR	0/37 wt x <u>fl</u> ^P		Mixed, spontaneous	
<u>hist-2</u>	<u>T51M152</u> (t)	IC	0/600 wt x Y152M14	Temperature-sensitive	74A, X-rays	" un(20)
<u>inos</u>	P47231	VR	0/200 wt x 37401		Mixed, spontaneous	
<u>os-1</u>	NM204(t)	IR	0/90 wt x B135	Nearly wild at 25°	Ems, UV	
<u>os-5</u>	P5341	IR	0/82 wt x NM2160		Mixed, spontaneous	
<u>rib-1</u>	76R5	VIR	0/193 x <u>pan-2</u>		Mixed, UV	
<u>thr-3</u> (probable)	<u>T42M59</u> (t)	IIL	0/73 wt x 44104(t)	Temperature-sensitive	74A, UV	Inoue et al. 1970 un(14)
<u>un-2</u>	ALS9(t)	IC	0/87 wt x 46006(t), D. Newmeyer		SL, UV	
<u>wc</u>	P4723w	VIIR	0/47 wt x P829		Mixed, spontaneous	
<u>ylo-1</u>	P5241	VIL	0/38 wt x Y30539y		Mixed, spontaneous	

Table 2. Data from 2- and 3-point crosses.

Zygote genotype and % recombination		Paren- tals	Crossovers			Isolation Numbers
			1	2	1,2	
LINKAGE GROUP I:						
A - flm-2	5	2	2	0	mt	
a - acr-3 +	1"	0	0	0	KH14(r)	
10 1"					Y256M223	
+ flm-2	43	1			66702	
suc +	6"	2			Y256M223	
8						
+ flm-2		2			T(I;IV)cut	
cut +					Y256M223	
3 (126 coral)						
+ aur arom-8	17	14	9	9	Y256M223	
flm-2 + +	12	1"	1"	9	34508	
47 41					DH8	
(B. c. Turner)						
A atr-1 +		3		0	mt	
a + un-2	22		5		RC2(r)	
10 17					46006(t)	
+ + bs		2	2		B123	
cr un-1 +	27			1	44409(t)	
9 9					AR62	
+ os-5 +	3"	4	2	0	KH52(r)	
act-1 + al-1	33	5	1	0	NM216o	
12 4					ALS4	
+ aur os-1	2"	2	6	0	T53M50(t)	
un-7 + +	24	0	7	0	34508	
3 22					B135	
+ al-2 arg-6		9	2		T53M50(t)	
un-7 + +					T5300	
Selective, un ⁺ arg ⁺ .						
Therefore, <u>un</u> not						
between.						
+ al-2		0			NM216o	
os-5 +		1			153"	
0.4 (206 total)						
+ aur	21	3			P5341	
os-5 +	26	0			34508	
6						

zygote genotype and % recombination	Paren- tals	Crossovers			Isolation Numbers
		1	7.	1,2	
LINKAGE GROUP 11:					
+ col-10	32	1			B101
pi +	48	0			R2438
1.2					
het-c + +			1		het-c
het-C pyr-4 r o - 3			1		36601
0.8					R2354
(236 total)					T(II-W)
Therefore ro-3 not between.					homozygous
+ ro-3 thr-2	52	2	4	0	36601
pyr-4 + +	64	0	6	0	R2354
2 8					35423
+ + da	32	7	2	0	36601
pyr-4 thr-3 +	4"	5	0	1	44104(t)
15 3					R2375
+ + lp		6	4		36601
pyr-4 thr-3 +	3"			0	44104(t)
15 10					P736
+ + m	26	1	3	0	36601
pyr-4 thr-3 +					44104(t)
3 10					NM218
+ thr-3 +	15	6	1	0	R2354
ro-3 thr-2 arg-5	18	1	4	0	T42M59(t)
16 17					35423
					27947
+ arg-5 mo	23	2	1"	0	35423
thr-2 + +	26	0	6	0	27947
3 24					D309
+ arg-5 mo	21	7	0	0	35423
thr-2 + +	27	4	1	0	27947
18 2					NM220
+ arg-5 pe	28	1	1	0	R2526
ro-9 + +	-	0	0	0	27947
3 3					Y8743m
+ mo +	31	0	4	0	UM107
arg-12 + fl	38	0	6	0	D309
0 13					1.

Table 2 (continued).

Zygote genotype and % recombination	Paren- tals	Crossovers			Isolation Numbers
		1	2	1,2	
LINKAGE GROUP 111:					
<u>+</u> <u>arg-9</u> <u>+</u>	32	0	3	0	9185
Lhi-2 <u>+</u> ro-2	40	0	3	0	35401
0 8					B20
<u>+</u> <u>un</u> <u>dow</u>	13	1	5	0	Y6994
tyr-1 <u>+</u> <u>+</u>	29	1	8	0	T51M171(t)
5 16					P616
<u>+</u> <u>+</u> Chi-4	13	17	0	0	R2537
spco-15 <u>acr-2</u> <u>+</u>					85902
> 50 0					KH5
LINKAGE GROUP IV:					
<u>+</u> <u>pyr-1</u>	23	0			T27M9(t)
un-8 <u>+</u>	24	0			H263
0					
<u>+</u> <u>pdx</u> <u>un-12</u>		12	27		T27M9(t)
un-8 <u>+</u> <u>+</u>					37803
Selective, un ⁺ .					T51M118(t)
Therefore <u>un-12</u>					
not between.					
<u>+</u> <u>col-4</u> <u>+</u>	35	2	0	0	37803
pdx <u>+</u> <u>un-12</u>	35	1	0	0	70007
4 0					T51M118(t)
LINKAGE GROUP V:					
<u>+</u> <u>at</u>	10	9			R2365
spco-3 <u>+</u>					M111
<u>+</u> <u>+</u> <u>+</u>	37	5	5	0	33933
lys-1 <u>aspt</u> <u>iv-1</u>	31	4	10	0	44304(C)
10 16					16117
<u>+</u> <u>act-2</u> <u>leu-5</u>	40	0	0	0	37101
lys-2 <u>+</u> <u>+</u>	45	0	0	0	KH53(r)
0 0					45208
Selective, act ^R leu ⁺ .	-		9		
Therefore <u>leu-5</u> not	-			0	
between.					
Selective, lys ⁺ leu ⁺ .	-	33	7		
Therefore <u>lys-2</u> not	-				
between.					

Zygote genotype and % recombination	Paren- tals	Crossovers			Isolation Numbers
		1	2	1,2	
<u>+</u> <u>leu-5</u> <u>+</u>	45	0	3	0	KH53(r)
act-2 + sp	44	1	3	0	45208
1 6					B132
<u>+</u> <u>al-3</u> <u>tryp-5</u>	29	20			83201(L)
inos + +					RP100
Selective, inos ⁺					A420
tryp ⁺ . Therefore					
<u>inos</u> not between.					
<u>+</u> <u>al-3</u>	43				89601
inos +		1			RP100
2					
<u>+</u> <u>al-3</u>	129				83201(t)
inos +		1			RP100
1					
<u>+</u> <u>+</u>	45	2			RP100
al-3 tryp-5	37	1			A420
4					
<u>+</u> <u>un-11</u> <u>+</u>	34	0	13	0	RP100
al-3 + hist-6					T42M30(t)
0 27					Y152M105
<u>+</u> <u>pyr-6</u> <u>+</u>	12		1	0	RP100
al-3 + Mat-6	15	2	1	0	DFC37
19 6					Y152M105
<u>+</u> <u>asp</u> <u>+</u>	19	6	1	0	RP100
al-3 + hist-6	20	5	2	0	T51M147
21 6					Y152M105
<u>+</u> <u>cot-2</u> <u>pab-2</u>	42	1	5	0	B6
bis + +	38	6	1	0	R1006
8 6					"193
<u>+</u> <u>un-9</u> <u>+</u>	31	0	2	0	DFC37
pyr-6 + hist-6	14	1	1	0	T54M96(t)
2 6					Y152M105
LINKAGE GROUP "I:					
<u>+</u> <u>un</u> <u>+</u>	19	0	0	0	DS6-85
lys-5 + ylo-1	24	0	1	0	T51M154(t)
0 2					Y30539y
<u>+</u> <u>ylo-1</u>	48	1			T42M24(t)
un-13 +	56	1			Y30539y
2					

Zygote genotype and % recombination	Parentals	Crossovers			Isolation Numbers
		1	2	1,2	
+ + <u>rib-1</u>	•	-	79		P5241
yl o-1 ad-1 +	•	-		0	3254
Selective, ad+ rib+.					51602(t)
Therefore <u>rib</u> not between.					
+ <u>rib-1</u> +	141	9	0	0	Y30539y
yl o-1 + pan-2	124	5	0	0	76R5
5 0					Y--?--
+ <u>rib-1</u> +	•	-		0	Y30539y
yl o-1 + <u>pan-2</u>	•	-	39		51602(t)
Selective, rib ⁺ pan ⁺ .					Y--?--
Therefore <u>pan</u> not between.					
+ + <u>acr-7</u>	34	3	12	2	Y30539y
yl o-1 <u>tryp-2</u> +	27	4	13	0	75001
9 28					P1676(r)
LINKAGE GROUP VII:					
+ + <u>un-10</u>	15	8	1	0	R2367
spco-4 <u>wc</u> +	34	5	4	0	P829
19 7					T42M45(t)
+ + +	29	11	4	1	Y31881
nic-3 <u>wc</u> <u>arg-10</u>	33	6	6	2	P829
22 14					B317
+ + +	5	2	3	0	Y31881
nic-3 <u>wc</u> <u>sk</u>	7	5	3	0	P829
28 24					B234

[illegible]