

Quantitative determination of perithecium formation

C. M. DeSha
Texas Women's University

R. Fuerst
Texas Women's University

Follow this and additional works at: <https://newprairiepress.org/fgr>



This work is licensed under a [Creative Commons Attribution-Share Alike 4.0 License](https://creativecommons.org/licenses/by-sa/4.0/).

Recommended Citation

DeSha, C. M., and R. Fuerst (1972) "Quantitative determination of perithecium formation," *Fungal Genetics Reports*: Vol. 19, Article 9. <https://doi.org/10.4148/1941-4765.1867>

This Technical Note is brought to you for free and open access by New Prairie Press. It has been accepted for inclusion in *Fungal Genetics Reports* by an authorized administrator of New Prairie Press. For more information, please contact cads@k-state.edu.

Quantitative determination of perithecium formation

Abstract

Quantitative determination of perithecium formation

DeSha, C.M. and R. Fuerst. A technique for the quantitative determination of perithecium formation.

filtered through 2.5 cm of sterile cotton in a disposable 10 ml syringe in order to remove hyphal fragments from the conidial suspension and to obtain microconidia. Sterile water is used to dilute this suspension until an absorbance reading of 0.05 is reached on a Bausch and Lomb Spectrophotometer 20, set at 500 nm. One ml of the suspension is diluted 1:100 with sterile water. Sterile 35mm Brewer plates with metal tops holding absorbent discs are filled with 15 ml of sterile CoLab's Oxoid cornmeal agar, which is made according to the specifications on the label with 1.5% agar. The Brewer plates are used since the discs absorb the water that condenses in the plate. Fewer perithecia are formed when other brands of cornmeal agar, such as Difco's or BBL's are employed. Westergaard's medium (Westergaard and Mitchell 1947 Am. J. Botany 34:573) can not be used since conidial growth is enhanced, thereby making it more difficult to count the perithecia.

After the plates are cool, 0.05 ml aliquots of the Em 5297a suspension, diluted 1:100 and containing less than 150 but more than 100 microconidia, are delivered to the centers of the plates, using a Schwartz BioResearch Autopipettor. The disposable tips for the pipettor are sterilized in a petri plate for 24 hr before use with several drops of ethylene oxide, since the tips cannot be autoclaved. The automatic pipettor is used since it allows fast, accurate dispensation of the conidial suspension onto the plates. Inoculated cornmeal plates are left at 25°C until protoperithecia develop, after which 0.05 ml of N. crassa St. Lawrence 74A inoculum (with the same microconidial concentration and obtained in the same manner as described for Em5297a and with an absorbance of 0.05, is pipetted onto the surface of the agar in the inoculated plate. This time the conidial suspension is spread over the surface of the agar with an autoclaved 5 x 7.5 cm glass slide. The slide is used for the spreading procedures, instead of a rod, since the inoculum can be applied faster and more evenly, according to our experience in the laboratory, by this method.

Dark perithecia form in seven days on CoLab's cornmeal agar. The spreading technique allows for even distribution on the surface of the agar, so that the perithecia can be more easily counted with the aid of a bacterial colony counter. Before counting, however, the plates are washed with 2% Clorox to inactivate the conidiospores which have formed. A control plate, one in which the maternal strain has not been treated, is found to contain approximately 250 perithecia.

The 1:100 dilution of Em5297a conidia used in this technique might be varied by the investigator depending on the perithecia producing capacity of the strains used. This method has been found in our laboratory to be applicable to several strains that were tested. It may be concluded that this method is excellent for measuring changes in the ability of the chemically or physically-treated strain to produce perithecia when mated to an untreated paternal strain, as compared to an untreated control cross.

This work was supported in part by Research Grant No. M-190 from the Robert A. Welch Foundation and from an Institutional Grant from the Texas Woman's University. ■ ■ Department of Biology, Texas Woman's University, Denton, Texas 76204.