

Effect of cycloheximide and actinomycin D on germinating conidia

S. E. Hitchcock

V. W. Cochrane

Follow this and additional works at: <http://newprairiepress.org/fgr>

Recommended Citation

Hitchcock, S. E., and V.W. Cochrane (1969) "Effect of cycloheximide and actinomycin D on germinating conidia," *Fungal Genetics Reports*: Vol. 15, Article 22. <https://doi.org/10.4148/1941-4765.1925>

This Response of Neurospora to Various Antibiotics and other Toxic Chemicals is brought to you for free and open access by New Prairie Press. It has been accepted for inclusion in Fungal Genetics Reports by an authorized administrator of New Prairie Press. For more information, please contact cads@k-state.edu.

Effect of cycloheximide and actinomycin D on germinating conidia

Abstract

Effect of cycloheximide and actinomycin D on germinating conidia

Creative Commons License



This work is licensed under a [Creative Commons Attribution-Share Alike 4.0 License](https://creativecommons.org/licenses/by-sa/4.0/).

Hitchcock, S.E. and V.W. Cochrane. Effect of cycloheximide and actinomycin on germinating conidia.

Conidia were harvested at 7 days, washed, and incubated in Vogel's liquid medium N with (called "germinating") or without (called "non-germinating") 2% glucose at a spore concentration of 10 mg/ml wet weight.

Cycloheximide (Upjohn Co.) at concentrations of 1, 10, and 100 $\mu\text{g/ml}$ (0.1, 1.0, 10.0 $\mu\text{g/mg}$ wet weight) inhibits germination completely and inhibits the incorporation of leucine- $\text{U-}^{14}\text{C}$, phenylalanine- $\text{U-}^{14}\text{C}$, and proline- $\text{U-}^{14}\text{C}$ into protein (hot TCA insoluble, hot NaOH soluble material). Inhibition at 1 $\mu\text{g/ml}$ was usually greater than 84%, at 10 $\mu\text{g/ml}$ greater than 97%, and at 100 $\mu\text{g/ml}$ greater than 99%. The inhibitor had complex effects on the amino acid pools which have not been analyzed.

Figures 1 and 2 show the effect of cycloheximide on RNA synthesis in "germinating" conidia. When cycloheximide was added at the beginning of the incubation, germination was inhibited and RNA synthesis approximated that in "non-germinating" conidia (Figure 1). When the inhibitor was added after germination had begun, RNA synthesis continued at the control rate for a short period and then leveled off and never attained the level of the control (Figure 2). Addition of cycloheximide at 30 minutes inhibited germination completely, while if added at 80 minutes, it halted germination after 20 minutes. It may be concluded that, while some RNA synthesis occurs in the absence of protein synthesis, continued protein synthesis is required for RNA synthesis at the rate found in germinating conidia.

In a study of the germination of conidia of wild type strain Em 5297a (ATCC#10816), the synthetic capacities of conidia incubated in minimal medium with and without a carbon source were investigated. Conidia were grown on Vogel's medium N with 1% glucose and 2% agar.




Figure 1. The effect of cycloheximide on RNA synthesis. Incorporation of uracil-2-C¹⁴ (5 μ c/flask, 0.075 mM) into TCA insoluble material. G="germinating"; U="non-germinating"; CYCLO=cycloheximide, 100 μ g/ml; % G=percent germination of "germinating" conidia.




Figure 2. The effect of cycloheximide on RNA synthesis when added after the beginning of incubation. Cycloheximide was added to germinating conidia at 30 or 80 minutes. Experimental conditions and symbols are as described for Figure 1.

Table 1. Effect of actinomycin D on RNA synthesis and germination.

Actinomycin D (μ g/ml)	0	1	10	25	100	250
% germination at 2.5 hr.	40	40	38	40	--	23
CPM in TCA insoluble material at 0.5 hrs.	2,529	2,417	2,155	2,182	2,208	1,613
CPM in TCA insoluble material at 1.5 hrs.	32,092	32,056	32,498	28,053	28,549	16,630

Table 1 shows preliminary results of the effect of actinomycin D (gift of Merck, Sharpe and Dohme) on RNA synthesis and germination. Concentrations of up to 100 μ g/ml had a small, if any, effect on RNA synthesis and germination. An extremely high level, 250 μ g/ml, inhibited incorporation of uracil-2-C¹⁴ into TCA insoluble material only by 48.2% and germination by 57.5%. It was not determined if the actinomycin D was taken up by conidia.

This research was supported by NSF grant GE 1401. Department of Biology, Wesleyan University, Middletown, Connecticut 06457. Present address of SEH: Department of Biology, Case Western Reserve University, Cleveland, Ohio 44106.