Tyrosine transport in Neurospora

V. T. Brescia

Follow this and additional works at: <https://newprairiepress.org/fgr>

This work is licensed under a [Creative Commons Attribution-Share Alike 4.0 License](https://creativecommons.org/licenses/by-sa/4.0/).

Recommended Citation

Brescia, V. T. (1968) "Tyrosine transport in Neurospora," *Fungal Genetics Reports*: Vol. 13, Article 1.
<https://doi.org/10.4148/1941-4765.1934>

This Research Note is brought to you for free and open access by New Prairie Press. It has been accepted for inclusion in Fungal Genetics Reports by an authorized administrator of New Prairie Press. For more information, please contact cads@k-state.edu.

Tyrosine transport in Neurospora

Abstract

Tyrosine transport in Neurospora

Brescia, V. T. Tymsine transport in *Neurospora*.

Uptake of tyrosine by *Neurospora* conidia was studied using ^{14}C tyrosine in the manner described by DeBusk and DeBusk (1965 *Biochim. Biophys. Acta* 104: 139) for phenylalanine. Conidial suspensions

which consistently gave 0.19 - 0.28 mg dry weight of conidia per 5 ml sample were prepared by adjusting OD₃₉₇ to 0.9-0.95 (B and L Spectronic 20). The usual conditions were a temperature of 30°C and tyrosine concentration of 1 μmole per 25 ml ($4 \times 10^{-5} \text{ M}$) reaction mixture (Vogel's minimal + cells). The optimum temperature was later found to be between 31-33°C and the pH optimum 5.8.

Incubation at 45°C for 20 minutes did not inactivate the transport system - as little as 2 minutes at 50° did temporarily inactivate (uptake less than 70% of control at 20 minutes). Recovery occurred in cells held at 30° for 30 minutes following 50° heat inactivation. Concentrations from 0.2 $\mu\text{mole}/25\text{ml}$ to 3.2 $\mu\text{mole}/25 \text{ ml}$ gave increasing initial rates of uptake; no increase was observed above 5 $\mu\text{mole}/25 \text{ ml}$. A reciprocal plot of initial uptake vs tymsine concentration (Lineweaver-Burke) gave a straight line. By extrapolation, the K_m was estimated at 1.2-1.8 $\times 10^{-4} \text{ M}$ in three experiments. After 50 minutes uptake, the amount of label chromatographically identical with tyrosine that can be extracted with 5% TCA at room temperature in 10 minutes is at least 30 x the external concentration.

Glucose (final conc. 1%) added to an actively transporting system will inhibit further transport within 6 minutes and will continue inhibiting for at least 15 minutes, after which transport is resumed, apparently at the same rate. Sodium azide and 2,4-dinitrophenol at 10^{-3} M restrict transport to about 10% of the control. With azide, at least the inhibition is almost instantaneous. A variety of compounds were tested at concentrations 25 x that of tymsine for their effects on uptake of ^{14}C L-tyrosine at a concentration of $4 \times 10^{-5} \text{ M}$. Shikimic acid and para-hydroxyphenylpyruvate, among others, had no effect whereas L-tryptophan and L-phenylalanine reduced uptake to 20% or less of control. Since all of the &we-mentioned compounds can supplement appropriate mutants, they must be capable of entering the cell. Therefore, the lack of effect of shikimic acid and para-hydroxyphenylpyruvate must reflect a stereospecificity of the tyrosine transport system. This is further demonstrated by the fact that D-tyrosine reduces uptake to 87% of control, whereas an equivalent amount of ^{12}C L-tyrosine reduces it to 25% of control. ■ ■ ■ Department of Biological Science, Florida State University, Tallahassee, Florida 32306.