

Linkage information for cysteine and methionine mutants

N. E. Murray

Follow this and additional works at: <http://newprairiepress.org/fgr>

Recommended Citation

Murray, N. E. (1968) "Linkage information for cysteine and methionine mutants," *Fungal Genetics Reports*: Vol. 13, Article 5.
<https://doi.org/10.4148/1941-4765.1938>

This Linkage Data is brought to you for free and open access by New Prairie Press. It has been accepted for inclusion in Fungal Genetics Reports by an authorized administrator of New Prairie Press. For more information, please contact cads@k-state.edu.

Linkage information for cysteine and methionine mutants

Abstract

Linkage information for cysteine and methionine mutants

Creative Commons License



This work is licensed under a [Creative Commons Attribution-Share Alike 4.0 License](https://creativecommons.org/licenses/by-sa/4.0/).

Murray, N. E. Linkage information for cysteine and methionine mutants.

cys-9 (T156). cys-9 is located between cr (crisp) and thi-1 (thiamine-1) in lininkaa group IR (see Table 1).

cys-10 (39816). The tentative location of this locus (Murray 1965 Genetics 52: 801) as the most distal marker in the left arm of linkage group IV is supported by information from other workers.

cys-11 (NM86). A cluster of cysteine mutants is located in the cys-5 region between leu-3 and mating type. The evidence is consistent with the region comprising two loci, cys-5 and cys-11. The mutants NM44 and R83R 1-1-271 gave very low recombination frequencies when crossed to cys-5 (35001). The recombinants from the latter cross were scored for flanking markers and all four flanking marker combinations were represented. When a fourth mutant (NM86) was crossed to cys-5 (35001) the recombination frequency was much higher and there was no or little "negative interference". Complementation tests showed that NM86 is physiologically different from the heterocaryon compatible cys-5 (NM44) strain, and more specifically Lein-weber (personal communication) has shown that while NM86 lacks ATP-sulfurylase, the cys-5 alleles tested (35001 and NM44) lack PAPS-reductase. It is proposed that NM86 is an allele at locus cys-11. The combination of flanking markers found for cysteine independent recombinants from a cross of cys-5 by cys-11 indicate the order mating type, cys-11, cys-5, leu-3. Adequate genetic information is lacking for a cross of cys (85518) by cys-5, but cys (85518) gave a very low recombination frequency (1 in 200,000) when crossed to cys-11 (NM86).

cys-12 (NM268). cys-12 is an additional cysteine locus in linkage group I distal to ad-9 and close to al (0 recombinants among 76 isolates) (see Table 1).

me-6 (35809) and mac (65108). These mutants are closely linked. Methionine independent recombinants have been isolated from crosses of me-6 by mac and classified with respect to the flanking markers thi-1 and ad-9 (adenine-9). The order indicated is thi-1, mac, me-6, ad-9, but it is probable that mac and me-6 is allelic.

me-7 and me-9. Methionine independent recombinants have been isolated from crosses of me-7 (NM73) by me-9 (NM43t) and classified with respect to flanking markers (thiamine-3 and white collar). The methionine loci are very closely linked in the order thi-3, me-7, me-9, wc.

Table 1. Linkage data on random segregants from crosses involving cys-9 or cys-12.

Zygote genotype and % recombination	Parental combinations	Recombination			Total and % germination	Marker isolation numbers
		Singles region 1	Singles region 2	Doubles regions 1 and 2		
<u>+</u> <u>thi-1</u> <u>ad-9</u>	34	7	10	0	93	T156
<u>cys-9</u> <u>+</u> <u>+</u>	33	5	4	0	(93%)	56501
12.9 15.1						Y 154M37
<u>+</u> <u>cys-9</u> <u>+</u>	26	2	20	0	88	8122
<u>cr</u> <u>+</u> <u>os-1</u>	23	1	16	0	(63%)	T156
3.4 40.9						B135
<u>+</u> <u>+</u> <u>cys-12</u>	48	8	8	0	120	56501
<u>thi-1</u> <u>ad-9</u> <u>+</u>	44	5	6	1	(83%)	Y 154M37
11.7 12.5						NM268

(The top number in each p-air represents the class that has the + allele of the leftmost marker).