

A method for detection of suppressor genes closely linked to the suppressed locus

M. Allison

Follow this and additional works at: <http://newprairiepress.org/fgr>

Recommended Citation

Allison, M. (1967) "A method for detection of suppressor genes closely linked to the suppressed locus," *Fungal Genetics Reports*: Vol. 12, Article 15. <https://doi.org/10.4148/1941-4765.1960>

This Technical Note is brought to you for free and open access by New Prairie Press. It has been accepted for inclusion in Fungal Genetics Reports by an authorized administrator of New Prairie Press. For more information, please contact cads@k-state.edu.

A method for detection of suppressor genes closely linked to the suppressed locus

Abstract

Detection of closely linked suppressor genes

Creative Commons License



This work is licensed under a [Creative Commons Attribution-Share Alike 4.0 License](https://creativecommons.org/licenses/by-sa/4.0/).

Allison, M. A method for detection of suppressor genes closely linked to the **suppressed** locus.

Diepoxybutane-induced reversions to adenine independence at the **odeninc-34** locus (allele 38701) of *N. crassa* were examined for closely linked suppressor mutations. The scheme used screened against parental type ascospores, allowing only crossovers in a small region surrounding the **ad-3A** locus to germinate in the selective medium.

The revertant was buck-crossed to an **od-3A (38701)**, **nic-2 (43002)** strain and homocaryotic **ad-3A** revertant, **nit-2** progeny were isolated. The **nis-2** locus, approximately 3 map units distal to **ad-3A**, provided one closely-m marker. The adenine-independent, **nit-2** strain was then crossed to a strain carrying the mutant allele **hist-2 (Y152M14)** which is approximately 2 map units proximal to the **ad-3A** locus. The progeny were plated on a medium supplemented with adenine only. Consequently the only survivors were the products of crossovers between the **hist-2** and **nit-2** loci.

+	ad-3A revertant?	nit-2
<hr/>		
hist-2	+	+

Around 10,000 spores were tested from one such cross. 27 of the 259 crossovers recovered proved to be adenine-requiring. One adenine-requirer isolated from this revertant showed the very strong mutational response to diepoxybutane which is specific for **adenine-3A (38701)**. The results suggest that there is a suppressor gene located approximately 0.54 map units distal to the **ad-3A** locus.

In 1955, Kølmark and Giles found no evidence for suppressors of **ad-3A (38701)**. Since, however, they examined only about 100 ascospores from each revertant, they could not exclude closely linked suppressors. - - - Institute of Animal Genetics, University of Edinburgh, West Mains Road, Edinburgh 9, Scotland.