

Kansas Agricultural Experiment Station Research Reports

Volume 0
Issue 1 *Cattleman's Day (1993-2014)*

Article 561

1996

Pregnancy rates in heifers and suckled beef cows after synchronized ovulation using PGF², GnRH, and norgestomet

D.P. Hoffman

Jeffrey S. Stevenson

C.L. Krehbiel

See next page for additional authors

Follow this and additional works at: <https://newprairiepress.org/kaesrr>



Part of the [Other Animal Sciences Commons](#)

Recommended Citation

Hoffman, D.P.; Stevenson, Jeffrey S.; Krehbiel, C.L.; Nichols, David A.; and McKee, R.M. (1996) "Pregnancy rates in heifers and suckled beef cows after synchronized ovulation using PGF², GnRH, and norgestomet," *Kansas Agricultural Experiment Station Research Reports*: Vol. 0: Iss. 1. <https://doi.org/10.4148/2378-5977.1964>

This report is brought to you for free and open access by New Prairie Press. It has been accepted for inclusion in Kansas Agricultural Experiment Station Research Reports by an authorized administrator of New Prairie Press. Copyright 1996 Kansas State University Agricultural Experiment Station and Cooperative Extension Service. Contents of this publication may be freely reproduced for educational purposes. All other rights reserved. Brand names appearing in this publication are for product identification purposes only. No endorsement is intended, nor is criticism implied of similar products not mentioned. K-State Research and Extension is an equal opportunity provider and employer.



Pregnancy rates in heifers and suckled beef cows after synchronized ovulation using PGF2", GnRH, and norgestomet

Abstract

Suckled cows and virgin heifers received a novel treatment that included PGF 2", GnRH, and norgestomet, with the objective of inducing estrus in prepubertal heifers and anestrous suckled cows, as well as synchronizing ovulation in estrus-cycling females. The treatment consisted of two injections of PGF 2" (day 14 and 0) plus 100 Fg of GnRH and a 6-mg norgestomet ear implant on day 7. The implant was removed 24 h after the second injection of PGF2" (day 0), and a second injection of GnRH was given 30 hours after implant removal. The treated females were inseminated 18 hours after the second injection of GnRH or at estrus, if it was detected before the second GnRH injection. Pregnancy rate in the treated females was greater than in control females that had received two injections of PGF 2" 14 days apart and were inseminated at estrus or at one fixed time (60.2 vs. 48%). The treatment successfully induced a fertile ovulation in previously prepubertal heifers and anestrous cows, resulting in 63.5% pregnancies vs. 26.5% for controls. In addition, in females not showing estrus, the treatment increased pregnancy rate following a fixed-time insemination (treatment vs. control; 60.0 vs. 3.8%). We concluded that treatment with PGF2", GnRH, and norgestomet induced estrus and increased pregnancy rates in prepubertal heifers, anestrous cows, and cycling females.

Keywords

Cattlemen's Day, 1996; Kansas Agricultural Experiment Station contribution; no. 96-334-S; Report of progress (Kansas State University. Agricultural Experiment Station and Cooperative Extension Service); 756; Beef; GnRH; Norgestomet; PGF2; Heat synchronization; Prepubertal heifers; Anestrous suckled cows

Creative Commons License



This work is licensed under a [Creative Commons Attribution 4.0 License](https://creativecommons.org/licenses/by/4.0/).

Authors

D.P. Hoffman, Jeffrey S. Stevenson, C.L. Krehbiel, David A. Nichols, and R.M. McKee

**PREGNANCY RATES IN HEIFERS AND SUCKLED
BEEF COWS AFTER SYNCHRONIZED OVULATION
USING PGF_{2α}, GnRH, AND NORGESTOMET**

*D. P. Hoffman, J. S. Stevenson, C. L. Krehbiel,
D. A. Nichols, and R. M. McKee*

Summary

Suckled cows and virgin heifers received a novel treatment that included PGF_{2α}, GnRH, and norgestomet, with the objective of inducing estrus in prepubertal heifers and anestrus suckled cows, as well as synchronizing ovulation in estrus-cycling females. The treatment consisted of two injections of PGF_{2α} (day 14 and 0) plus 100 μg of GnRH and a 6-mg norgestomet ear implant on day 7. The implant was removed 24 h after the second injection of PGF_{2α} (day 0), and a second injection of GnRH was given 30 hours after implant removal. The treated females were inseminated 18 hours after the second injection of GnRH or at estrus, if it was detected before the second GnRH injection. Pregnancy rate in the treated females was greater than in control females that had received two injections of PGF_{2α} 14 days apart and were inseminated at estrus or at one fixed time (60.2 vs. 48%). The treatment successfully induced a fertile ovulation in previously prepubertal heifers and anestrus cows, resulting in 63.5% pregnancies vs. 26.5% for controls. In addition, in females not showing estrus, the treatment increased pregnancy rate following a fixed-time insemination (treatment vs. control; 60.0 vs. 3.8%). We concluded that treatment with PGF_{2α}, GnRH, and norgestomet induced estrus and increased pregnancy rates in prepubertal heifers, anestrus cows, and cycling females.

(Key Words: GnRH, Norgestomet, PGF_{2α}, Heat Synchronization, Prepubertal Heifers, Anestrus Suckled Cows.)

Introduction

Estrus-synchronization programs improve reproductive efficiency by reducing the length of breeding and calving seasons, increasing calf weaning weights, and grouping cows and heifers so artificial insemination (AI) can be used more efficiently. They are not designed to induce estrus in prepubertal heifers or anestrus suckled beef cows. Treatments involving single or multiple injections of gonadotropin-releasing hormone (GnRH) given 10 to 12 days apart and/or implants of norgestomet have been used to jump start (induce estrus) noncycling heifers and cows. The result of injecting GnRH is to induce secretion of LH and FSH and cause ovulation of mature follicles. Norgestomet primes the hypothalamic-pituitary axis for release of the endogenous GnRH, LH, and FSH necessary for follicle growth. In prepubertal heifers and anestrus suckled cows, the norgestomet implant also prevents the short luteal phase or short estrous cycle that normally follows the first pubertal or postpartum ovulation. That short cycle prevents the continuation of pregnancy, even when fertilization occurs. Therefore, our objective was to test the effect of this novel treatment using prostaglandin F_{2α} (PGF_{2α}), GnRH, and norgestomet for its ability to induce estrus and increase conception in prepubertal heifers and anestrus suckled cows, as well to synchronize estrus in cycling females before one fixed-time insemination.

Experimental Procedures

In a 2-year study, purebred Angus, Hereford, and Simmental heifers and suckled cows were assigned to two treatments: 1) two injections of PGF_{2α} 14 days apart (control); or 2) two injections of PGF_{2α} (days 14 and 0) plus 100 μg of GnRH and a 6-mg norgestomet

implant on day 7 (ovulation synchronization; Figure 1). The norgestomet implant was removed 24 h after the second injection of PGF_{2α} (day 0). A second injection of 100 μg of GnRH was given 30 h after implant removal. Three blood samples were collected (24, 14, and 7 days) before the first GnRH injection to determine estrus-cycling status.

Control females were inseminated 12 to 16 h (AM-PM rule) after first detected estrus until 80 h after the second PGF_{2α} injection, when all remaining females were inseminated. The females in the ovulation synchronization treatment were inseminated either at estrus or at 18 h after the second injection of GnRH (48 h after implant removal or 72 h after the second PGF_{2α} injection). Pregnancy status was determined by intrarectal ultrasonography on days 34 or 35 after insemination.

Results and Discussion


The majority of heifers (87%) and cows (66%) were cycling at the time the treatments were initiated, based on serum progesterone concentrations 24, 14, and 7 days before the time of the GnRH injection and implant (Figure 1). Similar proportions of noncycling control and ovulation-synchronized heifers showed estrus and were inseminated at estrus (46.9 vs. 34.6%). For noncycling cows, more (P<.05) control than ovulation-synchronized cows were inseminated at estrus (80 vs. 39.5%). The remaining heifers and cows were inseminated at a fixed time after PGF_{2α}.

No differences in pregnancy rates were detected among breeds or parity groups (heifers and cows). Pregnancy rate was greater (P<.01) in the ovulation-synchronized females than in controls (60.2 vs. 48%). Control and treated females that were already cycling had similar pregnancy rates (56.2 vs. 58.8%). The experimental treatment increased pregnancy rates in noncycling females (26.5 vs. 63.5%; Table 1). Our results indicate that the ovulation synchronization treatment successfully induced a fertile ovulation in previously prepubertal heifers and anestrous suckled cows.

Pregnancy rates were similar in ovulation-synchronized females (66.1 vs. 60.6%), regardless of whether inseminated at estrus or at one fixed time (72 hours after PGF_{2α}; Table 2). In contrast, controls inseminated at estrus had greater pregnancy rates than controls inseminated at one fixed time (80 hours after PGF_{2α}), 60 vs. 3.8%.

These results demonstrate that our novel ovulation synchronization treatment induced a fertile ovulation in both noncycling and cycling heifers and cows. Furthermore, conception rate by fixed timed insemination after the ovulation synchronization equalled that achieved when inseminations were made at estrus in controls. Therefore, our treatment synchronizes ovulation with estrus. We conclude that treatment with PGF_{2α}, GnRH, and norgestomet induced estrus and increased pregnancy rates in prepubertal heifers, anestrous cows, and cycling females. In addition, the treatment increased pregnancy rates following one fixed-time insemination.

Ovulation Synchronization



A = 25 mg of Lutalysefi (PGF_{2α})

B = 100 μg of Cystorelinfi (GnRH)+ 6-mg ear implant of norgestomet () ■


C = 25 mg of Lutalyse

D = Removal of implant

E = 100 μg of Cystorelin

F = Insemination

Control



A = 25 mg of Lutalyse

B = 25 mg of Lutalyse

C = Insemination (in absence of detected estrus)

Figure 1. Treatment Protocols for Ovulation Synchronization Treatment and Control

Table 1. Pregnancy Rates in Previously Noncycling and Estrus-Cycling Heifers and Suckled Cows after Synchronized Ovulation using PGF_{2α}, GnRH, and Norgestomet^a

Parity	Treatment ^b			
	Control		Ovulation Synchronized	
	No.	% Pregnant	No.	% Pregnant
Noncycling	49	26.5 ^x	52	63.5 ^y
Heifers	4	0.0	5	100.0
Cows	45	28.9	47	59.6
Cycling	130	56.2 ^x	119	58.8 ^x
Heifers	32	56.3	32	59.4
Cows	98	56.1	87	58.6

^aHeifers and suckled cows were classified as noncycling or estrus-cycling based on serum concentrations of progesterone measured in three samples (24, 14, and 7 days before the second injection of PGF_{2α}).

^bSee Figure 1 for details.

^{x,y}Interaction (P=.08) of treatment and estrus-cycling status.

Table 2. Effect of Fixed-Time Insemination on Pregnancy Rates in Heifers and Suckled Cows

AI Time ^b	Treatment ^a			
	Control		Ovulation Synchronized	
	No.	% Pregnant	No.	% Pregnant
Estrus	127	66.1 ^x	66	60.6 ^x
Fixed time	52	3.8 ^y	91	60.0 ^x
Total	179	48.0	171	60.2

^aSee Figure 1 for details.

^bControls were inseminated at one fixed time (80 hours after the second injection of PGF_{2α}) in the absence of estrus. Ovulation synchronized females were inseminated 18 hours after the second GnRH injection (48 hours after the norgestomet implant was removed or 72 hours after the second injection of PGF_{2α}) in the absence of estrus.

^{x,y}Interaction (P<.01) of treatment and insemination time.