

Nature of a leucine-requiring strain derived from a phen-I stock

K.K. Jha

Follow this and additional works at: <http://newprairiepress.org/fgr>

Recommended Citation

Jha, K. K. (1967) "Nature of a leucine-requiring strain derived from a phen-I stock," *Fungal Genetics Reports*: Vol. 11, Article 3.
<https://doi.org/10.4148/1941-4765.1968>

This Research Note is brought to you for free and open access by New Prairie Press. It has been accepted for inclusion in Fungal Genetics Reports by an authorized administrator of New Prairie Press. For more information, please contact cads@k-state.edu.

Nature of a leucine-requiring strain derived from a *phen-1* stock

Abstract

Leucine requirer derived from *phen-1* stock

Creative Commons License



This work is licensed under a [Creative Commons Attribution-Share Alike 4.0 License](https://creativecommons.org/licenses/by-sa/4.0/).

Jho K. K. Nature of a leucine-requiring strain derived from a phen-1 stock.

Previously it was reported (Jha 1965 NN#7:16) that crosses between two phen-1 strains, H6196 (FGSC#492) and UA119 (FGSC#1167), yielded isolates which, unlike the phen-1 parents, had an obligate requirement for leucine. These derivatives were of mating type a and were presumed to originate from the H6196 stock. One of these "leucine" derivatives has been tested and confirmed to be a Leucine-3 isolate. It appears that the H6196 stock used by the author carries a very small proportion of leu-3 nuclei in heterocaryotic association but the ratio of leu-3 nuclei to phen-1 nuclei is not sufficiently high to give complementation. These derivatives can be neglected while considering other modifications of the phen-1 phenotype. - - - Department of Genetics, John Curtin School of Medical Research, Australian National University, Canberra, A.C.T., Australia.