

Kansas Agricultural Experiment Station Research Reports

Volume 0

Issue 1 *Cattleman's Day* (1993-2014)

Article 610

1996

Amino acid supplementation to growing and finishing steers

C.G. Campbell

C.T. Milton

Evan C. Titgemeyer

Follow this and additional works at: <https://newprairiepress.org/kaesrr>

 Part of the [Other Animal Sciences Commons](#)

Recommended Citation

Campbell, C.G.; Milton, C.T.; and Titgemeyer, Evan C. (1996) "Amino acid supplementation to growing and finishing steers," *Kansas Agricultural Experiment Station Research Reports*: Vol. 0: Iss. 1. <https://doi.org/10.4148/2378-5977.2013>

This report is brought to you for free and open access by New Prairie Press. It has been accepted for inclusion in Kansas Agricultural Experiment Station Research Reports by an authorized administrator of New Prairie Press. Copyright 1996 Kansas State University Agricultural Experiment Station and Cooperative Extension Service. Contents of this publication may be freely reproduced for educational purposes. All other rights reserved. Brand names appearing in this publication are for product identification purposes only. No endorsement is intended, nor is criticism implied of similar products not mentioned. K-State Research and Extension is an equal opportunity provider and employer.



Amino acid supplementation to growing and finishing steers

Abstract

One hundred British and British cross steers, averaging 631 lb (initial wt) were used in a growing and finishing study to evaluate the effects of unprotected amino acid supplementation on cattle performance and carcass characteristics. All diets contained 1% of a nonprotein nitrogen source, and treatments were: no additional supplemental protein (UREA), 2) supplemental protein from soybean meal (SBM), 3) 13 grams/day of an amino acid supplement (Low AA), and 4) 26 grams/day of an amino acid supplement (High AA). The Low AA treatment supplied 2 grams methionine, 8 grams lysine, 2 grams threonine, and 1 gram tryptophan per day, whereas the High AA treatment provided twice those amounts. The grower diet was based on whole-plant sorghum silage, and the finishing diet was based on rolled corn and corn silage. During the growing period, gains were higher ($P < .05$) for SBM-supplemented steers than for UREA steers and intermediate for amino acid-supplemented steers. Intakes were higher for steers supplemented with Low AA than for those supplemented with UREA or High AA. Few significant differences among treatments were observed in cattle performance during the finishing period. Hot carcass weights, dressing percentage, KPH fat, and yield grade were unaffected by amino acid supplementation. In this study, supplementing growing and finishing cattle with unprotected amino acids did not significantly improve steer performance or carcass characteristics, suggesting either that these amino acids were not limiting in these steers or that not enough of these supplemented amino acids escaped ruminal degradation to affect steers' performance.

Keywords

Cattlemen's Day, 1996; Kansas Agricultural Experiment Station contribution; no. 96-334-S; Report of progress (Kansas State University. Agricultural Experiment Station and Cooperative Extension Service); 756; Beef; Amino acids; Steers; Feedlot; Performance

Creative Commons License

This work is licensed under a [Creative Commons Attribution 4.0 License](https://creativecommons.org/licenses/by/4.0/).

AMINO ACID SUPPLEMENTATION TO GROWING AND FINISHING STEERS

C. G. Campbell, E. C. Titgemeyer, and C. T. Milton

Summary

One hundred British and British cross steers, averaging 631 lb (initial wt) were used in a growing and finishing study to evaluate the effects of unprotected amino acid supplementation on cattle performance and carcass characteristics. All diets contained 1% of a nonprotein nitrogen source, and treatments were: no additional supplemental protein (UREA), 2) supplemental protein from soybean meal (SBM), 3) 13 grams/day of an amino acid supplement (Low AA), and 4) 26 grams/day of an amino acid supplement (High AA). The Low AA treatment supplied 2 grams methionine, 8 grams lysine, 2 grams threonine, and 1 gram tryptophan per day, whereas the High AA treatment provided twice those amounts. The grower diet was based on whole-plant sorghum silage, and the finishing diet was based on rolled corn and corn silage. During the growing period, gains were higher ($P < .05$) for SBM-supplemented steers than for UREA steers and intermediate for amino acid-supplemented steers. Intakes were higher for steers supplemented with Low AA than for those supplemented with UREA or High AA. Few significant differences among treatments were observed in cattle performance during the finishing period. Hot carcass weights, dressing percentage, KPH fat, and yield grade were unaffected by amino acid supplementation. In this study, supplementing growing and finishing cattle with unprotected amino acids did not significantly improve steer performance or carcass characteristics, suggesting either that these amino acids were not limiting in these steers or that not enough of these supplemented amino acids escaped ruminal degradation to affect steers' performance.

(Key Words: Amino Acids, Steers, Feedlot, Performance.)

Introduction

The type of cattle fed in feedlots has changed considerably in recent years. Improvements in cattle nutrition, management, and genetics, along with newer feed additives and hormonal implants have resulted in average daily gains in the feedlot that can exceed 4 lb. Further, the composition of gain has shifted from fat to protein, and this has led some researchers to suggest that certain amino acids may be limiting performance. Supplementing protected amino acids to growing cattle in commercial situations is often cost prohibitive. However, if similar performance could be obtained by feeding higher levels of unprotected amino acids, then amino acid supplementation to growing and finishing cattle might be economical. Additionally, those supplemental amino acids might supply the rumen ecosystem with an essential nitrogen source, thereby enhancing ruminal fermentation.

Experimental Procedure

One hundred British and British cross steers averaging 631 lb initial weight were used in a randomized block design. Steers were allotted to one of five blocks based on initial weight and stratified into one of four pens within each block (five steers/pen). All growing diets contained 1% urea. Treatments were: 1) no supplemental protein (UREA), 2) 4.7% soybean meal (SBM), 3) 13 grams/day of an amino acid supplement (Low AA), and 4) 26 grams/day of an amino acid supplement (High AA). All finishing diets contained .8% urea and .2% ammonium sulfate, and treatments were the same as in the growing phase. The Low AA

treatment supplied 2 grams methionine, 8 grams lysine, 2 grams threonine, and 1 gram tryptophan per day, whereas the High AA treatment supplied 4, 16, 4, and 2 grams/steer/day of methionine, lysine, threonine, and tryptophan, respectively. It has been suggested that these amino acids most limit growth in cattle. The levels used were based on estimates of the amount of each amino acid that would be required to meet the steers' supplemental amino acid requirement, assuming 25% escaped ruminal destruction. Steers remained on the same treatment throughout the trial. For the growing phase, the diet was based on whole-plant sorghum silage (Table 1) and was fed for 85 days prior to a 13-day step-up to a finishing diet based on rolled corn and corn silage (Table 1). Steers remained on the finishing diet for 89 days prior to slaughter. Steers were weighed on 2 consecutive days at the initiation and end of the growing and finishing periods. During the step-up period, steers were moved into one of four dirt lots and remained on their respective treatments. All steers were implanted with Synovex-S at the initiation of the growing period and reimplanted with Revalor-S at the initiation of the finishing phase.

Results and Discussion

During the growing period, dry matter intakes were higher ($P < .05$) for steers fed SBM than for steers fed the UREA diet and also tended to be higher for steers fed the Low AA diet (Table 2). During the finishing period, however, intakes were similar across treatments. The higher intakes during the growing period and the higher intakes observed during the step-up period for SBM fed steers resulted in higher intakes for SBM fed steers for the whole study. For the whole study, intakes were higher for steers fed the Low AA diet than for steers fed the UREA or High AA diets. During the growing

phase, average daily gains were higher ($P < .05$) for SBM-supplemented steers than for UREA steers. Gains were intermediate for amino acid-supplemented steers. During the finishing period and the total study, no significant differences were observed among nitrogen sources for daily gains. However, for the total study, daily gains were numerically (but not statistically) higher for SBM-supplemented steers than for steers fed UREA or supplemented with amino acids.

During the growing phase, feed to gain conversions were numerically better for SBM-fed cattle than UREA-fed steers. However, during the step-up and the finishing periods, the UREA-fed steers had numerically improved efficiencies relative to SBM-supplemented steers, which probably represents a compensation for poorer gains during the growing period. For the whole trial, conversion efficiency was poorer for steers on the Low AA diet than for steers fed the UREA or High AA diets. Marbling score and 12th rib back fat were higher for SBM-supplemented steers than for steers supplemented with only UREA ($P < .10$). Marbling score was higher for steers supplemented with the Low AA treatment than for steers fed UREA or High AA. Quality grade was poorer for UREA-supplemented steers than for SBM-supplemented steers. Quality grade was similar for amino acid-supplemented steers and SBM-supplemented steers. No differences were observed between treatments for hot carcass weight; dressing percentage; or percents kidney, pelvic, and heart fat.

The lack of significant response to amino acid supplementation in this study suggests that either the supplemented amino acids were not limiting growth in these cattle or that not enough of these amino acids escaped ruminal degradation to alter cattle performance. If these amino acids stimulated ruminal fermentation, their effect on steer performance was not apparent.

Table 1. Composition of Diets Fed to Steers (% of Diet DM)

Item	Treatment			
	UREA	Low AA	High AA	SBM
<u>Grower period</u>				
Whole-plant sorghum silage	83.5	83.5	83.4	78.9
Rolled milo	9.3	9.2	9.1	9.2
Soybean meal				4.7
Molasses	3.0	3.0	3.0	3.0
Vitamin and mineral mix ^a	2.2	2.2	2.2	2.2
Urea	1.0	1.0	1.0	1.0
Rumensin and Tylan premix ^b	1.0	1.0	1.0	1.0
Amino acid mixture		.1	.3	
Crude protein	10.8	10.8	10.9	12.6
<u>Finishing period</u>				
Dry-rolled corn	79.7	79.7	79.6	75.5
Corn silage	10.1	10.1	10.1	10.1
Soybean meal				4.2
Ground sorghum	3.6	3.5	3.4	3.7
Vitamin and mineral mix ^a	2.5	2.5	2.5	2.4
Molasses	2.2	2.2	2.2	2.2
Rumensin and Tylan premix ^b	.9	.9	.9	.9
Urea	.8	.8	.8	.8
Ammonium sulfate	.2	.2	.2	.2
Amino acid mixture		.1	.3	
Crude protein	11.2	11.3	11.4	13.1

^aTo supply complete diets containing .8% Ca and .4% P.

^bTo supply 275 mg Rumensin and 90 mg Tylan/steer/day.

Table 2. Effects of Unprotected Amino Acids on Growing and Finishing Steer Performance and Carcass Characteristics

Item	Treatment				SEM
	UREA	Low AA	High AA	SBM	
<u>Grower period (85days)</u>					
Beginning wt, lb	632	631	630	630	1.1
Ending wt, lb ^a	834	846	840	860	9.2
Dry matter intake, lb/d ^{bc}	20.2	21.5	20.8	21.8	.41
Gain, lb/d ^c	2.37	2.53	2.47	2.70	.10
Feed:gain	8.51	8.55	8.43	8.15	.30
<u>Step-up period (13 days)^d</u>					
Dry matter intake, lb/d	22.9	23.0	22.5	25.5	
Gain, lb/d	3.09	2.78	2.54	3.04	
Feed:gain	7.41	8.27	8.85	8.40	
<u>Finishing period (89 days)</u>					
Beginning wt, lb ^{ae}	874	882	873	899	9.4
Ending wt, lb	1223	1225	1226	1242	13.3
Dry matter intake, lb/d	22.9	23.4	23.1	23.5	.71
Gain, lb/d	3.92	3.85	3.97	3.85	.08
Feed:gain	5.84	6.09	5.85	6.12	.14
<u>Total feeding trial (187 days)</u>					
Dry matter intake, lb/d ^{bc}	21.6	22.6	22.0	22.9	.33
Gain, lb/d	3.16	3.17	3.19	3.27	.07
Feed:gain ^b	6.85	7.11	6.93	7.04	.11
<u>Carcass</u>					
Hot carcass wt, lb	742	735	742	747	9.8
KPH, %	2.12	2.08	2.12	2.16	.04
Dressing %	60.6	60.0	60.5	60.1	.31
Backfat, in ^a	.43	.47	.47	.50	.03
REA, sq in ^b	12.48	11.82	12.10	12.14	.19
Yield grade	2.8	3.3	3.0	3.1	.12
Marbling score ^{abf}	2.7	3.2	2.9	3.1	.14
Quality grade ^{cg}	2.8	2.3	2.4	2.3	.14

^aEffect of UREA vs SBM (P<.10).

^bQuadratic effect of amino acid supplement (P<.11).

^cEffect of UREA vs SBM (P<.05).

^dNo statistics because steers were grouped by treatments into a single pen per treatment.

^eEffect of amino acid supplement vs SBM (P<.09).

^f2 = slight, 3 = small.

^g2 = choice, 3 = select.