

Temperature-sensitive mutant strains for isolation of additional mutants of a given site

S. Brody

Follow this and additional works at: <https://newprairiepress.org/fgr>



This work is licensed under a [Creative Commons Attribution-Share Alike 4.0 License](https://creativecommons.org/licenses/by-sa/4.0/).

Recommended Citation

Brody, S. (1966) "Temperature-sensitive mutant strains for isolation of additional mutants of a given site," *Fungal Genetics Reports*: Vol. 9, Article 10. <https://doi.org/10.4148/1941-4765.2033>

This Note on Methods is brought to you for free and open access by New Prairie Press. It has been accepted for inclusion in *Fungal Genetics Reports* by an authorized administrator of New Prairie Press. For more information, please contact cads@k-state.edu.

Temperature-sensitive mutant strains for isolation of additional mutants of a given site

Abstract

Use of temperature-sensitive strains

Brody, et temperature-sensitive mutant strains for isolation of additional mutants at a given site.

mapping and/or biochemical screening since the colonial phenotype can be due to mutation at many different genes (40 at least). The selection and isolation of temperature-sensitive revertants circumvented these problems. The rationale for this approach is as follows: revertants from a presumed point mutation in a structural gene may be due to mutations at the original site, elsewhere in the same gene, or at another locus. Quite often compensatory mutations at another site in the gene, the so-called second-site revertants, lead to the production of temperature-sensitive proteins. Therefore, some of the temperature-sensitive revertants may be second-site revertants. Approximately 40 wild-type revertants were isolated at 25°C and 4 of these were found to be colonial at 35°C. One of these temperature-sensitive strains was the strain desired; i.e., it had a temperature-sensitive glucose-6-P dehydrogenase.

Another temperature-sensitive revertant, which proved to be a temperature-sensitive suppressor strain, was helpful in that it was used as a source of conidia for the inositol-less death mutant selection technique. Selection of mutants in most colonial strains is difficult due to the lack of conidiation. However, this difficulty is bypassed by harvesting conidia from this suppressor strain grown at 25°C and then performing the mutant selection at 35°. Selection of certain mutants in colonial strains may be advantageous, since the altered metabolism of the colonial strains might not allow the growth of certain "leaky" mutants during inositol deprivation. * * * Rockefeller University, New York, New York. 10021.

For purposes of correlation with the results of studies on *N. crassa* strain col-2 (Y5331), it was desirable to isolate additional strains altered-e col-2 locus and to examine the properties of their glucose-6-P dehydrogenases. However, the isolation of a particular colonial strain would involve numerous mutant hunts and extensive