Isolation of nuclei from conidia

S. K. Dutta

Follow this and additional works at: <https://newprairiepress.org/fgr>

This work is licensed under a [Creative Commons Attribution-Share Alike 4.0 License](https://creativecommons.org/licenses/by-sa/4.0/).

Recommended Citation

Dutta, S. K. (1969) "Isolation of nuclei from conidia," *Fungal Genetics Reports*: Vol. 14, Article 11.
<https://doi.org/10.4148/1941-4765.2042>

This Technical Note is brought to you for free and open access by New Prairie Press. It has been accepted for inclusion in Fungal Genetics Reports by an authorized administrator of New Prairie Press. For more information, please contact cads@k-state.edu.

Isolation of nuclei from conidia

Abstract

Isolation of nuclei from conidia

Dutta, S. K. The isolation of nuclei
from *Neurospora crassa* conidia.

A greater degree of purity of nuclei isolated from conidial cells of *N. crassa* is obtained by combining the procedures described by Reich and Tsuda (1961 Biochem. Biophys. Acta 53:574) and Munkres et al. (1966 Neurospora Newsl. 9: 14) for mycelial cells with some modifications made in our laboratory. The

strain used was wild type 74A and all operations are conducted in the cold at 0-4°C. The conidial mass is squeezed dry between Whatman blotting papers and then ground gently with twice its volume of acid-washed sea sand (prepared by powdering the commercially obtained sea sand in a Wiley Mill and passing through a 60-mesh screen) until a smooth paste is obtained. About five volumes of sucrose-EDTA (0.5M sucrose, 1mM NaEDTA, 0.01M Tris HCl, pH 6.5) is added gradually, stirred into a thick paste, and filtered through four layers of silk cloth. The filtrate is centrifuged at 2000 x g for 25 minutes in a refrigerated centrifuge. This crude nuclear pellet is then suspended in a solution containing 0.5 M sucrose, 2 mM EDTA, and 5 mM CaCl_2 at pH 6.5 and centrifuged at low speed (500 x g) for two minutes. Two kinds of pellet were noticed. A hard pellet was formed below the loose pellet. The nuclei contained in the loose pellet were dispersed in their own supernatant and centrifuged again at 500 x g for two minutes. This low-speed centrifugation was repeated until no more hard pellet was noticed.

Comparatively pure nuclear pellet was obtained by passing this final loose pellet through 1.70 M sucrose solution containing 1 mM EDTA. This nuclear pellet was further cleaned from any cytoplasmic attachments by suspending and stirring for two hours in 10 volumes of saline EDTA (0.08 M NaCl, 0.02 M NaEDTA, pH 6.2), sedimenting at 2000 x g and resuspending in fresh solution of saline EDTA. In our process of chromatin isolation from these nuclei, we use triton-X-100 (0.01 %) along with saline EDTA in order to reduce the surface tension of nuclear membranes. The yield of nuclei was low by this process but a consistent purity (as judged by electron microscopy) was obtained, showing a 5:1 (total protein: DNA) ratio. Supported by NSF Grant No. Gy3894. ■ ■ ■ Department of Botany, Howard University, Washington, D. C. 20001.