

## Arginine transport in *Neurospora conidia*

W. B. Roess

A. G. DeBusk

Follow this and additional works at: <http://newprairiepress.org/fgr>

---

### Recommended Citation

Roess, W. B., and A.G. DeBusk (1965) "Arginine transport in *Neurospora conidia*," *Fungal Genetics Reports*: Vol. 8, Article 11.  
<https://doi.org/10.4148/1941-4765.2098>

This Research Note is brought to you for free and open access by New Prairie Press. It has been accepted for inclusion in *Fungal Genetics Reports* by an authorized administrator of New Prairie Press. For more information, please contact [cads@k-state.edu](mailto:cads@k-state.edu).

---

# Arginine transport in Neurospora conidia

## **Abstract**

Arginine transport in Neurospora conidia

## **Creative Commons License**



This work is licensed under a [Creative Commons Attribution-Share Alike 4.0 License](https://creativecommons.org/licenses/by-sa/4.0/).

Roess, W. B. and A. G. DeBusk. Arginine transport  
in *Neurospora conidia*.

*stereospecific transport processes, coupled with the amino acid growth inhibition data now in the literature, stimulated experiments in our laboratory on amino acid transport. The techniques used for the experiments reported here are similar to those previously reported for the characterization of the phenylalanine permease system ( DeBusk and DeBusk 1965 Biochim. Biophys. Acta 104: 139 ).*

The permease concept of metabolite entry into cells was first proposed by Rickenberg and co-workers in 1956 ( *Ann. Inst. Pasteur* 9: 829). The concept of genetically controlled

Arginine is actively transported into *Neurospora crassa* (74A) conidio by a constitutive, stereospecific permease system showing characteristic Michaelis-Menton kinetics and a Km of  $2 \times 10^{-6}$  M. The Process is temperature-dependent with an optimum at 35°C. A pH optimum occurs at 5.6. The amino acid is transported against a concentration gradient, resulting in an intracellular arginine concentration some 1450-fold greater than that of the external medium. The transport process is energy dependent as shown by its complete inhibition by  $\text{NaN}_3$  and DNP. No influx of previously accumulated arginine occurs either in the absence of external substrate or in the presence of energy uncoupling agents.

Stereospecificity of the transport system is indicated by transport competition studies with a number of amino acids. All L-isomers tested showed varying degrees of inhibition except proline which is characteristically a poor inhibitor for all permease systems studied. D-arginine, at concentrations 5-fold that of L-arginine, does not inhibit the transport of the L-isomer. The basic amino acids lysine and ornithine were very effective inhibitors, while glutamic acid was a poor inhibitor. The reduction in arginine transport at various inhibitor-to-arginine ratios is summarized in Table 1.

Simultaneous transport of pairs of amino acids was studied in order to further evaluate specificity and possible overlap of transport families. In all cases, the concentration of each amino acid was sufficiently high to saturate the permease enzyme(s) (rate independent of concentration). When lysine- $\text{C}^{14}$  and arginine- $\text{C}^{14}$  were simultaneously transported, the resulting rate was the average of their independent rates. This would indicate that arginine and lysine are transported by a common permease system. Very different results were obtained when phenylalanine- $\text{C}^{14}$  and arginine- $\text{C}^{14}$  were simultaneously transported. The initial rate of  $\text{C}^{14}$  transport was 80% of the sums of the independent rates for the individual amino acids. After 30 minutes the rate was nearly equal to the rate of arginine transport alone. This would suggest the existence of separate permeases for phenylalanine and arginine.

The inhibition of arginine transport by phenylalanine and other amino acids might be explained by the existence of general as well as specific permeases. Such a case has been clearly demonstrated for the aromatic amino acids in *Salmonella* (Ames, G. F. 1964 Arch. Biochem. Biophys. 104: 1).

This investigation was supported by a Public Health Service fellowship (GM-15,934) from the National Institutes of General Medical Sciences. ■ ■ ■ Genetics Laboratories, Department of Biological Sciences, Florida State University, Tallahassee, Florida.

Table 1, Arginine transport expressed as % of control,

	10:1	20:1	50:1	100:1
Lysine	28.0	15.2	—	5.2
Ornithine	—	29.6	18.3	12.0
Histidine	—	40.0	27.5	28.0
Phenylalanine	43.5	42.6	—	33.6
Tryptophan	—	32.0	32.0	27.5
Citrulline	—	55.4	41.6	36.0
Alanine	55.8	49.0	—	41.0
Isoleucine	—	54.0	44.5	37.2
Leucine	—	46.8	41.0	42.2
Methionine	—	44.5	40.0	39.2
Serine	—	68.0	55.0	47.0
Glutamic Acid	—	96.0	78.0	54.0
Glycine	—	75.5	60.3	53.0
Threonine	—	73.0	65.0	50.0
Proline	—	101.0	98.2	101.0