

Kansas Agricultural Experiment Station Research Reports

Volume 0
Issue 1 *Cattleman's Day (1993-2014)*

Article 1008

1987

Effect of stocker receiving diet on subsequent pasture gains

F. Brazle

Gerry L. Kuhl

D. Harmon

Follow this and additional works at: <https://newprairiepress.org/kaesrr>



Part of the [Other Animal Sciences Commons](#)

Recommended Citation

Brazle, F.; Kuhl, Gerry L.; and Harmon, D. (1987) "Effect of stocker receiving diet on subsequent pasture gains," *Kansas Agricultural Experiment Station Research Reports*: Vol. 0: Iss. 1. <https://doi.org/10.4148/2378-5977.2411>

This report is brought to you for free and open access by New Prairie Press. It has been accepted for inclusion in Kansas Agricultural Experiment Station Research Reports by an authorized administrator of New Prairie Press. Copyright 1987 Kansas State University Agricultural Experiment Station and Cooperative Extension Service. Contents of this publication may be freely reproduced for educational purposes. All other rights reserved. Brand names appearing in this publication are for product identification purposes only. No endorsement is intended, nor is criticism implied of similar products not mentioned. K-State Research and Extension is an equal opportunity provider and employer.



Effect of stocker receiving diet on subsequent pasture gains

Abstract

Seven diets were fed for 27 days to newly arrived heifers to evaluate their impact on stocker gains during the receiving and pasture periods. The diets were as follows: soybean hulls plus soybean meal; sun-dried alfalfa plus wheat middlings; dehydrated alfalfa plus grain sorghum; distillers dried grain plus cottonseed hulls; brome hay plus 2 lb protein supplement/day; prairie hay plus 2 lb protein supplement/day; and a commercial receiving ration. During the receiving period, diets of soybean hulls plus soybean meal, distillers dried grain plus cottonseed hulls, and the commercial receiving ration produced the highest cattle gains. The soybean hull plus soybean meal diet and the commercial receiving ration resulted in the best combined dry lot and pasture gains as well.


Keywords

Cattlemen's Day, 1987; Kansas Agricultural Experiment Station contribution; no. 87-309-S; Report of progress (Kansas State University. Agricultural Experiment Station and Cooperative Extension Service); 514; Beef; Stocker cattle; Diet; Pasture gains

Creative Commons License



This work is licensed under a [Creative Commons Attribution 4.0 License](https://creativecommons.org/licenses/by/4.0/).



Effect of Stocker Receiving¹ Diet on Subsequent Pasture Gains

Frank Brazle², Gerry Kuhl,
and David Harmon

Summary

Seven diets were fed for 27 days to newly arrived heifers to evaluate their impact on stocker gains during the receiving and pasture periods. The diets were as follows: soybean hulls plus soybean meal; suncured alfalfa plus wheat middlings; dehydrated alfalfa plus grain sorghum; distillers dried grain plus cottonseed hulls; brome hay plus 2 lb protein supplement/day; prairie hay plus 2 lb protein supplement/day; and a commercial receiving ration. During the receiving period, diets of soybean hulls plus soybean meal, distillers dried grain plus cottonseed hulls, and the commercial receiving ration produced the highest cattle gains. The soybean hull plus soybean meal diet and the commercial receiving ration resulted in the best combined drylot and pasture gains as well.

Introduction

Receiving rations for feedlot cattle are designed to adapt the rumen's microflora to a grain diet. Adapting the rumen's microflora to a lush pasture diet may be just as important. We evaluated seven receiving diets to determine their affect on pasture gains of intensive-early grazed stocker cattle.

Experimental Procedures

In March, 301 yearling heifers averaging 500 lbs were purchased in Oklahoma City and shipped to Moline, Kansas. The stockers were fed prairie hay and water on arrival and were processed the following morning. The cattle were vaccinated for Infectious Bovine Rhinotracheitis (IBR); Bovine Virus Diarrhea (BVD); Parainfluenza₃ (PI₃); Leptospirosis pomona; Clostridium chauvei, septicum, novyi, sordellii, and perfringens C and D (7-way); dewormed with Tramisol[®] injectable; and implanted with Ralgro[®].

The heifers were randomly allotted to 7 receiving rations with 3 pens of cattle on each diet. The major ingredients in the receiving rations were as follows:

¹ Appreciation is expressed to Delmer Miller, Scott City, KS and John ReQua Wichita, KS for providing cattle and to Ranch Aid, Inc., Eureka, KS; Bung Inc., Emporia, KS; Longton Dehydrating and Milling Co., Longton, KS; and American Diversified Co., Ethanol Plant, Hastings, NE for funding support. A sincere thank you to Jim Williams and Ed Durbin, Moline, KS for use of their facilities and to Dr. Williams, Moline, KS for assistance in processing and treating the cattle.

Appreciation is also extended to Agricultural Extension Agents' KelLee Parr, Elk Co.; Jeff Davidson, Greenwood Co.; Dale Lanham, Woodson Co.; and David Pace, Chautauqua Co.; and to Terry Goehring, Extension Assistant for data collection.

² Extension Livestock Specialist, Southeast Kansas.

1) soybean hulls (83%) plus soybean meal (10%); 2) suncured alfalfa (47%) plus wheat middlings (47%); 3) dehydrated alfalfa (55%) plus grain sorghum (39%); 4) distillers dried grain (41%) plus cottonseed hulls (51%); 5) a commercial receiving ration; 6) poor quality brome hay plus 2 lb soybean meal-based supplement per head daily; and 7) prairie hay plus 2 lb soybean meal-based supplement daily (Table 26.1). Cattle on the first 5 diets also received 2 lb of prairie hay per head daily.

All concentrate mixtures were pelleted and fortified with Deccox® (125 mg/head daily), salt, and vitamin A. All diets except the two hay-based rations were designed to contain minimum nutritional levels (dry basis) of 14% crude protein, 15% crude fiber, 60% TDN, .4% calcium, .3% phosphorus, and 1% potassium. Heifers were individually weighed at the start and finish of the 27-day receiving period, and at the end of 82-day grazing trial. Rumen fluid was collected from 30% of the heifers at the start and end of the receiving period. All heifers were intensive-early grazed in one native tallgrass pasture stocked at 2 acres/head.

Results and Discussion

Heifers fed the diets containing soybean hulls plus soybean meal, the commercial receiving ration, and distillers dried grain plus cottonseed hulls gained the fastest during the 27-day receiving period. Cattle fed the prairie and brome hay plus protein supplement diets were intermediate in gains during the receiving period, and those consuming wheat middlings plus suncured alfalfa or grain sorghum plus dehydrated alfalfa had the lowest gains (Table 26.2).

Highest cattle intake during the first 27 days was achieved on the commercial receiving ration, followed by the brome and prairie hay diets. The wheat middlings plus dehydrated alfalfa diet resulted in lowest heifer intake (Table 26.2). The best cattle feed conversions were obtained on the soybean hulls plus soybean meal, commercial receiving ration, and distillers dried grain plus cottonseed hull diets. Heifers fed the soybean hulls plus soybean meal and wheat middlings plus suncured alfalfa diets had the lowest ruminal pH and highest rumen propionic acid levels on day 27. Results suggest that these two rations had higher digestible energy levels than the other rations and may explain the lower heifer intakes, even though the soybean hulls plus soybean meal ration was one of the highest in crude fiber. About 25% of the heifers were treated for shipping fever, with no differences among treatments.

The heifers started on distillers dried grain plus cottonseed hulls had the lowest pasture gains. This may be explained by the fact that, although this ration was the highest in crude protein, its high ruminal bypass potential resulted in the lowest ($P < .05$) rumen ammonia level at the end of the receiving period, which likely inhibited ruminal microbial growth.

This trial confirms that the receiving ration fed to stockers before going to grass can affect short-term grass gains. Feeding a higher energy ration that results in faster gains for 3 to 4 weeks before grass will not necessarily reduce subsequent grass gains, as illustrated by the heifers fed diets containing soybean hulls plus soybean meal, and the commercial receiving ration. Feeding prairie hay plus 2 lb of protein supplement did not optimize stocker gains on grass, as has been suggested. However, stockers fed high energy starting rations for a longer period would be fleshier going to grass, which could result in lower grass gains.

Table 26.1. Ingredient Composition and Analysis of Receiving Rations

Items	Prairie Hay + 2 lb Prot. Suppl. ¹	Brome Hay + 2 lb Prot. Suppl. ¹	Distillers Dried Grain + Cottonseed Hulls	Soybean Hulls + Soybean Meal	Wheat Midds + Suncured Alfalfa	Grain Sorghum + Dehydrated Alfalfa	Commercial ² Receiving Ration
Ration Ingredients, lb/ton							
Soybean Meal	1835	1835	-----	200	-----	-----	-----
Soybean Hulls	-----	-----	-----	1667	-----	-----	-----
Cottonseed Hulls	-----	-----	1025	-----	-----	-----	-----
Dist. Dried Grain	-----	-----	832	-----	-----	-----	-----
Suncured Alfalfa	-----	-----	-----	-----	940	-----	-----
Wheat Midds	-----	-----	-----	-----	948	-----	-----
Dehydrated Alfalfa	-----	-----	-----	-----	-----	1100	-----
Grain Sorghum	-----	-----	-----	-----	-----	778	-----
Liquid Molasses	-----	-----	100	100	100	100	-----
White Salt	50	50	10	10	10	10	-----
Dicalcium Phosphate	100	100	-----	20	-----	10	-----
Limestone	-----	-----	20	-----	-----	-----	-----
Potassium Chloride	-----	-----	10	-----	-----	-----	-----
Vitamin A&D Premix	10	10	2	2	1	1	-----
Deccox Premix	5	5	1	1	1	1	-----
Chemical Analysis, Dry Matter Basis							
Crude Protein, %	5.0 ²	6.0 ²	17.9	15.2	14.5	15.2	16.2
Crude Fiber, %	36.0 ²	35.0 ²	24.0	33.0	22.0	19.0	26.0
Potassium, %	.83 ²	1.29 ²	1.51	1.41	1.63	2.77	1.17

¹Protein supplement ingredients only.²Purina Receiving Chow[®] with Deccox[®].³Hay analysis only.

Table 26.2. Effect of Starting Ration on Stocker Performance During Receiving and Grazing Periods

Item	Prairie Hay + 2 lb Prot. Suppl.	Brome Hay + 2 lb Prot. Suppl.	Distillers Dried Grain + Cottonseed Hulls	Soybean Hulls + Soybean Meal	Wheat Midds + Suncured Alfalfa	Grain Sorghum + Dehydrated Alfalfa	Commercial Receiving Ration
No. Heifers	43	43	43	43	43	43	43
Starting Wt., lb	493	475	485	483	483	491	490
Daily Gain, lb:							
Receiving-27 Days	2.16 ^{ab}	2.09 ^{ab}	2.30 ^a	2.30 ^a	1.88 ^b	1.92 ^b	2.45 ^a
Grazing-82 Days	1.24 ^a	1.36 ^a	1.08 ^b	1.34 ^a	1.24 ^a	1.36 ^a	1.31 ^a
Overall-109 Days	1.48 ^{ab}	1.54 ^{ab}	1.38 ^b	1.58 ^a	1.40 ^b	1.49 ^{ab}	1.59 ^a
Daily Feed Intake, lb:							
Receiving-27 Days	14.2 ^{ab}	14.5 ^a	14.0 ^b	13.8 ^b	13.4 ^b	13.8 ^b	14.6 ^a
% of Body Wt.	2.88	3.05	2.88	2.86	2.78	2.83	2.98
Feed/Gain, lb	6.6 ^b	7.0 ^{ab}	5.0 ^c	6.0 ^c	7.2 ^a	7.2 ^a	6.0 ^c
Rumen Parameters, Day 27:							
pH	6.78 ^b	6.72 ^b	6.82 ^b	6.45 ^a	6.51 ^a	6.83 ^b	6.72 ^b
Ammonia, mmoles/l	3.39 ^b	5.30 ^{ab}	.48 ^c	5.25 ^{ab}	3.65 ^b	5.69 ^{ab}	3.14 ^b
Total VFA's, mmoles/l	72.7 ^b	73.3 ^b	55.6 ^c	87.7 ^{ab}	76.6 ^b	60.6 ^c	64.6 ^c
Propionic Acid, mmoles/l	10.9 ^b	12.1 ^b	10.9 ^b	17.2 ^a	16.9 ^a	12.2 ^b	10.2 ^b

abc Means with different superscripts differ (P<.05).