

Kansas Agricultural Experiment Station Research Reports

Volume 0

Issue 1 *Cattleman's Day* (1993-2014)

Article 1054

1985

Action of decoquinate in altering feed efficiency of ruminants

D.L. Harmon

Follow this and additional works at: <https://newprairiepress.org/kaesrr>

 Part of the [Other Animal Sciences Commons](#)

Recommended Citation

Harmon, D.L. (1985) "Action of decoquinate in altering feed efficiency of ruminants," *Kansas Agricultural Experiment Station Research Reports*: Vol. 0: Iss. 1. <https://doi.org/10.4148/2378-5977.2457>

This report is brought to you for free and open access by New Prairie Press. It has been accepted for inclusion in Kansas Agricultural Experiment Station Research Reports by an authorized administrator of New Prairie Press. Copyright 1985 Kansas State University Agricultural Experiment Station and Cooperative Extension Service. Contents of this publication may be freely reproduced for educational purposes. All other rights reserved. Brand names appearing in this publication are for product identification purposes only. No endorsement is intended, nor is criticism implied of similar products not mentioned. K-State Research and Extension is an equal opportunity provider and employer.



Action of decoquinate in altering feed efficiency of ruminants

Abstract

The effects of adding decoquinate (Deccox®) to the diet at 0, 0.5, or 5 mg per kg body weight were evaluated with diets of 30 and 80% concentrate. Decoquinate did not influence rumen or plasma metabolites, diet digestibility, or volatile fatty acid production. No metabolic effects were seen that would suggest a role for decoquinate in altering rumen fermentation and feed efficiency of ruminants.

Keywords

Cattlemen's Day, 1985; Kansas Agricultural Experiment Station contribution; no. 85-319-S; Report of progress (Kansas State University. Agricultural Experiment Station and Cooperative Extension Service); 470; Beef; Feed efficiency; Decoquinate; Digestibility; Fermentation

Creative Commons License



This work is licensed under a [Creative Commons Attribution 4.0 License](https://creativecommons.org/licenses/by/4.0/).

K**S****U**

Action of Decoquinat in Altering Feed Efficiency of Ruminants

David L. Harmon

Summary

The effects of adding decoquinat (Deccox®) to the diet at 0, 0.5, or 5 mg per kg body weight were evaluated with diets of 30 and 80% concentrate. Decoquinat did not influence rumen or plasma metabolites, diet digestibility, or volatile fatty acid production. No metabolic effects were seen that would suggest a role for decoquinat in altering rumen fermentation and feed efficiency of ruminants.

Introduction

Decoquinat is a synthetic compound marketed under the name Deccox® as a feed additive to control coccidia infestations. Previous research has shown that decoquinat, added to the ration at the level of 0.5 mg/kg body weight per day, resulted in improved feed intake and animal performance. It is unclear whether these improvements were due to the action of decoquinat as a coccidiostat, or to independent effects. Our objective was to find out if decoquinat altered rumen fermentation and digestive efficiency.

Experimental Procedures

Two experiments were conducted using ruminally fistulated cattle (570 lb) consuming two levels of dietary concentrate; 30% in Experiment 1 and 80% in Experiment 2. Corn was the concentrate and soybean meal was the supplemental protein in both diets. Animals received either 0, 0.5, or 5 mg of decoquinat per kg body weight daily. Feed was offered in two equal portions daily with each animal receiving half of the daily dose of decoquinat in 0.45 lb of grain immediately prior to feeding.

We measured volatile fatty acids (VFA), pH, lactic acid, and ammonia levels as indicators of rate and efficiency of rumen fermentation, as well as rate of VFA production and digestibility of dry matter, starch, and several fiber components. Plasma lactate and glucose levels were also determined.

Results and Discussion

The data on rumen and serum metabolites (Table 16.1), rumen VFA concentrations (Table 16.2), starch, dry matter, and fiber digestibilities (Table 16.3), and VFA production (Table 16.4), indicate that decoquinat had little, if any, effect on digestion or rumen metabolism.

For the diets we evaluated, we see no evidence that would suggest a role for decoquinat, other than as a coccidiostat.

Table 16.1. Rumen and Plasma Metabolites in Steers Supplemented with Decoquinat

Item	Level of Decoquinat ^a					
	30% Concentrate			80% Concentrate		
	0	0.5	5.0	0	0.5	5.0
Rumen pH	6.61	6.60	6.64	6.53	6.49	6.38
Rumen NH ₃ , mg/dl	5.06	5.03	5.49	2.71	3.50	1.76
Rumen L(+) lactate, mM	.01	.01	.01	0.05	0.02	0.03
Plasma L(+) lactate, mM	1.12	0.55	0.85	0.57	0.59	0.68
Plasma Glucose, mg/dl	61.82	56.76	63.81	83.30	92.75	85.66

^amg per kg body weight daily.

Table 16.2. Rumen Volatile Fatty Acids in Steers Supplemented with Decoquinat

Item	Level of Decoquinat ^a					
	30% Concentrate			80% Concentrate		
	0	0.5	5.0	0	0.5	5
Acetate, molar%	75.56	75.12	75.69	65.78	64.28	70.24
Propionate, molar%	13.19	13.26	13.35	20.25	20.20	17.85
Butyrate, molar%	9.78	10.01	9.45	9.57	10.67	7.75
Others, molar%	1.47	1.60	1.51	4.39	4.85	4.16
Total VFA, mM	87.17	85.59	80.42	92.16	84.09	97.82

^amg per kg body weight daily.

Table 16.3. Intake and Digestibility of Steers Supplemented With Decoquinat

Item	Level of Decoquinat ^a					
	30% Concentrate			80% Concentrate		
	0	0.5	5.0	0	0.5	5.0
Dry Matter Intake, lb	13.34	12.94	13.58	15.92	15.81	16.05
Dry Matter Digestibility, %	54.99	53.53	52.87	64.70	65.93	64.96
Neutral Detergent Fiber Digestibility, %	55.27	47.89	56.19	62.65	68.63	61.22
Acid Detergent Fiber Digestibility, %	47.27	40.82	46.60	56.03	52.04	51.92
Hemicellulose Digestibility, %	60.71	59.14	61.40	67.20	79.36	62.39
Starch Digestibility, %	60.71	59.14	61.40	80.46	82.63	82.42

^amg per kg body weight daily.

Table 16.4. Rate of In Vitro Volatile Fatty Acid Production in Steers Supplemented with Decoquinat

Volatile Fatty Acid	Level of Decoquinat ^a					
	30% Concentrate			80% Concentrate		
	0	0.5	5	0	0.5	5.0
Acetate	24.27	28.00	34.67	28.20	30.09	29.68
Propionate	5.52	5.77	6.01	6.09	11.70	10.07
Butyrate	3.97	7.07	4.01	4.48	5.84	3.61
Total VFA	34.11	44.63	45.23	40.69	49.49	45.33

^amg per kg body weight daily.

^bmmole per liter rumen fluid per hour.