Kansas Agricultural Experiment Station Research Reports

Volume 0
Issue 1 *Cattleman's Day (1993-2014)*

Article 1152

1982

Trials on commercial silage additives

K. Bolsen

Follow this and additional works at: <https://newprairiepress.org/kaesrr>

Part of the [Other Animal Sciences Commons](#)

Recommended Citation

Bolsen, K. (1982) "Trials on commercial silage additives," *Kansas Agricultural Experiment Station Research Reports*: Vol. 0: Iss. 1. <https://doi.org/10.4148/2378-5977.2555>

This report is brought to you for free and open access by New Prairie Press. It has been accepted for inclusion in Kansas Agricultural Experiment Station Research Reports by an authorized administrator of New Prairie Press. Copyright 1982 the Author(s). Contents of this publication may be freely reproduced for educational purposes. All other rights reserved. Brand names appearing in this publication are for product identification purposes only. No endorsement is intended, nor is criticism implied of similar products not mentioned. K-State Research and Extension is an equal opportunity provider and employer.

K**S****U**Trials on Commercial Silage Additives¹

Keith Bolsen

Introduction

Numerous commercial silage additives, manufacturers of which make various claims for their improving silage quality, are available to Kansas farmers and ranchers. In 2 previous years (1980 and 1981), we reported on nine trials involving six additives (Progress Reports 377 and 394). Each additive improved the silage in at least one of the four criteria we evaluated: 1) ensiling temperature, 2) dry matter recovery, 3) feeding value, and 4) aerobic stability. Two more trials with additives are reported on in this Progress Report (pages 6 and 11).

Summary

Summarized in Table 5.1 are dry matter recoveries for 20 additive silages and their 11 controls in trials conducted from 1975 to 1981. Recovery of feedable dry matter averaged 85.0% for controls and 87.9% for additives. Fifteen additive silages had higher DM recoveries than did controls; only five silages, the same or lower recoveries.

Table 5.1. Dry matter recovery for control and additive silages in 11 trials conducted from 1975 to 1981.*

conducted from 1975 to 1981.*					
Silage and dry matter	Additive treatment	Recovery of feedable DM**	Silage and dry matter	Additive treatment	Recovery of feedable DM**
		%			%
Corn (38%)	control	80.9	Sorghum (33%)	control	91.0
	Silo Best ^R	87.5		Cold-flo	84.9
Corn (35%)	control	87.4	Alfalfa (36%)	Sila-bac	90.7
	Silo Guard ^R	93.7		control	84.6
Corn (44%)	control ^R	88.7		Ensila Plus	90.0
	Cold-flo ^R	91.5		Silo Guard ^R	89.7
	Sila-bac ^R	91.7	Sila-lator ^R	90.4	
	Silo-Best	91.3	Alfalfa (33%)	control	82.0
Corn (37%)	control	93.3		Sila-bac	82.0
	Cold-flo	88.5		Silo Guard	86.2
	Ensila Plus ^R	94.1	<hr/> 1981 trials <hr/>		
Wheat (42%)	control	77.6	Corn (32%)	control	87.3
	Ensila Plus	79.4		Silo-Best ^R	88.7
Sorghum (29%)	control	84.1		Sila-ferm ^R	87.4
	Silo Guard	92.0	Sorghum (30%)	control	78.1
		Sila-bac		81.1	
				LSA-100 ^R	77.2
			11 -Trial avg.	control	85.0
				additive	87.9

* All silages were made in 10 ft x 50 ft concrete stave silos.

** Percent of the dry matter ensiled.

¹ Mention of products and companies is made with the understanding that no discrimination or endorsement is intended. Also, no criticism is implied of products and companies not mentioned.