

# Kansas Agricultural Experiment Station Research Reports

---

Volume 0  
Issue 1 *Cattleman's Day (1993-2014)*

Article 1164

---

1982

## Effect of Bovatec on grazing steer performance

Lyle W. Lomas

Follow this and additional works at: <https://newprairiepress.org/kaesrr>



Part of the [Other Animal Sciences Commons](#)

---

### Recommended Citation

Lomas, Lyle W. (1982) "Effect of Bovatec on grazing steer performance," *Kansas Agricultural Experiment Station Research Reports*: Vol. 0: Iss. 1. <https://doi.org/10.4148/2378-5977.2567>

This report is brought to you for free and open access by New Prairie Press. It has been accepted for inclusion in Kansas Agricultural Experiment Station Research Reports by an authorized administrator of New Prairie Press. Copyright 1982 Kansas State University Agricultural Experiment Station and Cooperative Extension Service. Contents of this publication may be freely reproduced for educational purposes. All other rights reserved. Brand names appearing in this publication are for product identification purposes only. No endorsement is intended, nor is criticism implied of similar products not mentioned. K-State Research and Extension is an equal opportunity provider and employer.



---

## Effect of Bovatec on grazing steer performance

### Abstract

Steers fed 100 mg or 200 mg of Bovatec per head daily while grazing bromegrass gained 16.4% and 23.9% faster, respectively, during 112 days of grazing than did controls.

### Keywords

Cattlemen's Day, 1982; Report of progress (Kansas State University. Agricultural Experiment Station); 413; Beef; Steer Performance; Brome grass; Gain

### Creative Commons License



This work is licensed under a [Creative Commons Attribution 4.0 License](https://creativecommons.org/licenses/by/4.0/).

---

**K****S****U**

---

Effect of Bovatec<sup>1,3</sup> on Grazing Steer PerformanceLyle W. Lomas<sup>2</sup>Summary

Steers fed 100 mg or 200 mg of Bovatec<sup>1</sup> per head daily while grazing brome brome grass gained 16.4% and 23.9% faster, respectively, during 112 days of grazing than did controls.

Introduction

Previous research has shown Bovatec to be effective in increasing rate of gain and improving feed efficiency in feedlot cattle. Limited research has shown Bovatec to be effective in increasing rate of gain in grazing cattle. Our trial evaluated two levels of Bovatec in a grazing study.

Procedure

Seventy-two Simmental steers, averaging 645 lb, from one ranch were randomly allotted by weight to nine 10-acre brome grass pastures on June 24, 1981; so each pasture contained eight steers. Their treatments were: 1) control; 2) 100 mg of Bovatec per head daily; and 3) 200 mg of Bovatec per head daily. Each treatment was replicated three times.

All cattle received 3 lb of dry, rolled milo per head daily for the first 84 days and 5 lb for the last 28 days. Cattle on Bovatec received the additive daily; it was mixed with the rolled milo.

All cattle had free access to a mixture of equal parts steamed bone meal and trace-mineral salt and were provided fly control by dust bags.

Steers were weighed every 28 days. Initial and final weights were taken after a 16-hour shrink off feed and water. The study was terminated after 112 days, on October 14, 1981.

Results

During this 112-day grazing study, steers fed 100 mg of Bovatec per head daily gained 16.4% ( $P < .05$ ) and those fed 200 mg gained 23.9% ( $P < .01$ ) more than control steers did (Table 22.1). There was no statistically significant difference ( $P > .20$ ) between Bovatec levels. Bovatec caused no palatability problems.

---

<sup>1</sup>Bovatec is the trademark name for lasalocid sodium produced by Hoffmann-LaRoche, Inc., Nutley, N.J. 07110. Feed additive and partial financial assistance provided by Hoffmann-LaRoche, Inc.

<sup>2</sup>Southeast Kansas Branch Experiment Station, Parsons, KS 67357.

<sup>3</sup>Bovatec is not currently cleared by the FDA for use in cattle.

Table 22.1. Effect of Bovatec on Steer Performance - 112 days

Item	Level of Bovatec (mg/head/day)		
	0	100	200
No. of steers	24	24	24
Initial wt, lb	644	639	642
Final wt, lb	794 <sup>a,c</sup>	813 <sup>b,c,d</sup>	827 <sup>b,d</sup>
Total gain, lb	150 <sup>a,c</sup>	174 <sup>b,c,d</sup>	185 <sup>b,d</sup>
Average daily gain, lb	1.34 <sup>a,c</sup>	1.56 <sup>b,c,d</sup>	1.66 <sup>b,d</sup>

<sup>a,b</sup> Values in the same row with different superscripts differ significantly ( $P < .05$ ).

<sup>c,d</sup> Values in the same row with different superscripts differ significantly ( $P < .01$ ).