Kansas Agricultural Experiment Station Research Reports

Volume 0
Issue 1 *Cattleman's Day (1993-2014)*

Article 1177

1981

Continuous versus intermittent electrical stimulation of beef carcasses and their effect on hot-boned Muscle-pH decline

J.E. Bowles

Melvin C. Hunt

J.L.A. Kendall

See next page for additional authors

Follow this and additional works at: <https://newprairiepress.org/kaesrr>

Part of the [Other Animal Sciences Commons](#)

Recommended Citation

Bowles, J.E.; Hunt, Melvin C.; Kendall, J.L.A.; Lyon, M.; Kastner, Curtis L.; and Dikeman, Michael E. (1981) "Continuous versus intermittent electrical stimulation of beef carcasses and their effect on hot-boned Muscle-pH decline," *Kansas Agricultural Experiment Station Research Reports*: Vol. 0: Iss. 1. <https://doi.org/10.4148/2378-5977.2580>

This report is brought to you for free and open access by New Prairie Press. It has been accepted for inclusion in Kansas Agricultural Experiment Station Research Reports by an authorized administrator of New Prairie Press. Copyright 1981 the Author(s). Contents of this publication may be freely reproduced for educational purposes. All other rights reserved. Brand names appearing in this publication are for product identification purposes only. No endorsement is intended, nor is criticism implied of similar products not mentioned. K-State Research and Extension is an equal opportunity provider and employer.

Continuous versus intermittent electrical stimulation of beef carcasses and their effect on hot-boned Muscle-pH decline

Authors

J.E. Bowles, Melvin C. Hunt, J.L.A. Kendall, M. Lyon, Curtis L. Kastner, and Michael E. Dikeman

K**S****U**

Continuous Versus Intermittent
Electrical Stimulation of Beef Carcasses
and Their Effect on Hot-boned Muscle-pH Decline

J. E. Bowles, C. L. Kastner, M. E. Dikeman,
M. C. Hunt, J. L. A. Kendall, and M. Lyon

Summary

Short bursts of electricity (intermittent electrical stimulation) to beef carcasses accelerated pH decline and the onset of rigor mortis more than continuous stimulation did.

Introduction

Boning beef carcasses while they are still warm leads to large savings in power and cooler space. "Hot boning" processes and its advantages were discussed in the 1980 Cattlemen's Day publication. However, when a muscle undergoes rigor after being hot boned, it can become less tender. Rigor occurs because the muscle's chemical energy supply (glycogen) is converted to lactic acid, which causes the pH to drop, and rigor to occur. Using electrical stimulation to contract carcass muscles speeds up glycogen conversion to lactic acid, the subsequent pH drop, and rigor. Then, the carcass can be hot boned without decreasing tenderness.

The optimum combination of electrical stimulation (continuous or intermittent shock, voltage, postmortem stimulation time, duration of stimulation, amperage and frequencies) required for the fastest rigor onset has not been determined, so that is what we hoped to determine.

Experimental Procedure

Two groups of crossbred steers were slaughtered, electrically stimulated and hot boned. In group I, 46 sides were stimulated 1 hour postmortem with 400 volts of continuous, 60 cycle, alternating current for 2 minutes. Approximately .6 amp was delivered through the carcass. Two hours after slaughter the strip loin was hot boned. PH was monitored on the strip loin longissimus muscle.

In group II, 24 sides were electrically stimulated for 2 minutes at 45 minutes postmortem, using the same current characteristics except the current was pulsed, 1.6 second "on" and 0.8 second "off."

Results and Discussion

Both continuous and intermittent ES accelerated postmortem pH decline, but intermittent stimulation decreased the muscle pH slightly faster. Intermittent stimulation at 45 minutes postmortem may be more effective in accelerating postmortem pH decline than continuous stimulation at 1 hour postmortem. Allowing muscles to relax between contractions may allow the chemical energy from glycogen to be used up faster than when the muscle remains contracted.