

# Kansas Agricultural Experiment Station Research Reports

---

Volume 0  
Issue 1 *Cattleman's Day (1993-2014)*

Article 1189

---

1981

## Feeding MGA to grazing heifers

L. Corah

F. Brazle

Jeffrey L. Davidson

Follow this and additional works at: <https://newprairiepress.org/kaesrr>



Part of the [Other Animal Sciences Commons](#)

---

### Recommended Citation

Corah, L.; Brazle, F.; and Davidson, Jeffrey L. (1981) "Feeding MGA to grazing heifers," *Kansas Agricultural Experiment Station Research Reports*: Vol. 0: Iss. 1. <https://doi.org/10.4148/2378-5977.2592>

This report is brought to you for free and open access by New Prairie Press. It has been accepted for inclusion in Kansas Agricultural Experiment Station Research Reports by an authorized administrator of New Prairie Press. Copyright 1981 Kansas State University Agricultural Experiment Station and Cooperative Extension Service. Contents of this publication may be freely reproduced for educational purposes. All other rights reserved. Brand names appearing in this publication are for product identification purposes only. No endorsement is intended, nor is criticism implied of similar products not mentioned. K-State Research and Extension is an equal opportunity provider and employer.



---

## Feeding MGA to grazing heifers

### Abstract

Feeding MGA to grazing heifers suppressed estrus but did not improve gain.

### Keywords

Cattlemen's Day, 1981; Report of progress (Kansas State University. Agricultural Experiment Station); 394; Beef; MGA; Heifers

### Creative Commons License



This work is licensed under a [Creative Commons Attribution 4.0 License](https://creativecommons.org/licenses/by/4.0/).

**K**

## Feeding MGA to Grazing Heifers

**S**Larry Corah, Frank Brazle<sup>1</sup>, Jeff Davidson<sup>2</sup>**U**Summary

Feeding MGA to grazing heifers suppressed estrus but did not improve gain.

Introduction

MGA is widely used to promote growth and feed efficiency in feedlot heifers. But this trial was designed to study the efficacy of MGA on grazing heifers--an area that has received little attention.

Experimental Procedure

The trial was conducted at the Harold Engle Jr. ranch near Madison, Kans. April 28, 1980, 70 head of predominantly crossbred heifers were individually weighed, tagged, and allotted randomly in equal numbers to treatment or control groups. The two groups were pastured separately in 160-acre native grass pastures, and rotated every 20-30 days to eliminate pasture effects.

The heifers fed MGA received 1.4 pounds of a 14% range pellet formulated to supply .5 mg MGA/heifer/day. The control heifers received the same supplement without the MGA. Eighty days after the trial started, the heifers received an additional 3 pounds of a 17% range cube/heifer/day. Because of the extremely dry summer conditions, they were weighed off pasture August 2, 1980, much earlier than planned.

Results

Results are shown in Table .1. In spite of dry grazing conditions, the average daily gain of the heifers was excellent. The 3% gain improvement by heifers fed MGA was not statistically significant.

Table 18.1. Effect of MGA on the performance of grazing heifers.

Treatment	No.	Starting weight, lbs	Final weight, lbs	Pounds gained	A.D.G.
Control	32	500.6	664.5	163.9	2.48
MGA-fed	34	506.9	675.7	168.8	2.56

<sup>1</sup>Southeast Area Livestock Extension Specialist, Chanute, Kans.

<sup>2</sup>Greenwood County Extension Agricultural Agent, Eureka, Kans.