

Kansas Agricultural Experiment Station Research Reports

Volume 0
Issue 1 *Cattleman's Day (1993-2014)*

Article 1274

1977

Effect of using one versus two growth promoting implants on the gains of nursing calves

L.R. Corah

M. McKee

R.R. Schalles

Follow this and additional works at: <https://newprairiepress.org/kaesrr>



Part of the [Other Animal Sciences Commons](#)

Recommended Citation

Corah, L.R.; McKee, M.; and Schalles, R.R. (1977) "Effect of using one versus two growth promoting implants on the gains of nursing calves," *Kansas Agricultural Experiment Station Research Reports*: Vol. 0: Iss. 1. <https://doi.org/10.4148/2378-5977.2677>

This report is brought to you for free and open access by New Prairie Press. It has been accepted for inclusion in Kansas Agricultural Experiment Station Research Reports by an authorized administrator of New Prairie Press. Copyright 1977 Kansas State University Agricultural Experiment Station and Cooperative Extension Service. Contents of this publication may be freely reproduced for educational purposes. All other rights reserved. Brand names appearing in this publication are for product identification purposes only. No endorsement is intended, nor is criticism implied of similar products not mentioned. K-State Research and Extension is an equal opportunity provider and employer.



Effect of using one versus two growth promoting implants on the gains of nursing calves

Abstract

One hundred twenty-seven suckling calves were allotted to one of three treatments: (1) Control group - not implanted (2) Implanted (Ralgro) once during suckling period (3) Implanted (Ralgro) twice during suckling period - at average age of 44 days and 70 days later Using one Ralgro implant improved the weight at weaning by 8.4 pounds, while re-implanting, thus utilizing two implants during the suckling phase, resulted in an extra 43.0 pounds.

Keywords

Report of progress (Kansas State University. Agricultural Experiment Station); 291; Cattlemen's Day, 1977; Beef; Growth; Implants; Gain

Creative Commons License



This work is licensed under a [Creative Commons Attribution 4.0 License](https://creativecommons.org/licenses/by/4.0/).

K**Effect of Using One Versus Two Growth Promoting
Implants on the Gains of Nursing Calves****S**

Larry R. Corah, Miles McKee, and R. R. Schalles

U

Summary

One hundred twenty-seven suckling calves were allotted to one of three treatments:

- (1) Control group - not implanted
- (2) Implanted (Ralgro) once during suckling period
- (3) Implanted (Ralgro) twice during suckling period - at average age of 44 days and 70 days later

Using one Ralgro implant improved the weight at weaning by 8.4 pounds, while re-implanting, thus utilizing two implants during the suckling phase, resulted in an extra 43.0 pounds.

Introduction

Previous research published by Kansas State University and other Universities has shown that either DES or Ralgro will improve the weight gains of suckling calves 15 to 25 pounds. In most cases, calves are left on cows for 200 to 250 days, while the normal life span of most implants is 80 to 100 days.

Thus this trial was designed to compare the effect of using one versus two implants during the suckling phase. The implants used in the trial were 36 mg Ralgro implants.

Experimental Procedure

Forty-nine steer and 87 heifers, Polled Hereford or crossbreed calves, were divided into three groups based on breed, age, and sex. Sex of the calf had no effect on response to implanting so no further reference is made to calf sex. The three treatments were:

Treatment 1. Control group - not implanted

Treatment 2. One implant used during the suckling period when the calves were approximately 2½ months old.

Treatment 3. Two implants used during the suckling period. Initial implant when the calves were an average age of 44 days and re-implanted approximately 70 days later.

Two groups of calves were used. One group was suckling cows on native grass; the other group was suckling cows confined in drylot.

The calves that received one implant were implanted for a total of 114 days before weaning. Calves receiving two implants were implanted a total of 149 days during the suckling period.

Results and Discussion

Calves receiving one implant gained 8.4 pounds more than non-implanted calves at the end of the suckling period. This response is slightly under what has normally been shown in other research tests. Calves implanted when they averaged 1½ months and again approximately 70 days later, weighed 43.0 pounds more at weaning than calves not implanted.

No side effects were noted on any of the calves receiving one or two Ralgro implants during the suckling period.

Table 6.1 Results from using 0, 1, and 2 implants during suckling period of calves.

	<u>No. calves</u>	<u>Average age at initial implanting</u>	<u>No. days implanting to weaning</u>	<u>Birth wt. lbs.</u>	<u>Weaning wt.*</u>	<u>Gain lbs.*</u>	<u>Treatment advantage</u>
Not implanted control	41	---	---	72	372.6	300.6	---
One implant during suckling period	43	79 days	114 days	71	380.0	309.0	+8.4
Two implants during suckling period	43	44 days	149 days	71	414.6**	343.6**	+43.0**

* Weaning weights and lbs. gained were adjusted based on calf age.

**P<0.05.