

Kansas Agricultural Experiment Station Research Reports

Volume 0
Issue 1 *Cattleman's Day (1993-2014)*

Article 1287

1977

Feeding Monensin to yearling cattle on summer grass

F. Schwartz

E. Smith

Jack G. Riley

See next page for additional authors

Follow this and additional works at: <https://newprairiepress.org/kaesrr>



Part of the [Other Animal Sciences Commons](#)

Recommended Citation

Schwartz, F.; Smith, E.; Riley, Jack G.; and Corah, L. (1977) "Feeding Monensin to yearling cattle on summer grass," *Kansas Agricultural Experiment Station Research Reports*: Vol. 0: Iss. 1. <https://doi.org/10.4148/2378-5977.2690>

This report is brought to you for free and open access by New Prairie Press. It has been accepted for inclusion in Kansas Agricultural Experiment Station Research Reports by an authorized administrator of New Prairie Press. Copyright 1977 the Author(s). Contents of this publication may be freely reproduced for educational purposes. All other rights reserved. Brand names appearing in this publication are for product identification purposes only. No endorsement is intended, nor is criticism implied of similar products not mentioned. K-State Research and Extension is an equal opportunity provider and employer.



Feeding Monensin to yearling cattle on summer grass

Authors

F. Schwartz, E. Smith, Jack G. Riley, and L. Corah

K**Feeding Monensin to Yearling Cattle on Summer Grass****S**

Frank Schwartz, Ed Smith, Jack Riley and Larry Corah

U

Summary

Two grazing trials conducted at different locations in Kansas evaluated feeding Monensin to grazing yearling cattle. In one trial Monensin was fed with and without implant treatments of diethylstilbestrol or Ralgro. Monensin increased weight gain on summer grass; diethylstilbestrol and Ralgro implants also increased weight gains, and the combination of Monensin with either implant was more effective than Monensin alone.

Introduction

Previous research conducted at Kansas State with the newly released feed additive, Monensin showed average daily gain of cattle on growing rations increased by 5-8% and feed efficiency improved by 10-12%.

The current experiments reported here were conducted during the summer grazing season of 1976 at two locations in Kansas to study the effect of feeding Monensin to yearling cattle during the grazing season.

Experimental Procedure

Trial 1

Thirty-six Hereford and Angus-Santa Gertrudis cross steers were randomly allotted into two groups. Both groups were implanted with Ralgro and each grazed a 60-acre native bluestem pasture from April 28 to October 6, 1976. They had available in covered boxes commercial feed blocks¹ composed primarily of cane molasses; soybean meal feed, 20%; salt, 16-20%; and other feed ingredients. One group had Monensin added to the feed block at 327 mg. per pound. All steers were gathered the first of each month, penned overnight without feed or water, weighed the next morning, and rotated between pastures each month.

Trial 2

Sixty-two yearling cattle of mixed origin were allotted at random to

¹Feed blocks supplied by A. E. Staley Mfg. Co., Decatur, Ill., whose support is greatly appreciated.

two pasture groups at a ranch in northwestern Kansas.² The yearling cattle consisted of 52 steers and 10 heifers with the heifers randomly dispersed through the treatments. They were weighed directly off pasture initially and at the end of the trial. When weighed initially they were implanted, wormed, and vaccinated for black leg. Approximately one-third of the cattle in each pasture group received no implant, one-third were implanted with 30 mg. of diethylstilbestrol (DES), and one-third with Ralgro. The pasture was a typical northwestern Kansas short grass pasture; the two pasture groups were separated by an electric fence.

Two lbs of cracked corn were hand fed daily per head to both pasture groups. Monensin was mixed with the cracked corn to provide 100 mg. per animal per day to one of the pasture groups. The trial started May 2 with the cattle weighed directly off pasture July 21. The Monensin and grain were initially fed to cattle May 17 and then daily through July 21. Thus, Monensin was fed only for 65 days while the implanting comparison covered the full 80-day growing period.

Results and Discussion

Trial 1

Gain by steers getting Monensin on native bluestem pasture was 1.60 lbs. per steer daily compared with 1.32 for those not getting Monensin (Table 11.1). Intake of feed blocks available free choice averaged 0.32 lb. per steer daily for the Monensin group, slightly less for the non-Monensin. At that level steers received 123 mg. of Monensin per steer daily.

Trial 2

Use of Monensin over a 65-day trial increased the average daily gain .26 lb. which resulted in an extra 20.5 lbs. for the full 80-day trial period (Table 11.2). DES and Ralgro implants increased summer gains per animal 12.5 and 18.4 lbs., respectively (Table 11.3). Implants and Monensin in combination gave the best improvement in summer gains--37.6 extra pounds with the DES-Monensin and 35.2 extra pounds with Ralgro-Monensin (Table 11.4).

²Appreciation is expressed to Mr. Stan Albers and Ed Karnes, Hoxie, Kansas, for providing the cattle and pasture and to Sheridan County Extension Agent, Jim Grider, for assistance in conducting the trial.

Table 11.1. Feeding Monensin to steers on summer bluestem pasture - Trial 1.

	<u>No Monensin</u> lbs.	<u>Monensin</u> lbs.
Initial weight	471	476
Final weight	683	733
Daily gain	1.32**	1.60**
Feed block, lbs. consumed per steer daily		
May	0.17	0.23
June	0.22	0.23
July	0.26	0.46
Aug.	0.43	0.30
Sept.	0.24	0.40
Avg.	0.26	0.32

** Differ significantly ($P < 0.01$).

Table 11.2. Effect of Monensin on summer gains - Trial 2.

	<u>Treatment</u>	
	<u>Control</u>	<u>100 mg. Monensin/ hd/day</u>
No. cattle	31	31
Starting weight, lbs.	533	514
Final weight, lbs.	681.9	683.4
Lbs. gained	148.9*	169.4*
Average daily gain, lbs.	1.86*	2.12*
Treatment response, lbs.		+20.5

* $P < 0.05$.

Table 11.3. Effect of implants on summer gains - Trial 2.

	Treatment		
	<u>Control</u>	<u>DES</u>	<u>Ralgro</u>
No. cattle	19	20	23
Starting weight, lbs.	503.8	535.5	528.0
Final weight, lbs.	652.4	696.6	695.0
Lbs. gained	148.6 ^a	161.1 ^b	167.0 ^b
Average daily gain, lbs.	1.86 ^a	2.01 ^b	2.09 ^b
Treatment response, lbs.		+12.5	+18.4

^{ab}Means followed by dissimilar letters differ significantly ($P < 0.05$).

Table 11.4 Effect of using Monensin and implants in combination - Trial 2.

	<u>Control</u>	<u>DES and Monensin</u>	<u>Ralgro and Monensin</u>
	No. cattle	10	10
Starting weight, lbs.	510	524.5	517.5
Final weight, lbs.	646	698.1	688.7
Lbs. gained	136 ^a	173.6 ^b	171.2 ^b
Average daily gain, lbs.	1.70 ^a	2.17 ^b	2.14 ^b
Treatment response, lbs.		+37.6*	+35.2*

^{ab}Means followed by dissimilar letters differ ($P < 0.05$).