

Kansas Agricultural Experiment Station Research Reports

Volume 0
Issue 1 *Cattleman's Day (1993-2014)*

Article 1296

1977

Micronized milo, urea and prairie hay for growing beef heifers

K. Bolsen

J. Oltjen

Jack G. Riley

Follow this and additional works at: <https://newprairiepress.org/kaesrr>



Part of the [Other Animal Sciences Commons](#)

Recommended Citation

Bolsen, K.; Oltjen, J.; and Riley, Jack G. (1977) "Micronized milo, urea and prairie hay for growing beef heifers," *Kansas Agricultural Experiment Station Research Reports*: Vol. 0: Iss. 1. <https://doi.org/10.4148/2378-5977.2699>

This report is brought to you for free and open access by New Prairie Press. It has been accepted for inclusion in Kansas Agricultural Experiment Station Research Reports by an authorized administrator of New Prairie Press. Copyright 1977 Kansas State University Agricultural Experiment Station and Cooperative Extension Service. Contents of this publication may be freely reproduced for educational purposes. All other rights reserved. Brand names appearing in this publication are for product identification purposes only. No endorsement is intended, nor is criticism implied of similar products not mentioned. K-State Research and Extension is an equal opportunity provider and employer.



Micronized milo, urea and prairie hay for growing beef heifers

Abstract

Twenty-four individually fed heifers were used in two heifer trials to evaluate four combinations of micronized or dry-rolled milo and soybean meal or urea supplements in prairie hay growing rations. Feeding 4 to 5 lbs. of micronized milo produced an average of 17% faster and 16% more efficient gains than feeding 4 to 5 lbs. of dry-rolled milo.

Keywords

Report of progress (Kansas State University. Agricultural Experiment Station); 291; Cattlemen's Day, 1977; Beef; Milo; Urea; Prairie hay; Heifers

Creative Commons License



This work is licensed under a [Creative Commons Attribution 4.0 License](https://creativecommons.org/licenses/by/4.0/).

K

Micronized Milo, Urea and Prairie
Hay for Growing Beef Heifers

S

Keith Bolsen, Jim Oltjen and Jack Riley

U

Summary

Twenty-four individually fed heifers were used in two heifer trials to evaluate four combinations of micronized or dry-rolled milo and soybean meal or urea supplements in prairie hay growing rations. Feeding 4 to 5 lbs. of micronized milo produced an average of 17% faster and 16% more efficient gains than feeding 4 to 5 lbs. of dry-rolled milo.

Introduction

Previous research at Kansas State University and other midwest research stations has shown properly gelatinized milo superior to dry-rolled milo in high-grain, beef finishing rations. Also, soybean meal and urea supplements have generally given similar performance in high-grain rations; in high-roughage or silage rations, soybean meal has supported faster and more efficient gains than urea. Limited data are available comparing gelatinized milo and dry-rolled milo in high-roughage, beef cattle growing rations.

Our objective in this trial was to repeat a previous trial (Prog. Rpt. 262, Kan. Agr. Expt. Sta., 1976) evaluating four combinations of micronized or dry-rolled milo and soybean meal or urea supplements in hay rations for growing beef heifers.

Experimental Procedures and Results

Twenty-four Hereford and Hereford-Simmental heifers were allocated by breed and weight to sheltered, individual feeding pens. Six pens were assigned to each of these four treatments:

<u>Milo</u>		<u>Supplement</u>
1. dry-rolled	+	soybean meal (SBM)
2. dry-rolled	+	urea
3. micronized	+	soybean meal (SBM)
4. micronized	+	urea

All heifers were fed twice daily and received chopped prairie hay to appetite, 4 lbs. of the appropriate milo and 2 lbs. of the appropriate supplement daily. Both supplements contained 32% crude protein (as-fed basis).^a Initial and final weights of the heifers were taken after they

^aSoybean meal supplement: rolled milo, 688 lbs.; soybean meal, 1186 lbs.; dicalcium phosphate, 54 lbs.; salt, 42 lbs.; trace minerals, 8 lbs.; soybean oil, 21 lbs.; and vitamin A, 1 lb. Urea supplement: urea mix (100% CP), 514 lbs.; cane molasses, 390 lbs.; calcium lignin sulfate, 423 lbs.; trace minerals, 2 lbs.; 10-34-0, 70 lbs.; distillers' solubles, 600 lbs. and vitamin A, 1 lb.

had gone 15 hrs. without feed or water. The 98-day trial was conducted during the summer of 1976 with results shown in Table 22.1 (parts a and b).

Data from the 1975 and 1976 trials are compared in Table 22.2. In the 1975 trial, the heifers received 5 lbs. of milo daily and were fed 84 days. Results of the two trials were similar, although differences between treatments were smaller in 1976. Over both trials, heifers fed micronized milo + SBM or micronized milo + urea gained faster than heifers fed dry-rolled milo + urea (Table 22.2; part a). Heifers receiving micronized milo + SBM were more efficient than those receiving dry-rolled milo + urea.

Heifers receiving micronized milo gained faster and more efficiently than heifers receiving dry-rolled over both trials (Table 22.2; part b). Heifers fed SBM gained .07 lb. per day faster and required .82 lb. less dry matter per lb. of gain than those fed urea. Heifers fed the urea-containing liquid supplement consumed more hay than heifers fed the SBM supplement (10.55 vs. 10.15 lbs. daily in 1975; 11.24 vs. 10.53 lbs. daily in 1976). However, the higher moisture content of the urea supplement compared with the SBM supplement (45% vs. 13%), resulted in similar total dry matter consumptions.

Table 22.1. Performance of yearling heifers fed dry-rolled or micronized milo with soybean meal (SBM) or urea.¹

Part a:	Dry-rolled milo		Micronized milo	
	SBM	Urea	SBM	Urea
No. of heifers	6	6	6	6
Initial wt., lbs.	618.0	614.3	606.3	610.0
Final wt., lbs.	757.7	746.0	757.7	761.0
Avg. daily gain, lbs.	1.42	1.34	1.54	1.54
<u>Avg. daily feed, lbs.²</u>				
prairie hay	10.73	11.25	10.32	11.23
milo	3.63	3.75	3.66	3.80
supplement	1.59	1.10	1.55	1.08
Total	15.96	16.10	15.53	16.11
Feed/lb. of gain, lbs. ²	11.72	12.72	10.35	10.50
Part b:	Milo		Supplement	
	Dry-rolled	Micronized	SBM	Urea
No. of heifers	12	12	12	12
Avg. daily gains, lbs.	1.38 ^b	1.54 ^a	1.48	1.44
<u>Avg. daily feed, lbs.²</u>				
prairie hay	10.99	10.78	10.53	11.24
milo	3.69	3.73	3.65	3.78
supplement	1.34	1.31	1.57	1.08
Total	16.02	15.82	15.75	16.10
Feed/lb. of gain, lbs. ²	12.22 ^b	10.43 ^a	11.03	11.61

¹ 98 days (May 19 to August 25, 1976).

² 100% dry matter basis.

a,b Means on the same line with different superscripts differ significantly (P<.05).

Table 22.2. Comparison of the 1975 and 1976 heifer trials.

Part a:		Dry-rolled milo		Micronized milo	
		SBM	Urea	SBM	Urea
Avg. daily gain, lbs.	1975	1.31	1.25	1.66	1.50
	1976	1.42 _{bc}	1.34 _c	1.54 _a	1.54 _{ab}
	Avg.	1.36 ^{bc}	1.30 ^c	1.60 ^a	1.52 ^{ab}
Avg. daily ₁ feed, lbs.	1975	15.62	15.77	16.01	15.86
	1976	15.95	16.10	15.53	16.11
	Avg.	15.78	15.92	15.77	15.99
Feed/gain ¹	1975	12.12	12.55	9.69	10.98
	1976	11.72 _{ab}	12.72 _b	10.35	10.50 _{ab}
	Avg.	11.93 ^{ab}	12.63 ^b	10.02 ^a	10.74 ^{ab}
Part b:		Milo		Supplement	
		Dry-rolled	Micronized	SBM	Urea
Avg. daily gain, lbs.	1975	1.28	1.58	1.48	1.38
	1976	1.38 _b	1.54 _a	1.48	1.44
	Avg.	1.33 ^b	1.56 ^a	1.48	1.41
Avg. daily ₁ feed, lbs.	1975	15.70	15.93	15.81	15.81
	1976	16.02	15.82	15.75	16.10
	Avg.	15.85	15.88	15.78	15.95
Feed/gain ¹	1975	12.55	10.33	10.92	11.96
	1976	12.22 _b	10.43 _a	11.03	11.61
	Avg.	12.38 ^b	10.38 ^a	10.97	11.79

¹ 100% dry matter basis.

a,b,c Means on the same line with different superscripts differ significantly (P<.05).