

# Kansas Agricultural Experiment Station Research Reports

---

Volume 0  
Issue 1 *Cattleman's Day (1993-2014)*

Article 1297

---

1977

## Micronized corn and urea-liquid supplements in growing rations for calves

K. Bolsen

Leslie J. Chyba

W. Ibbetson

Follow this and additional works at: <https://newprairiepress.org/kaesrr>



Part of the [Other Animal Sciences Commons](#)

---

### Recommended Citation

Bolsen, K.; Chyba, Leslie J.; and Ibbetson, W. (1977) "Micronized corn and urea-liquid supplements in growing rations for calves," *Kansas Agricultural Experiment Station Research Reports*: Vol. 0: Iss. 1. <https://doi.org/10.4148/2378-5977.2700>

This report is brought to you for free and open access by New Prairie Press. It has been accepted for inclusion in Kansas Agricultural Experiment Station Research Reports by an authorized administrator of New Prairie Press. Copyright 1977 the Author(s). Contents of this publication may be freely reproduced for educational purposes. All other rights reserved. Brand names appearing in this publication are for product identification purposes only. No endorsement is intended, nor is criticism implied of similar products not mentioned. K-State Research and Extension is an equal opportunity provider and employer.



**K**

## Micronized Corn and Urea-liquid Supplements in Growing Rations for Calves

**S**

Keith Bolsen<sup>1</sup>, Les Chyba<sup>2</sup> and Wes Ibbetson<sup>2</sup>

**U**

### Summary

In growing rations containing fescue hay, micronized corn + urea-liquid gave: (1) 20.5% faster gains and 11.5% more efficient gains than cracked corn + urea-liquid and (2) similar rate and efficiency of gains as cracked corn + soybean meal.

### Introduction

Previous research at Kansas State University with high-grain rations showed micronized milo supporting a 16% more efficient gain than dry rolled milo when fed to finishing steers. In other trials reported here (Micronized Milo, Urea and Prairie Hay for Growing Beef Heifers) calves fed 4 to 5 lbs. of micronized milo gained 17% faster and 16% more efficiently than calves fed 5 lbs. of dry rolled milo. In the same trials, micronized milo plus a urea-liquid supplement supported slightly better performance than dry rolled milo plus a soybean meal supplement.

Our objective in this trial was to compare micronized and cracked corn in hay rations for growing calves.

### Experimental Procedures and Results

Fifty Hereford, Angus and HXA steer and heifer calves were allotted by breed, sex and weight to six pens. Two pens were assigned to each of the following corn and supplement treatments:

- (1) cracked corn + soybean meal (SBM)
- (2) cracked corn + urea-liquid
- (3) micronized corn + urea-liquid

All calves were fed twice daily and received long fescue hay to appetite, 4 lbs. of the appropriate corn and 2 lbs. of the appropriate supplement. Both supplements<sup>3</sup> contained 32% crude protein on an as-fed basis.

<sup>1</sup>Animal Science and Industry Dept., Kansas State University, Manhattan.

<sup>2</sup>Southeast Kansas Branch Experiment Station, Mound Valley.

<sup>3</sup>Soybean meal supplement: rolled milo, 674 lbs.; soybean meal, 1186 lbs.; dicalcium PO<sub>4</sub>, 54 lbs.; salt, 42 lbs.; trace minerals, 8 lbs.; molasses, 35 lbs. and vitamin A, 1 lb. Urea supplement: urea mix (100% CP), 514 lbs., cane molasses, 390 lbs.; calcium lignin sulfonate, 423 lbs.; trace minerals, 2 lbs.; 10-34-0, 70 lbs.; distillers' solubles, 600 lbs. and vitamin A, 1 lb.

The 105-day trial was conducted at the Southeast Kansas Branch Experiment Station from December 29, 1975 to April 12, 1976 (Table 23.1).

The overall performance of the calves was low, reflecting poor quality fescue hay.

Calves fed micronized corn + urea-liquid tended to gain faster and more efficiently than calves fed cracked corn + urea-liquid. Calves receiving micronized corn + urea-liquid or cracked corn + SBM had similar performance. SBM supported better performance than urea-liquid when each was fed with cracked corn. Calves fed urea-liquid rations consumed less feed ( $P < .05$ ) than those fed SBM.

Table 23.1. Performance of calves fed cracked or micronized corn and SBM or urea-liquid supplements.

	Cracked corn + SBM	Cracked corn + urea-liquid	Micronized corn + urea-liquid
No. of calves	16	18	16
Initial wt., lbs.	509	497	501
Final wt., lbs.	615	578	599
Avg. total gain, lbs.	106	81	98
Avg. daily gain, lbs.	1.01	.78	.94
<u>Avg. daily feed, lbs.<sup>1</sup></u>			
fescue hay	7.92	6.98	7.53
corn	3.40	3.40	3.60
supplement	1.78	1.10	1.10
Total	13.10 <sup>a</sup>	11.48 <sup>c</sup>	12.23 <sup>b</sup>
Feed/lb. of gain, lbs. <sup>1</sup>	13.06	14.72	13.02

<sup>1</sup>100% dry matter basis.

abc, Means on the same line with different superscripts differ significantly ( $P < .05$ ).