

Kansas Agricultural Experiment Station Research Reports

Volume 0
Issue 1 *Cattleman's Day (1993-2014)*

Article 1305

1976

Synchronizing estrus in beef heifers with prostaglandin and syncromate B

R.C. DeBenedetti

G.H. Kiracofe

V. Hultine

See next page for additional authors

Follow this and additional works at: <https://newprairiepress.org/kaesrr>



Part of the [Other Animal Sciences Commons](#)

Recommended Citation

DeBenedetti, R.C.; Kiracofe, G.H.; Hultine, V.; McKee, R.M.; and Schalles, R.R. (1976) "Synchronizing estrus in beef heifers with prostaglandin and syncromate B," *Kansas Agricultural Experiment Station Research Reports*: Vol. 0: Iss. 1. <https://doi.org/10.4148/2378-5977.2708>

This report is brought to you for free and open access by New Prairie Press. It has been accepted for inclusion in Kansas Agricultural Experiment Station Research Reports by an authorized administrator of New Prairie Press. Copyright 1976 Kansas State University Agricultural Experiment Station and Cooperative Extension Service. Contents of this publication may be freely reproduced for educational purposes. All other rights reserved. Brand names appearing in this publication are for product identification purposes only. No endorsement is intended, nor is criticism implied of similar products not mentioned. K-State Research and Extension is an equal opportunity provider and employer.



Synchronizing estrus in beef heifers with prostaglandin and syncromate B

Abstract

Forty-five of 50 heifers were in estrus 1 to 5 days after 7-day synchronization implants were removed. Prostaglandin was injected one day before implants were removed. Thirty-one of the 45 (68.9%) heifers conceived to first artificial insemination service.

Keywords

Cattlemen's Day, 1976; Report of progress (Kansas State University. Agricultural Experiment Station); 262; Beef; Synchronize; Prostaglandin; Syncromate B

Creative Commons License



This work is licensed under a [Creative Commons Attribution 4.0 License](https://creativecommons.org/licenses/by/4.0/).

Authors

R.C. DeBenedetti, G.H. Kiracofe, V. Hultine, R.M. McKee, and R.R. Schalles

K

Synchronizing Estrus in Beef Heifers with Prostaglandin and Syncromate B

S

R. C. DeBenedetti, G. H. Kiracofe, V. Hultine,
R. M. McKee and R. R. Schalles

U

Summary

Forty-five of 50 heifers were in estrus 1 to 5 days after 7-day synchronization implants were removed. Prostaglandin was injected one day before implants were removed. Thirty-one of the 45 (68.9%) heifers conceived to first artificial insemination service.

Introduction

We have previously reported that an ear implant (Syncromate B, G. D. Searle Co.) for 7 days followed by intramuscular injection of prostaglandin F_{2α} (PGF, The UpJohn Co.) synchronized estrus with no effect on conception rates in beef heifers. Heifers with a corpus luteum on the ovary when the PGF was injected showed estrus 12-24 hrs. later than those with no corpus luteum. The injected PGF is to regress the corpus luteum to allow estrus; we thought injecting the PGF one day before removing the implant would more closely synchronize estrus than previously reported.

Experimental Procedure

Fifty cycling heifers were implanted in the ear with Syncromate B, a synthetic progestogen that prevents estrus. Six days later each heifer was injected with 33.3 mg. of prostaglandin THAM salt. One day later we removed the implant. Heifers were checked frequently during the day for estrus and were artificially inseminated 12 to 24 hours after estrus was first detected. Conception rate was determined by rectal palpation.

Results and Discussion

Forty-one of the 50 heifers were in estrus in a three-day period and 45 of 50, in a five-day period; 31 of the 45 (68.9%) conceived the first insemination; 41 of the 45 (91.1%) conceived in two inseminations.

Five heifers not synchronized exhibited estrus within the next five days. Four of the 5 conceived with first service. Conception rate for the 50 heifers for the entire 65-day breeding season was 94%.

These results do not differ from those previously reported when prostaglandin was given when the implant was removed.

This procedure resulted in approximately 90% of cycling heifers showing estrus in a five-day period, and conception rates were the same as for nonsynchronized heifers.

Table 2.1 Occurrence of estrus and conception rates in heifers treated with Syncro-mate B and prostaglandin

	Days post treatment ^a										Total
	AM 1	PM 1	AM 2	PM 2	AM 3	PM 3	AM 4	PM 4	AM 5	PM 5	
No. in estrus	6	19	3	10	0	3	0	3	1	0	45 ^b
No. conceived 1st service	5	12	3	6	0	3	0	1	1	0	31

^a Implant removed AM day 0.

^b Five of 50 heifers showed no signs of estrus during the 5 days, four of the 5 later conceived with first service.