Kansas Agricultural Experiment Station Research Reports

Volume 0
Issue 1 *Cattleman's Day (1993-2014)*

Article 1355

1974

Influence of rumen fluid inoculation on incidence of sickness in newly arrived feeder calves

K.K. Bolsen

S. Armbruster

Homer K. Caley -1922

See next page for additional authors

Follow this and additional works at: <https://newprairiepress.org/kaesrr>

Part of the [Other Animal Sciences Commons](#)

Recommended Citation

Bolsen, K.K.; Armbruster, S.; Caley, Homer K. -1922; Fink, G.; and Riley, Jack G. (1974) "Influence of rumen fluid inoculation on incidence of sickness in newly arrived feeder calves," *Kansas Agricultural Experiment Station Research Reports*: Vol. 0: Iss. 1. <https://doi.org/10.4148/2378-5977.2758>

This report is brought to you for free and open access by New Prairie Press. It has been accepted for inclusion in Kansas Agricultural Experiment Station Research Reports by an authorized administrator of New Prairie Press. Copyright 1974 the Author(s). Contents of this publication may be freely reproduced for educational purposes. All other rights reserved. Brand names appearing in this publication are for product identification purposes only. No endorsement is intended, nor is criticism implied of similar products not mentioned. K-State Research and Extension is an equal opportunity provider and employer.

Influence of rumen fluid inoculation on incidence of sickness in newly arrived feeder calves

Authors

K.K. Bolsen, S. Armbruster, Homer K. Caley -1922, G. Fink, and Jack G. Riley

K**S****U**

Influence of Rumen Fluid Inoculation on Incidence of Sickness in Newly Arrived Feeder Calves

J. G. Riley, K. K. Bolsen, S. Armbruster, H. Caley and G. Fink

Summary

Inoculating newly arrived feeder calves with 250 ml. rumen fluid did not stimulate weight gain or decrease sickness. One hundred seventy steer calves were used in the 31 day study.

Introduction

Inoculating or drenching with various products is being promoted as a beneficial stimulus for increased weight gain and reduced sickness in newly arrived feeder cattle, particularly light weight calves. Theoretically, if rumen fluid could be obtained from steers already adapted to the ration to be fed and put directly into rumens of the new arrivals, the rumen fluid may stimulate development of favorable rumen micro-organisms.

The objective of this trial was to evaluate the feasibility and effectiveness of inoculating newly arrived feeder calves with rumen fluid.

Experimental Procedure

Two groups of feeder steers totaling 170 head were purchased from Texarkana, Texas in late July, 1973. They were offered long hay and fresh water free choice 24 hours before being vaccinated for IBR, Leptospirosis, Blackleg and Malignant Edema. Steers with temperatures above 103.5°F were inoculated with either 250 ml. of freshly collected rumen fluid or 250 ml. of tap water. Data from steers with temperatures below 103.5°F were taken to judge losses due to high temperatures. The steers were weighed individually at the beginning and end of the test. Steers were assigned at random to pens of 8-10 each for close daily observation. Thereafter, all steers with temperatures above 103.5°F were treated with antibiotics and records were kept on the number of treatments per steer. All steers were fed a ration composed of sorghum silage, rolled milo and protein supplement.

Results and Discussion

The effect of rumen fluid inoculation on weight gain and incidence of sickness in newly arrived feeder calves is shown in table 3.1. Steers with initial temperatures below 103.5°F gained 5 lb. (36 lb. vs. 31 lbs.) more than either group with initial temperatures above 103.5°F. Steers inoculated with rumen fluid required the most antibiotic treatments during the 31 day trial.

Results indicate that inoculating with rumen fluid had no benefits.

Table 3.1. Effects of Rumen Fluid Inoculation on Weight Gain and Incidence of Sickness in Newly Arrived Feeder Steer Calves (31 days)

	No. Steers	Initial wt lb.	31-day Gain, lb.	% treated 2 or more times with antibiotics
Non-Innoculated	84	467.8	36	42
Rumen Fluid Innoculated, 250 ml.	45	456.2	31	47
Water Innoculated, 250 ml.	41	462.9	31	34