

# Kansas Agricultural Experiment Station Research Reports

---

Volume 0  
Issue 1 *Cattleman's Day (1993-2014)*

Article 1362

---

1974

## Soybean meal and Starea (R) blocks compared by wintering cows on bluestem pasture: effects on birth and weaning weights of calves

L.H. Harbers

E.L. Shiawoya

K. Conway

*See next page for additional authors*

Follow this and additional works at: <https://newprairiepress.org/kaesrr>



Part of the [Other Animal Sciences Commons](#)

---

### Recommended Citation

Harbers, L.H.; Shiawoya, E.L.; Conway, K.; and McKee, R.M. (1974) "Soybean meal and Starea (R) blocks compared by wintering cows on bluestem pasture: effects on birth and weaning weights of calves," *Kansas Agricultural Experiment Station Research Reports*: Vol. 0: Iss. 1. <https://doi.org/10.4148/2378-5977.2765>

This report is brought to you for free and open access by New Prairie Press. It has been accepted for inclusion in Kansas Agricultural Experiment Station Research Reports by an authorized administrator of New Prairie Press. Copyright 1974 Kansas State University Agricultural Experiment Station and Cooperative Extension Service. Contents of this publication may be freely reproduced for educational purposes. All other rights reserved. Brand names appearing in this publication are for product identification purposes only. No endorsement is intended, nor is criticism implied of similar products not mentioned. K-State Research and Extension is an equal opportunity provider and employer.



---

## Soybean meal and Starea (R) blocks compared by wintering cows on bluestem pasture: effects on birth and weaning weights of calves

### Abstract

Similar gains and adjusted weaning weights can be expected from calves whose dams are wintered on soybean meal or Starea protein blocks. Rebreeding time is expected to be similar for both groups of dams.

### Keywords

Cattlemen's Day, 1974; Report of progress (Kansas State University. Agricultural Experiment Station); 210; Beef; Soybean meal; Starea; Bluestem pasture; Birth weight; Weaning weight

### Creative Commons License



This work is licensed under a [Creative Commons Attribution 4.0 License](https://creativecommons.org/licenses/by/4.0/).

### Authors

L.H. Harbers, E.L. Shiawoya, K. Conway, and R.M. McKee

**K****S****U**

Soybean Meal and Starea (R) Blocks Compared by Wintering  
Cows on Bluestem Pasture: Effects on Birth and Weaning  
Weights of Calves

L. H. Harbers, E. L. Shiawoya, K. Conway, and R. M. McKee

### Summary

Similar gains and adjusted weaning weights can be expected from calves whose dams are wintered on soybean meal or Starea portein blocks. Rebreeding time is expected to be similar for both groups of dams.

### Introduction

During the winter of 1972-73, 52 pregnant beef cows (Hereford and Angus) from 3 to 14 years old (average 5 years) were divided into two groups to compare soybean meal and Starea blocks during gestation (KAES Bulletin 568, 1972, p. 15). Summarized here are data collected on the growth performance of calves and the time cows rebred the next spring and summer.

### Methods

Estimates of adjusted 205-day weaning weights and rate of gain by calves were obtained from calf weights at weaning. Breeding dates were calculated from palpation estimates during September.

### Results

Average daily gains and adjusted weaning weights (steer equivalent) were similar for both groups (table 7.1). The average rebreeding date was the same for cows fed soybean meal or Starea.

Table 7.1. Rebreeding Dates of Cows and Performance of Calves Born to Cows Receiving Soybean Meal or Starea Blocks During Gestation

Item	Soybean meal block	Starea block
Calf weaning wt., lb.	500 (401-644) <sup>a</sup>	528 (436-627)
Calf daily gain, lb.	2.23 (1.66-3.19)	2.34 (1.98-2.81)
Rebreeding day of year	157.6 (June 6)	157.1 (June 6)

<sup>a</sup>Average (range).