

Kansas Agricultural Experiment Station Research Reports

Volume 0
Issue 1 *Cattleman's Day (1993-2014)*

Article 1374

1973

Influence of fly control on incidence of pinkeye and on calf performance

R.R. Schalles

M. McKee

J. Evans

See next page for additional authors

Follow this and additional works at: <https://newprairiepress.org/kaesrr>



Part of the [Other Animal Sciences Commons](#)

Recommended Citation

Schalles, R.R.; McKee, M.; Evans, J.; Davis, Duane L.; and Pitts, C.W. Jr. (1973) "Influence of fly control on incidence of pinkeye and on calf performance," *Kansas Agricultural Experiment Station Research Reports*: Vol. 0: Iss. 1. <https://doi.org/10.4148/2378-5977.2777>

This report is brought to you for free and open access by New Prairie Press. It has been accepted for inclusion in Kansas Agricultural Experiment Station Research Reports by an authorized administrator of New Prairie Press. Copyright 1973 the Author(s). Contents of this publication may be freely reproduced for educational purposes. All other rights reserved. Brand names appearing in this publication are for product identification purposes only. No endorsement is intended, nor is criticism implied of similar products not mentioned. K-State Research and Extension is an equal opportunity provider and employer.



Influence of fly control on incidence of pinkeye and on calf performance

Authors

R.R. Schalles, M. McKee, J. Evans, Duane L. Davis, and C.W. Jr. Pitts



Influence of Fly Control on Incidence of Pinkeye and on Calf Performance

R. R. Schalles, Miles McKee, Jack Evans,
D. L. Davis, and C. W. Pitts, Jr.¹

Summary

Controlling flies significantly decreased incidence of pinkeye in cattle on native Flint Hill range. There was no difference in the average weaning weight of groups sprayed or not sprayed. However individuals severely affected with pinkeye were much lighter than the average.

Introduction

Some type of fly control is usually recommended for cows and calves on summer pasture in Kansas. However the value of fly control is not well established. Nor have the relationships between fly control and incidence of pinkeye or between fly control and calf performance been evaluated. We studied those relationships for this report.

Experimental Procedure

Sixty-two yearling heifers and cows with 32 calves were sprayed with 4% Rabon as needed to control flies (every 7 to 14 days from May 15 to September 20). No fly control was used with another group of 62 yearling heifers and cows with 34 calves.

Cattle were checked several times a week. Individuals with pinkeye were treated with $\frac{1}{2}$ to 1 cc subconjunctival injection of antibiotic-corticosteroid mixture (dosage varied by size of animal).

Calves grazed spring-burned, native Bluestem pasture. All Cattle were weighed each month and the calves were weaned October 21 at an average age of 208 days.

Results

About 10% fewer cows had pinkeye in the fly-control group. Approximately 70% of the calves in each group had pinkeye; however those in the no-fly-control group averaged having pinkeye 2.3 times each, while those in the fly-control group averaged having pinkeye 1.6 times. Calves in the fly-

¹Department of Entomology

control group required 1.44 treatments per case of pinkeye compared with 1.74 treatments for those in the no-fly-control group. Sixty-four percent of all pinkeye treatments were on cattle in the no-fly-control group.

Greatest advantage of fly control in reducing pinkeye was early summer. Before July 15, 84% of the pinkeye treatments were on cattle in the no-fly-group. Cases of pinkeye were most numerous the last half of July. After July the incidence of pinkeye decreased and occurred about equally in both groups.

Average weaning weights of calves did not differ between groups, probably because most calves completely recover from pinkeye. However, some are permanently affected with impaired vision. In this study one calf was blind and had a 205-day adjusted weight 120 lbs. less than the average.

Because so many animals required treatment for pinkeye, both groups were handled about the same number of times.