

Kansas Agricultural Experiment Station Research Reports

Volume 0
Issue 1 *Cattleman's Day (1993-2014)*

Article 1387

1974

Effect of sound stress on ovulation, estrus, and conception in beef heifers

G. Heersche Jr.

G. Kiracofe

M. McKee

See next page for additional authors

Follow this and additional works at: <https://newprairiepress.org/kaesrr>



Part of the [Other Animal Sciences Commons](#)

Recommended Citation

Heersche, G. Jr.; Kiracofe, G.; McKee, M.; and Ames, D.R. (1974) "Effect of sound stress on ovulation, estrus, and conception in beef heifers," *Kansas Agricultural Experiment Station Research Reports*: Vol. 0: Iss. 1. <https://doi.org/10.4148/2378-5977.2790>

This report is brought to you for free and open access by New Prairie Press. It has been accepted for inclusion in Kansas Agricultural Experiment Station Research Reports by an authorized administrator of New Prairie Press. Copyright 1974 Kansas State University Agricultural Experiment Station and Cooperative Extension Service. Contents of this publication may be freely reproduced for educational purposes. All other rights reserved. Brand names appearing in this publication are for product identification purposes only. No endorsement is intended, nor is criticism implied of similar products not mentioned. K-State Research and Extension is an equal opportunity provider and employer.



Effect of sound stress on ovulation, estrus, and conception in beef heifers

Abstract

Thirty of the 50 heifers used in estrous synchronization studies and that received prostaglandin and Syncro-Mate B to synchronize estrus were subjected to sound stress for 48 hours after prostaglandin was injected. Fifty-nine percent of 29 sound-stressed heifers that showed estrus within 5 days conceived when artificially inseminated compared with 72 percent of 18 nonstressed heifers.

Keywords

Cattlemen's Day, 1974; Report of progress (Kansas State University. Agricultural Experiment Station); 210; Beef; Ovulation; Estrus; Conception; Heifers

Creative Commons License



This work is licensed under a [Creative Commons Attribution 4.0 License](https://creativecommons.org/licenses/by/4.0/).

Authors

G. Heersche Jr., G. Kiracofe, M. McKee, and D.R. Ames

K**S****U**

Effect of Sound Stress on Ovulation, Estrus, and Conception in Beef Heifers

G. Heersche, Jr., Guy Kiracofe, Mile McKee
and D. R. Ames

Summary

Thirty of the 50 heifers used in estrous synchronization studies and that received prostaglandin and Syncro-Mate B to synchronize estrus were subjected to sound stress for 48 hours after prostaglandin was injected. Fifty-nine percent of 29 sound-stressed heifers that showed estrus within 5 days conceived when artificially inseminated compared with 72 percent of 18 nonstressed heifers.

Introduction

Recent data in sheep indicate that exposing ewes to sound stress shortly before estrus increases ovulation but that does not affect conception rate or maintenance of pregnancy. We studied sound-stressed heifers.

Experimental Procedure

Fifty heifers received ear implants of Syncro-Mate B which were removed 7 days later. At implant removal, heifers were injected intramuscularly with 30 mgs of prostaglandin. Thirty of the 50 heifers were immediately placed in a barn and exposed to 8,000 Hz pure tones at 90 decibels intensity. The other 20 were protected from the sound (controls). The sound was played intermittently with the sound on 2 minutes and off 8 minutes. After 48 hrs of exposure, all heifers were returned to pens and bred 12 to 18 hours after the onset of estrus. Two weeks after the prostaglandin injection, sound-stressed and control heifers were rectally palpated to determine number of ovulations that had occurred at the last estrus. Numbers of corpora lutea on the ovaries were recorded as the number of ovulations. All heifers were pregnancy diagnosed 65 and 95 days after breeding.

Results and Discussion

Occurrence of estrus and conception rate are reported in Table 17.1. Rectal palpation gave no evidence that sound treatment affected ovulation rate. Onset of estrus occurred over a similar period of time and the duration of estrus was the same for sound stressed and control heifers. A lower percentage (59%) of heifers conceived in the sound-stressed group (17 of 29) than in the control group (72% or 13 of 18); however, work is needed to determine if sound caused the difference.

Table 17.1. Time of Estrus and First Service Conception Rate for Sound Stressed and Control Heifers

Item	Days after prostaglandin injected								Totals
	pm 1	am 2	pm 2	am 3	pm 3	am 4	pm 4	am 5	
No. Controls pregnant from first service	3	3	2	2	0	1	1	1	13
No. sound stressed in estrus	4	4	12	7	0	1	1	0	29 ^a
No. Controls in estrus	4	5	3	2	0	2	1	1	18 ^b
No. sound stressed pregnant from first service	3	1	10	3	0	0	0	0	17

^aOne heifer was not synchronized (estrus, am day 7).

^bOne heifer was not in estrus until day 19, another heifer did not show estrus.