

Kansas Agricultural Experiment Station Research Reports

Volume 0
Issue 1 *Cattleman's Day (1993-2014)*

Article 1425

1971

Comparison of Biuret and soybean meal for wintering cows on bluestem pasture II. Effect on birth and weaning weight of progeny

H.A. Thyault

L.H. Harbers

E.F. Smith

Follow this and additional works at: <https://newprairiepress.org/kaesrr>



Part of the [Other Animal Sciences Commons](#)

Recommended Citation

Thyault, H.A.; Harbers, L.H.; and Smith, E.F. (1971) "Comparison of Biuret and soybean meal for wintering cows on bluestem pasture II. Effect on birth and weaning weight of progeny," *Kansas Agricultural Experiment Station Research Reports*: Vol. 0: Iss. 1. <https://doi.org/10.4148/2378-5977.2828>

This report is brought to you for free and open access by New Prairie Press. It has been accepted for inclusion in Kansas Agricultural Experiment Station Research Reports by an authorized administrator of New Prairie Press. Copyright 1971 Kansas State University Agricultural Experiment Station and Cooperative Extension Service. Contents of this publication may be freely reproduced for educational purposes. All other rights reserved. Brand names appearing in this publication are for product identification purposes only. No endorsement is intended, nor is criticism implied of similar products not mentioned. K-State Research and Extension is an equal opportunity provider and employer.



Comparison of Biuret and soybean meal for wintering cows on bluestem pasture II. Effect on birth and weaning weight of progeny

Abstract

During the winter of 1969-70, 48 five-year-old cows were divided into three groups to compared soybean meal with biuret as a winter supplement in combination with sorghum grain (Bulletin 536, 1970, p.33). Soybean meal supplemented cows gained 31 lbs. each during 4 months while biuret-fed cows lost 15 lbs. each. Cows fed sorghum with biuret in a separate mineral mix (fed free choice) lost 62 lbs. each.

Keywords

Cattlemen's Day, 1971; Report of progress (Kansas State University. Agricultural Experiment Station); 546; Beef; Biuret; Soybean meal; Bluestem pasture; Birth weight; Weaning weight

Creative Commons License



This work is licensed under a [Creative Commons Attribution 4.0 License](https://creativecommons.org/licenses/by/4.0/).

K Comparison of Biuret and Soybean Meal for Wintering
Cows on Bluestem Pasture

S II. Effect On Birth and Weaning Weight of Progeny
(Project 253)

U H. A. Thyault, L. H. Harbers, and E. F. Smith

During the winter of 1969-70, 48 five-year-old cows were divided into three groups to compare soybean meal with biuret as a winter supplement in combination with sorghum grain (Bulletin 536, 1970, p. 33). Soybean supplemented cows gained 31 lbs. each during 4 months while biuret-fed cows lost 15 lbs. each. Cows fed sorghum grain with biuret in a separate mineral mix (fed free choice) lost 62 lbs. each.

Birth weight and 205-day, adjusted, weaning weight (steer equivalent) of calves born during and after the winter-supplementation study are given in table 24. When the supplements were mixed with sorghum grain, soybean meal and biuret were equally good; but cows given biuret in a mineral mix gave birth to smaller calves ($P < .05$). Weaning weights were lower in the biuret-mineral group but not significantly different from those fed protein supplements mixed with sorghum grain.

Table 24. Birthweight And Adjusted Weaning Weight of Calves Born to Cows Receiving Soybean Meal or Biuret as Protein Supplements

Item	Sorghum grain- soybean meal	Sorghum grain-biuret	Mineral-mix biuret
Birth Wt., Lb.	67 (55-60) ^a	67 (50-80)	62 (44-70)
Weaning Wt., Lb.	422 (316-476)	423 (329-481)	407 (286-461)

^a Average, with range

Biuret can be successfully added to complete wintering cow supplements but, in a separate mineral mix, biuret did not compare favorably with mixtures of either soybean meal and sorghum grain or biuret and sorghum grain.