Kansas Agricultural Experiment Station Research Reports

Volume 0
Issue 1 *Cattleman's Day (1993-2014)*

Article 1429

1970

Urea and soybean meal compared for cows on winter bluestem pasture

R.W. Swanson

E.F. Smith

Follow this and additional works at: <https://newprairiepress.org/kaesrr>

Part of the [Other Animal Sciences Commons](#)

Recommended Citation

Swanson, R.W. and Smith, E.F. (1970) "Urea and soybean meal compared for cows on winter bluestem pasture," *Kansas Agricultural Experiment Station Research Reports*: Vol. 0: Iss. 1. <https://doi.org/10.4148/2378-5977.2832>

This report is brought to you for free and open access by New Prairie Press. It has been accepted for inclusion in Kansas Agricultural Experiment Station Research Reports by an authorized administrator of New Prairie Press. Copyright 1970 the Author(s). Contents of this publication may be freely reproduced for educational purposes. All other rights reserved. Brand names appearing in this publication are for product identification purposes only. No endorsement is intended, nor is criticism implied of similar products not mentioned. K-State Research and Extension is an equal opportunity provider and employer.

Urea and Soybean Meal Compared for
Cows on Winter Bluestem Pasture
1968-1969 (Project 253)

R.W. Swanson and E.F. Smith

This test compared urea supplement (hand-fed), urea supplement (self-fed), and soybean meal (SBM) supplement (hand-fed) with cows on winter bluestem pasture. The supplements were formulated to supply the same amount of protein and total digestible nutrients. Salt was fed free choice with the hand-fed supplement.

The self-fed supplement presented many problems because limiting intake of the supplement to 3.0 lbs. per head per day required from .30 lb. to .85 lb. of salt per head per day.

Supplement compositions are shown in table 12; test results, in table 11.

The SBM supplement was superior to the urea supplements in maintaining cows' weight, percentage of calves weaned, and percentage of cows breeding back.

Self-feeding the urea supplement produced lighter calves and fewer cows breeding back than did hand-feeding the urea supplement.

Table 11. Results From Supplementing Winter Bluestem Pasture with Indicated Supplements

	<u>Urea supplement</u>	<u>Urea self-fed</u>	<u>Soybean meal supplement</u>
1968-Fall cows' wt., lb.	869	826	864
1969-Spring cows' wt., lb.	598	574	655
1969-Fall cows' wt., lb.	939	898	937
1969-% Calf crop weaned	87.5	87.5	91.7
1969-Adj. calf weaning wt., lb.	408	365	396
1969-% Cows bred	95.8	87.5	100

Table 12. Composition of Indicated Supplements

<u>Urea supplement, lbs.</u>		<u>Urea self-fed, lbs.</u>		<u>Soybean meal supplement, lbs.</u>	
Sorghum grain	940	Sorghum grain	625	Sorghum grain	655
Urea	42	Urea	28	SBM	327
Dicalcium	18	Dicalcium	12	Dicalcium	18
phosphate		phosphate		phosphate	
	<u>1000</u>	Salt ¹	<u>335</u>		<u>1000</u>
			<u>1000</u>		
Pounds of Supplement per cow daily	3.2		4.1		3.0

¹Salt was used to limit consumption to 3 lb./head/day. Salt needed varied from .30 to .85 lb./head/day.