

Kansas Agricultural Experiment Station Research Reports

Volume 0
Issue 1 *Cattleman's Day (1993-2014)*

Article 1482

2010

Dried distillers grains supplemented at different frequencies to stocker heifers grazing late-season Flint Hills native pastures

M.P. Epp

W.L. Metzen

Brandon E. Oleen

See next page for additional authors

Follow this and additional works at: <https://newprairiepress.org/kaesrr>



Part of the [Other Animal Sciences Commons](#)

Recommended Citation

Epp, M.P.; Metzen, W.L.; Oleen, Brandon E.; and Blasi, Dale A. (2010) "Dried distillers grains supplemented at different frequencies to stocker heifers grazing late-season Flint Hills native pastures," *Kansas Agricultural Experiment Station Research Reports*: Vol. 0: Iss. 1. <https://doi.org/10.4148/2378-5977.2885>

This report is brought to you for free and open access by New Prairie Press. It has been accepted for inclusion in Kansas Agricultural Experiment Station Research Reports by an authorized administrator of New Prairie Press. Copyright 2010 Kansas State University Agricultural Experiment Station and Cooperative Extension Service. Contents of this publication may be freely reproduced for educational purposes. All other rights reserved. Brand names appearing in this publication are for product identification purposes only. No endorsement is intended, nor is criticism implied of similar products not mentioned. K-State Research and Extension is an equal opportunity provider and employer.



Dried distillers grains supplemented at different frequencies to stocker heifers grazing late-season Flint Hills native pastures

Abstract

Wetter-than-normal summer grazing seasons can produce excessive forage beyond what the typical Flint Hills double-stock harvest rate can remove. These late-summer native grasses do not contain adequate protein to sustain economical gains for stocker cattle, but producers can extend the grazing season by using a protein-based supplement to generate economically feasible rates of gain. Use of dried distillers grains as a supplementation program can help promote overall productivity during the fall grazing season. However, the drawback is the potential high labor input for supplement delivery. Delivering supplements at reduced frequencies per week may be able to sustain daily gains of stocker cattle while reducing labor costs associated with supplement delivery.

Keywords

Cattlemen's Day, 2010; Kansas Agricultural Experiment Station contribution; no. 10-170-S; Report of progress (Kansas State University. Agricultural Experiment Station and Cooperative Extension Service); 1029; Beef Cattle Research, 2010 is known as Cattlemen's Day, 2010; Beef; Dried distillers grains; Stocker heifers; Protein; Daily gain

Creative Commons License



This work is licensed under a [Creative Commons Attribution 4.0 License](https://creativecommons.org/licenses/by/4.0/).

Authors

M.P. Epp, W.L. Metzen, Brandon E. Oleen, and Dale A. Blasi

Dried Distillers Grains Supplemented at Different Frequencies to Stocker Heifers Grazing Late-Season Flint Hills Native Pastures

M.P. Epp, D.A. Blasi, W.L. Metzner, and B.E. Olen

Introduction

Wetter-than-normal summer grazing seasons can produce excessive forage beyond what the typical Flint Hills double-stock harvest rate can remove. These late-summer native grasses do not contain adequate protein to sustain economical gains for stocker cattle, but producers can extend the grazing season by using a protein-based supplement to generate economically feasible rates of gain. Use of dried distillers grains as a supplementation program can help promote overall productivity during the fall grazing season. However, the drawback is the potential high labor input for supplement delivery. Delivering supplements at reduced frequencies per week may be able to sustain daily gains of stocker cattle while reducing labor costs associated with supplement delivery.

Experimental Procedures

One 72-day grazing period was conducted at the Kansas State University Beef Stocker Unit starting September 1, 2009, to determine the response to various frequencies of dried distillers grains supplementation of heifers grazing Flint Hills native grasses. All cattle were sourced from a local sale barn 12 days prior to the study, vaccinated for clostridial and respiratory diseases, dewormed, and fed a high-roughage diet free choice. To obtain a more uniform group of animals, cattle with extreme high or low weights or cattle that exhibited signs of prior illness, lameness, or poor disposition were removed from the pool of cattle eligible for the experiment.

Cattle were randomized and stratified by weight to obtain an equal stocking rate per paddock. The stocking rate per paddock was approximately 63 lb/acre. There were three treatment groups and three replications (paddocks) per treatment group. All cattle were fed dried distillers grains at a daily rate of 0.33% body weight (dry matter basis), but the dried distillers grains were delivered at one of three frequencies: every day, every other day, or every third day. On the basis of predicted gains of 1.5 lb/head daily, dried distillers grains supplementation was increased every 2 weeks during the grazing period. All cattle were hand-fed in bunks located in each pasture. Forage samples were collected at three different times during the grazing study from four different pastures and submitted to a commercial laboratory for nutrient analysis.

Performance data were analyzed by using the MIXED model procedure of SAS (SAS Institute, Inc., Cary, NC). Data were arranged in a randomized complete block design, and pasture was the experimental unit for growth outcomes as affected by treatment. In the model, fixed effects were treatment and pasture, and random effects were pasture \times treatment, pasture, and animal.

Results and Discussion

Average daily gain was not different among the treatment groups (Table 1). Cattle fed dried distillers grains every other day or every third day gained at essentially the same rate as cattle supplemented daily during the 72-day grazing period that was initiated in September 2009.

Crude protein and energy content of the clipped forage collected from the four paddocks declined at a marked rate over this study (Table 2).

Implications

If adequate grass resources are available in late summer, producers can reduce labor costs by supplementing dried distillers grains to cattle every second or third day and still achieve performance that is similar to feeding dried distillers grains daily.

Table 1. Performance of stocker heifers fed dried distillers grains (DDG) at a daily rate of 0.33% of body weight (dry basis) at different frequencies while grazing Flint Hills native grasses in late summer and fall

Item	DDG supplementation frequency			SEM	P-value
	Daily	Every other day	Every third day		
No. of pastures	3	3	3		
No. of cattle	17	17	17		
DDG ¹ , total lb fed	508	507	501		
Initial weight, lb	620	622	614	8.7	>0.54
Final weight, lb	727	728	715	10.1	>0.38
Average daily gain, lb	1.51	1.49	1.43	0.15	>0.71

¹ DDG nutrient content: 90% dry matter, 30% crude protein.

Table 2. Fall 2009 nutritional quality of native pastures at the Kansas State University Beef Stocker Unit

Item	Pasture 3			Pasture 7			Pasture 13			Pasture 17		
	Sampling date			Sampling date			Sampling date			Sampling date		
	8/31	9/29	11/2	8/31	9/29	11/2	8/31	9/29	11/2	8/31	9/29	11/2
Nutrient composition												
Dry matter, %	38.3	36.3	73.4	44.3	40.4	72.9	43.0	44.2	80.27	41.2	43.9	77.3
Crude protein, %	6.8	4.9	3.1	5.3	4.7	3.1	5.3	4.7	3.04	6.3	4.2	2.5
Soluble crude protein, % of crude protein	28.8	25.0	25.0	40.6	25.0	25.0	25.0	25.0	25.0	38.3	25.0	25.0
Fat, %	2.4	2.5	2.1	1.8	2.4	2.5	1.9	1.8	2.5	2.4	1.9	2.3
Acid detergent fiber, %	36.0	36.4	40.4	34.3	36.4	41.1	33.8	37.5	37.7	36.0	40.3	42.0
Neutral detergent fiber, %	64.6	62.2	68.2	62.4	62.1	69.3	61.4	63.9	64.1	62.3	68.3	70.7
Calculated analysis												
Net energy maintenance, Mcal/cwt	58.0	52.3	49.6	50.8	51.9	46.5	52.7	47.3	49.2	56.9	50.0	49.6
Net energy gain, Mcal/cwt	32.1	26.8	24.4	25.5	26.5	21.5	27.2	22.2	24.0	31.1	24.2	24.4
Total digestible nutrients, %	59.0	55.3	53.6	54.4	55.1	51.6	55.6	52.1	53.3	58.3	53.9	53.6
Calcium, %	0.38	0.82	0.43	0.47	1.03	0.97	0.83	0.84	1.24	0.56	0.47	0.47
Phosphorus, %	0.12	0.07	0.04	0.10	0.13	0.06	0.09	0.07	0.04	0.11	0.08	0.05
Magnesium, %	0.20	0.23	0.14	0.24	0.24	0.17	0.10	0.09	0.11	0.23	0.19	0.14
Potassium, %	1.16	0.69	0.11	1.12	0.65	0.25	0.83	0.67	0.12	1.09	0.66	0.19
Sulfur, %	0.12	0.08	0.05	0.11	0.09	0.07	0.08	0.07	0.07	0.10	0.07	0.05
Iron, ppm	178.57	209	153	234.29	291	263	238.37	216	175	---	174	118
Manganese, ppm	41.29	38	36	16.51	17	21	37.15	49	53	---	32	40
Copper, ppm	11.35	14	11	11.35	12	10	11.35	13	13	---	13	10
Zinc, ppm	29.93	38	34	32.00	35	43	28.89	33	39	---	30	30
Molybdenum, ppm	---	1	1	3.10	2	1	1.03	1	1	---	1	1
Cobalt, ppm	---	1	1	3.10	1	1	---	1	1	---	1	1
Selenium, ppm	---	0.05	0.04	---	0.03	0.03	---	0.02	0.03	---	---	0.03