

Kansas Agricultural Experiment Station Research Reports

Volume 0
Issue 1 *Cattleman's Day (1993-2014)*

Article 1487

2010

Initial heifer body composition has little impact on response to Zilmax

L.K. Thompson

C. Schneider

G. Parsons

See next page for additional authors

Follow this and additional works at: <https://newprairiepress.org/kaesrr>



Part of the [Other Animal Sciences Commons](#)

Recommended Citation

Thompson, L.K.; Schneider, C.; Parsons, G.; Miller, K.; Reinhardt, Christopher D.; and Drouillard, James S. (2010) "Initial heifer body composition has little impact on response to Zilmax," *Kansas Agricultural Experiment Station Research Reports*: Vol. 0: Iss. 1. <https://doi.org/10.4148/2378-5977.2890>

This report is brought to you for free and open access by New Prairie Press. It has been accepted for inclusion in Kansas Agricultural Experiment Station Research Reports by an authorized administrator of New Prairie Press. Copyright 2010 Kansas State University Agricultural Experiment Station and Cooperative Extension Service. Contents of this publication may be freely reproduced for educational purposes. All other rights reserved. Brand names appearing in this publication are for product identification purposes only. No endorsement is intended, nor is criticism implied of similar products not mentioned. K-State Research and Extension is an equal opportunity provider and employer.



Initial heifer body composition has little impact on response to Zilmax

Abstract

Using a growth promotant at the correct time of finishing is critical for maximizing profit potential. Previous studies have shown that zilpaterol-HCl (Zilmax; Intervet/ Schering-Plough Animal Health, Millsboro, DE) improves carcass characteristics. The objective of this study was to determine effects of prior body composition on subsequent changes in carcass weight, fatness, and muscle in heifers fed Zilmax so producers can introduce Zilmax at the level of finish that will result in the most desirable response. We hypothesized that fatter heifers use fat as the fuel for muscle growth.

Keywords

Cattlemen's Day, 2010; Kansas Agricultural Experiment Station contribution; no. 10-170-S; Report of progress (Kansas State University. Agricultural Experiment Station and Cooperative Extension Service); 1029; Beef Cattle Research, 2010 is known as Cattlemen's Day, 2010; Beef; Heifer; Zilmax; Growth promotant

Creative Commons License



This work is licensed under a [Creative Commons Attribution 4.0 License](https://creativecommons.org/licenses/by/4.0/).

Authors

L.K. Thompson, C. Schneider, G. Parsons, K. Miller, Christopher D. Reinhardt, and James S. Drouillard

Initial Heifer Body Composition Has Little Impact on Response to Zilmax

L. Thompson, C. Schneider, G. Parsons, K. Miller, C. Reinhardt, and J. Drouillard

Introduction

Using a growth promotant at the correct time of finishing is critical for maximizing profit potential. Previous studies have shown that zilpaterol-HCl (Zilmax; Intervet/Schering-Plough Animal Health, Millsboro, DE) improves carcass characteristics. The objective of this study was to determine effects of prior body composition on subsequent changes in carcass weight, fatness, and muscle in heifers fed Zilmax so producers can introduce Zilmax at the level of finish that will result in the most desirable response. We hypothesized that fatter heifers use fat as the fuel for muscle growth.

Experimental Procedures

Crossbred heifers ($n = 353$; 941 ± 19.5 lb) were fed finishing diets consisting of a combination of steam-flaked corn and processed grain by-products with 3% alfalfa hay and 6% corn silage. Diets provided 300 mg monensin, 90 mg tylosin, and 0.5 mg melengestrol acetate per animal daily. Before initiation of the study, heifers were assigned randomly to experimental diets and pens within weight block. Cattle were placed into 1 of 48 concrete-surfaced pens; there were seven to eight animals per pen. Cattle were fed once daily and had free-choice access to feed and water.

Prior to feeding Zilmax, cattle were weighed, and ribeye area, rump fat thickness, and 12th rib fat thickness were measured by ultrasound. Average daily gains for individual heifers were calculated for the 66-day period preceding initiation of Zilmax feeding, and pre-Zilmax hot carcass weights were estimated by regression using ribeye area, rib fat thickness, and pre-Zilmax body weight. Zilmax was added to finishing diets at 7.56 g/ton (dry matter basis) beginning 23 days before harvest and withdrawn for the final 3 days on feed. Heifers were slaughtered at a commercial abattoir, where carcass data were collected.

Regression formulas were developed to estimate the effects of the means of pre-Zilmax body weight, rump fat thickness, rib fat thickness, ribeye area, and average daily gain on subsequent post-Zilmax changes in 12th rib fat thickness, yield grade, ribeye area, and carcass average daily gain.

Results and Discussion

The following formulas were used to estimate the relationship between initial carcass measurements and changes in fatness or carcass weight that developed during the Zilmax feeding period:

$$\text{Change in 12th rib fat thickness (in inches)} = (0.43 \times \text{rump fat}) - 0.08$$

$$(R^2 = 0.09, P < 0.01).$$

Change in carcass average daily gain (lb/day) = $(0.007 \times \text{pre-Zilmax body weight}) - (0.18 \times \text{ribeye area}) - 3.67$
 $(R^2 = 0.13, P < 0.01)$. This formula is shown in Figure 1.

Yield grade change = $(2.21 \times \text{rump fat}) + (0.002 \times \text{pre-Zilmax body weight}) - (0.10 \times \text{pre-Zilmax average daily gain}) - (0.11 \times \text{ribeye area}) - 2.16$
 $(R^2 = 0.28, P < 0.01)$.

The following formulas illustrate that increases in ribeye areas were more pronounced in leaner, heavier heifers that started with smaller ribeyes:

Change in ribeye area (square inches) = $6.88 - (0.68 \times \text{ribeye area}) - (2.06 \times \text{rump fat}) - (2.96 \times \text{rib fat}) + (0.004 \times \text{pre-Zilmax body weight}) + (0.26 \times \text{pre-Zilmax average daily gain})$
 $(R^2 = 0.58, P < 0.01)$.

Change in carcass efficiency = $(-0.67 \times \text{ribeye area}) - (2.07 \times \text{rump fat}) - (2.93 \times \text{rib fat}) + (0.004 \times \text{pre-Zilmax body weight}) + (0.26 \times \text{pre-Zilmax average daily gain}) + 6.91$
 $(R^2 = 0.58, P < 0.01)$.

Implications

Feeding Zilmax can be an effective management tool for feedlot operators. This study suggests that benefits of Zilmax are similar for lean and fat heifers. Carcass gain efficiency is also similar for lean and fat heifers. Feeding Zilmax to fatter heifers does not elicit a better response that would justify costs of the extended feeding period. Zilmax should be fed to heifers with a typical degree of finish because degree of fatness does not drive the Zilmax response. Formulas from this study may help predict changes in carcass composition when feeding Zilmax to heifers.

NUTRITION




Figure 1. Changes in hot carcass weight as affected by ribeye size and live body weight of feedlot heifers.