Kansas Agricultural Experiment Station Research Reports

Volume 0
Issue 2 *Dairy Research (1984-2014)*

Article 47

1991

Leukocyte function and health status of calves supplemented with vitamins A and E

S. D. Eicher-Pruett

J. L. Morrill

Frank Blecha

See next page for additional authors

Follow this and additional works at: <https://newprairiepress.org/kaesrr>

Part of the [Dairy Science Commons](#)

Recommended Citation

Eicher-Pruett, S. D.; Morrill, J. L.; Blecha, Frank; and Chitko-McKown, C. G. (1991) "Leukocyte function and health status of calves supplemented with vitamins A and E," *Kansas Agricultural Experiment Station Research Reports*: Vol. 0: Iss. 2. <https://doi.org/10.4148/2378-5977.2972>

This report is brought to you for free and open access by New Prairie Press. It has been accepted for inclusion in Kansas Agricultural Experiment Station Research Reports by an authorized administrator of New Prairie Press. Copyright 1991 the Author(s). Contents of this publication may be freely reproduced for educational purposes. All other rights reserved. Brand names appearing in this publication are for product identification purposes only. No endorsement is intended, nor is criticism implied of similar products not mentioned. K-State Research and Extension is an equal opportunity provider and employer.

Leukocyte function and health status of calves supplemented with vitamins A and E

Authors

S. D. Eicher-Pruett, J. L. Morrill, Frank Blecha, and C. G. Chitko-McKown

LEUKOCYTE FUNCTION AND HEALTH STATUS OF CALVES SUPPLEMENTED WITH VITAMINS A AND E

*S. D. Eicher-Pruett, J. L. Morrill, F. Blecha¹,
C. G. Chitko-McKown¹, and N. V. Anderson²*

Summary

Forty-four Holstein calves were fed milk replacers with varied concentrations of vitamins A and E from 3 to 45 d of age to determine their effects on concentrations of plasma vitamin A (retinol and retinyl palmitate) and vitamin E (α -tocopherol), lymphocyte and neutrophil functions, and health of calves. Plasma α -tocopherol was unaffected by increased vitamin A supplementation. Fecal scores, and eye and nose membrane responses were improved with increased vitamin A and lower vitamin E concentration, whereas the same treatment tended to reduce neutrophil cytotoxic and bactericidal activity by 6 wk of age. Increased supplemental vitamin E tended to enhance neutrophil functions. However, age appeared to have an effect on response to both vitamins.

(Key Words: Calves, Leukocytes, Vitamins, Health.)

Introduction

Previous research has shown improved immune function of lymphocytes with increased vitamin E supplementation to young calves. However, research with other species indicated that absorption of α -tocopherol diminished with increased dietary vitamin A, leading to the hypothesis that increased dietary vitamin A may interfere with absorption of dietary vitamin E in the calf. Therefore, vitamin A may limit availability of vitamin E to enhance immune functions. Many milk replacers contain more than 10 times

the NRC requirement of vitamin A and amounts less than or equal to NRC recommendations of vitamin E. This experiment was conducted to determine if 1) increased vitamin A interferes with plasma α -tocopherol concentrations and 2) various concentrations of vitamins A and E in the diet affect lymphocyte and neutrophil functions and other health traits. All concentrations of vitamins that were used reflect concentrations present in milk replacers on the market.

Procedures

Forty-four Holstein calves, blocked by sex and age, were fed colostrum and then transition milk for 3 d. They were then fed experimental milk replacer at 10% of body weight, adjusted weekly. Vitamin A concentrations provided in milk replacers were low (LA; 3,200 IU/lb) or high (HA; 39,900 IU/lb) and vitamin E concentrations were low (LE; 5.1 IU/lb) or high (HE; 25.9 IU/lb). Concentrations of vitamin A and vitamin E reflect those amounts contained in milk replacers. The four experimental milk replacers were designated LA-LE, HA-LE, LA-HE, and HA-HE. Twice daily fecal scores and discharges of eyes and nose were recorded. Calves were weighed weekly. At 0, 3, and 6 wk, blood was sampled for determination of plasma retinol, retinyl palmitate, and α -tocopherol. Blood samples were collected at 3 and 6 wk to determine lymphocyte proliferation and neutrophil cytotoxicity and bactericidal and chemotactic functions (measures of immune health of calves). Concanavalin A was used as the mitogen for lymphocyte proliferation. The cytotoxicity assay was an

¹Department of Anatomy and Physiology.

²Department of Surgery and Medicine.

antibody-dependent cellular-cytotoxicity (ADCC) assay using chicken red blood cells as the target cells. The neutrophil bactericidal assay targeted *Staphylococcus aureus*. Chemotaxis was measured under agarose with zymosan-activated serum as the chemoattractant for directed:random migration.

Results and Discussion

Plasma Vitamin Concentrations

Concentrations of plasma α -tocopherol were not affected adversely by increased supplementation of vitamin A at 6 wk (Table 1) but reflected the supplementation of vitamin E. However, α -tocopherol concentrations tended to increase overall with high vitamin A supplementation and were higher ($P<.05$) at 3 wk. Plasma retinol and retinyl palmitate did not consistently reflect the increased supplementation of vitamin A. Some of the inconsistencies may have been due to a retinol ester that is formed or because of tissue stores (neither measured in our analysis).

Growth and Health

Gain in body weight was similar between treatments for the total 6-wk period (72, 71, 64, and 66 lb for LA-LE, HA-LE, LA-HE, and HA-HE, respectively). The mean fecal score (1=solid, 4=fluid) for the 6-wk period of the HA-LE calves was lower ($P<.10$) than the scores of both LA treatment groups. The HA-LE group tended to have the lowest fecal score at 2 to 5 wk (Figure 1). The increase for the LA-HE group at 2 wk may explain the decrease in gain of that group that occurred then. The eye discharges increased, beginning at 2 wk for all treatments and remained high through 5 wk (Figure 2). The discharges observed in this study were clear,

probably in response to fly irritation. Therefore, an increased discharge was considered a healthy response of the eye membrane. The HA-LE treatment tended to have the greatest occurrence of eye discharges. Total nasal discharges across weeks were greater for the LA-HE treatment ($P<.10$; data not shown). These discharges were thick mucous that occurred in few calves and for short periods of time and were considered a sign of infection.

Leukocyte Function

No differences in lymphocyte blastogenesis occurred among treatments at 3 or 6 wk (Table 2). Neutrophil phagocytosis and bactericidal activity tended to be lowest at 6 wk for calves on HA-LE treatment. Significant differences ($P<.05$) in bactericidal activity occurred between HA-HE and LA-HE treatments at 3 wk. The chemotaxis index indicated a greater response to a chemoattractant at 6 wk for LA-HE-supplemented calves.

Conclusion

Increased supplementation of vitamin A tended to improve responses that rely on a healthy mucous membrane. Simultaneously, the immune functions that utilize vitamin E tended to be improved by increased vitamin E and were inhibited when lower vitamin E and higher vitamin A concentrations were fed. The response of neutrophils to the chemoattractant, although enhanced by HE supplementation, was inhibited when HA was fed simultaneously, indicating possible interference of vitamin A with vitamin E utilization when both are fed at high concentrations. An age effect on vitamin E was seen both in plasma concentrations and leukocyte responses.

Table 1. Plasma Retinol, Retinyl Palmitate, and α -Tocopherol Concentrations in Calves Fed Experimental Milk Replacers

	Vitamin supplementation				SE
	LA-LE	HA-LE	LA-HE	HA-HE	
Vitamin & wk	----- (µg/dl) -----				
α -Tocopherol					
3 wk	266 ^c	255 ^c	298 ^b	354 ^a	7.6
6 wk	285 ^b	297 ^b	439 ^a	452 ^a	8.4
Retinol					
3 wk	101 ^b	95 ^c	102 ^b	109 ^a	2.2
6 wk	72 ^b	191 ^a	89 ^b	77 ^b	11.7
Retinyl Palmitate					
3 wk	51 ^b	55 ^b	52 ^b	63 ^a	3.1
6 wk	50 ^c	66 ^b	72 ^{ab}	84 ^a	5.5

^{a,b,c}Means within a row without a common superscript letter differ (P<.05).

Table 2. Cellular Functions Weeks 3 and 6 of Calves Fed Experimental Milk Replacers

Measurement and Wk	Vitamin supplement				SE
	LA-LE	HA-LE	LA-HE	HA-HE	
Lymphocyte Blastogenesis (CPM)					
3 wk	193244	179062	179798	167536	18715
6 wk	191916	170630	169899	195659	17076
ADCC (%Lysis)					
3 wk	40.1	42.4	35.1	34.5	4.1
6 wk	42.4	37.5	44.8	45.4	6.1
<u>S. aureus</u> (% Kill)					
3 wk	27.1 ^{ab}	20.5 ^b	19.7 ^{bd}	31.7 ^{ac}	6.3
6 wk	24.0	18.8	25.6	26.9	5.6
Chemotaxis Index ¹					
3 wk	3.8	2.5	3.8	3.2	.1
6 wk	4.2 ^{ab}	5.1 ^{ab}	7.9 ^a	4.0 ^b	.3

^{a,b,c,d}Means within row with different superscripts differ (^{ab}P<.10); ^{cd}P<.05).

¹For description of test see test.

Figure 1. Weekly Fecal Scores of Calves Fed Experimental Milk Replacers. Means Within a Week with Different Superscripts Differ ($P < .10$). 1 = Solid to 4 = Fluid.

Figure 2. Weekly Eye Discharges of Calves Fed Experimental Milk Replacers. Means Within a Week with Different Superscripts Differ ($P < .10$).