

Kansas Agricultural Experiment Station Research Reports

Volume 0
Issue 2 *Dairy Research (1984-2014)*

Article 115

1987

Early lactation somatic cell count should be low

James R. Dunham

Follow this and additional works at: <https://newprairiepress.org/kaesrr>



Part of the [Dairy Science Commons](#)

Recommended Citation

Dunham, James R. (1987) "Early lactation somatic cell count should be low," *Kansas Agricultural Experiment Station Research Reports*: Vol. 0: Iss. 2. <https://doi.org/10.4148/2378-5977.3040>

This report is brought to you for free and open access by New Prairie Press. It has been accepted for inclusion in Kansas Agricultural Experiment Station Research Reports by an authorized administrator of New Prairie Press. Copyright 1987 Kansas State University Agricultural Experiment Station and Cooperative Extension Service. Contents of this publication may be freely reproduced for educational purposes. All other rights reserved. Brand names appearing in this publication are for product identification purposes only. No endorsement is intended, nor is criticism implied of similar products not mentioned. K-State Research and Extension is an equal opportunity provider and employer.



Early lactation somatic cell count should be low

Abstract

Cows and heifers in milk for fewer than 50 days, as shown on the DHIA Somatic Cell Count report, should have a lower average Somatic Cell Count than cows in milk for more than 300 days, if the dry cow mastitis treatment and management programs are effective. The DHIA Somatic Cell Count (SCC) average for a dairy herd is a good evaluation of a mastitis control program. Since leucocytes migrate to the mammary system during periods of subclinical mastitis, the SCC average indicates the amount of subclinical mastitis in a herd. A realistic goal for a herd average SCC is less than 200,000. In many herds with average SCC over 400,000, the average SCC of early lactation cows is higher than that of late lactation cows. This indicates a problem during the dry period, either dry cows are not being cured of mastitis or they are becoming re-infected with mastitis during the dry period.; Dairy Day, 1987, Kansas State University, Manhattan, KS, 1987;

Keywords

Kansas Agricultural Experiment Station contribution; no. 88-114-S; Report of progress (Kansas Agricultural Experiment Station); 527; Dairy; Lactation; Somatic cell count; Dry cow

Creative Commons License



This work is licensed under a [Creative Commons Attribution 4.0 License](https://creativecommons.org/licenses/by/4.0/).

K**S****U**

EARLY LACTATION SOMATIC CELL COUNT
SHOULD BE LOW

J.R. Dunham

Summary

Cows and heifers in milk for fewer than 50 days, as shown on the DHIA Somatic Cell Count report, should have a lower average Somatic Cell Count than cows in milk for more than 300 days, if the dry cow mastitis treatment and management programs are effective.

Introduction

The DHIA Somatic Cell Count (SCC) average for a dairy herd is a good evaluation of a mastitis control program. Since leucocytes migrate to the mammary system during periods of subclinical mastitis, the SCC average indicates the amount of subclinical mastitis in a herd. A realistic goal for a herd average SCC is less than 200,000.

In many herds with average SCC over 400,000, the average SCC of early lactation cows is higher than that of late lactation cows. This indicates a problem during the dry period, either dry cows are not being cured of mastitis or they are becoming re-infected with mastitis during the dry period.

Evaluating the Dry Cow — Mastitis Control Program

The DHIA SCC report summarizes the herd on each test day according to the stage of lactation of each cow. This summary is the Days in Milk SCC Average shown in Table 1.

Table 1. Days in milk and Somatic Cell Count average

Days in Milk	SCC Average	
	Your Herd	Mid-States Top 25%
Fresh under 50 days	497,000	178,000
Fresh 50 to 100 days	390,000	159,000
Fresh 101 to 200 days	435,000	166,000
Fresh 201 to 300 days	444,000	183,000
Fresh over 300 days	466,000	206,000

The effectiveness of a dry cow mastitis control program can be evaluated by comparing the SCC average of cows fresh less than 50 days with that cows fresh over 300 days. If this report shows a consistently higher average SCC for early fresh cows, than an evaluation of the dry cow program should include:

1. Drying Cows Off. Cows should be turned dry by discontinuing milking abruptly. Those cows producing more than 40 lb at dry-off should be fed less grain to decrease milk flow.
2. Dry Cow Treatment. Every quarter should be treated with a dry cow preparation shown to be effective by sensitivity tests. Sanitary procedures during treatment must be followed. Cows showing high SCC during late lactation should be re-treated 10-14 days following dry-off.
3. Lots and Housing. The lots and housing facilities must be clean and as dry as possible for springing heifers and dry cows.
4. Prepartum Milking. Springing cows and heifers making up udders that show signs of mastitis should be milked prepartum, and a treatment program should be initiated.

