Kansas Agricultural Experiment Station Research Reports

Volume 0
Issue 2 *Dairy Research (1984-2014)*

Article 158

1984

Effect of processing temperature on utilization of whole soybeans by young calves

I.E.O. Abdelgadir

J.L. Morrill

J.A. Stutts

See next page for additional authors

Follow this and additional works at: <https://newprairiepress.org/kaesrr>

Part of the [Dairy Science Commons](#)

Recommended Citation

Abdelgadir, I.E.O.; Morrill, J.L.; Stutts, J.A.; and Morrill, B. (1984) "Effect of processing temperature on utilization of whole soybeans by young calves," *Kansas Agricultural Experiment Station Research Reports*: Vol. 0: Iss. 2. <https://doi.org/10.4148/2378-5977.3083>

This report is brought to you for free and open access by New Prairie Press. It has been accepted for inclusion in Kansas Agricultural Experiment Station Research Reports by an authorized administrator of New Prairie Press. Copyright 1984 the Author(s). Contents of this publication may be freely reproduced for educational purposes. All other rights reserved. Brand names appearing in this publication are for product identification purposes only. No endorsement is intended, nor is criticism implied of similar products not mentioned. K-State Research and Extension is an equal opportunity provider and employer.

Effect of processing temperature on utilization of whole soybeans by young calves

Authors

I.E.O. Abdelgadir, J.L. Morrill, J.A. Stutts, and B. Morrill

K**S****U**

EFFECT OF PROCESSING TEMPERATURE ON UTILIZATION OF WHOLE SOYBEANS BY YOUNG CALVES¹

I. O. Abdelgadir, J. L. Morrill, J. A. Stutts,
M. B. Morrill, D. E. Johnson²,
and K. C. Behnke³

Summary

Two trials were conducted to determine the optimum conditions for processing whole soybeans for young calves. In the first trial, laboratory procedures and a nitrogen balance study were used. This information was used to design the second trial in which raw and processed soybeans were evaluated in an 8-wk growth trial using 96 Holstein day-old calves. Starters contained soybean meal (SBM), SBM with added fat, raw soybeans, or soybeans processed at 280, 340, or 375°F in a California Pellet Mill Jet-Sploder. Calves fed the starters containing soybeans processed at 340 consumed more feed, gained faster, had lower fecal scores (less scours), and less mortality.

Introduction

Soybeans have a high protein and energy content. However, raw soybeans contain several anti-nutritional factors, which may lower their feed value, especially for young ruminants and nonruminants. Proper heat treatment will destroy these factors and also will improve protein utilization by ruminants by increasing the amount of protein that escapes degradation in the rumen. These experiments were conducted in an attempt to determine the optimal heat treatment for soybeans to be fed to young calves and to evaluate a particular method for performing this treatment.

Procedures

In the first trials, soybeans were heat-treated using various combinations of temperature and time. The processed beans then were subjected to various laboratory tests and to metabolism studies to determine the optimum set of processing conditions.

¹We acknowledge the help of California Pellet Mill, San Francisco, CA in providing financial assistance and of Simonsen Feeds, Quimby, Iowa for processing some of the soybeans.

²Department of Statistics

³Department of Grain Science and Industry

Using the information gained earlier, a growth trial was conducted with 96 Holstein calves. They were fed milk until weaning at 5 wk of age and all of one of 6 calf starters (Table 1) they would consume from birth until 8 wk of age. The 6 starters were alike except for the supplementary protein sources which were either SBM, SBM with added fat, raw soybeans, or soybeans processed at 280, 340, or 375°F. A California Pellet Mill Jet-Sploder was used to process the soybeans. Growth rates, amounts of feed consumed, and various measurements of health were recorded.

TABLE 1. Ingredient and chemical composition of starters^a

Item	Starters ^b					
	SBM	SBM + fat	Raw	280	340	375
	%					
Ingredients						
Corn cobs, ground	9.95	9.97	9.99	10.07	10.10	10.12
Corn, rolled	40.23	40.32	40.39	40.72	40.87	40.95
Oats, rolled	20.12	20.16	20.20	20.36	20.43	20.47
Soybean meal	15.29	15.85	—	—	—	—
Animal fat	—	1.57	—	—	—	—
Soybean, ground, raw	—	—	19.05	—	—	—
Soybean, ground, 280 ^c	—	—	—	18.39	—	—
Soybean, ground, 340 ^c	—	—	—	—	18.1	—
Soybean, ground, 375 ^c	—	—	—	—	—	17.93
Sorghum grain, rolled	8.04	5.75	3.98	4.01	4.03	4.04
Dry molasses	4.66	4.67	4.68	4.72	4.74	4.75
Minerals and vitamins	1.71	1.71	1.71	1.73	1.73	1.74
Chemical analysis						
Dry matter	87.3	87.3	87.9	88.4	86.2	87.6
Crude protein ^d	17.2	15.5	14.3	15.6	15.4	16.6
Ether extract ^d	3.1	3.8	6.2	7.4	5.9	6.5
Crude fiber ^d	8.8	7.8	9.3	7.9	6.1	8.4
Ash ^d	4.9	7.3	5.1	5.2	4.7	4.8
Nitrogen free extract ^d	66.0	65.6	65.1	63.9	67.9	63.7

^aAs fed basis, except as indicated.

^bRefers to supplementary protein sources. SBM=soybean meal; Raw=raw soybeans; 280, 340 and 375 = soybeans processed at those temperatures.

^cIndicates processing temperature.

^dDry matter basis.

Results and Discussion

Results in the earlier trials indicated that the optimum treatment should be setting the Jet-Sploder to produce beans with an exit temperature of 340°F. Therefore, that treatment, a treatment with less heat, and one with more heat were chosen to be evaluated in the growth trial. The results of this trial are in Table 2. Growth rate, feed consumption, and health of calves were improved when the calves were fed starters containing soybeans processed to exit at 340°F. To reach that exit temperature the processing time is about 1 min. These results show that soybeans can be used to supply protein in calf starters, that the soybeans should be heat treated, and recommend the desired conditions for processing those beans. The CPM Jet-Sploder, which is energy-efficient, is satisfactory for processing the beans.

Table 2. Average weekly weight gains, feed consumption, and fecal scores, and mortality in growth study

Treatment ¹	Weight gain (lb)	Starter consumed (lb)	Fecal score ²	Mortality ³
340	7.6 ^a	12.8 ^a	1.58 ^a	0
375	6.6 ^{ab}	12.3 ^a	1.86 ^{ab}	0
280	5.1 ^{bc}	9.7 ^{ab}	1.97 ^b	3
Raw	4.8 ^{bc}	9.6 ^{ab}	2.02 ^b	2
SBM	4.3 ^{bc}	8.8 ^b	2.17 ^b	4
SBM + fat	4.2 ^c	8.4 ^b	2.19 ^b	3

¹ Refers to protein source in starter; see Table 1.

² 1=normal, 2=moderately soft, 3=semi-fluid, 4=watery.

³ Number of calves that died.

^{abc} Means within column with the same letter are not different ($P < .05$ for gain and fecal score, $P < .07$ for starter consumption).