

Kansas Agricultural Experiment Station Research Reports

Volume 0
Issue 10 *Swine Day (1968-2014)*

Article 42

1969

Simple and complex pig starters compared

B A. Koch

Robert H. Hines

Follow this and additional works at: <https://newprairiepress.org/kaesrr>



Part of the [Other Animal Sciences Commons](#)

Recommended Citation

Koch, B A. and Hines, Robert H. (1969) "Simple and complex pig starters compared," *Kansas Agricultural Experiment Station Research Reports*: Vol. 0: Iss. 10. <https://doi.org/10.4148/2378-5977.3462>

This report is brought to you for free and open access by New Prairie Press. It has been accepted for inclusion in Kansas Agricultural Experiment Station Research Reports by an authorized administrator of New Prairie Press. Copyright 1969 Kansas State University Agricultural Experiment Station and Cooperative Extension Service. Contents of this publication may be freely reproduced for educational purposes. All other rights reserved. Brand names appearing in this publication are for product identification purposes only. No endorsement is intended, nor is criticism implied of similar products not mentioned. K-State Research and Extension is an equal opportunity provider and employer.



Simple and complex pig starters compared

Abstract

Baby pigs have the ability to grow rapidly and efficiently. Creep rations at an early age enhance their potential. Most commercial creep rations contain feedstuffs to increase palatability and acceptability. Are these ingredients necessary or will rather simple starters produce similar gains? Five starters were compared, varying from a simple corn-milo- soybean meal ration to a ration containing several palatability aiding ingredients--from 10 days of age to weaning at 28-35 days; and from 6 weeks to 10 weeks of age.; Swine Day, Manhattan, KS, September 25, 1969

Keywords

Swine day, 1969; Report of progress (Kansas State University. Agricultural Experiment Station and Cooperative Extension Service); 151; Swine; Creep rations; Palatability

Creative Commons License



This work is licensed under a [Creative Commons Attribution 4.0 License](https://creativecommons.org/licenses/by/4.0/).

Simple and Complex Pig Starters Compared
R. H. Hines and B. A. Koch

Baby pigs have the ability to grow rapidly and efficiently. Creep rations at an early age enhance their potential. Most commercial creep rations contain feedstuffs to increase palatability and acceptability. Are these ingredients necessary or will rather simple starters produce similar gains?

Five starters were compared, varying from a simple corn-milo-soybean meal ration to a ration containing several palatability aiding ingredients--from 10 days of age to weaning at 28-35 days; and from 6 weeks to 10 weeks of age.

Experimental Procedure

Phase I. Creep rations were assigned randomly. Birth weight of litter was used as the initial weight and litter weaning weight as terminal weight. Rations used are listed in Table 12. All creeps were placed in the same location in the farrowing pens. The first few days that creep was offered, the ration was freshened every other day.

Phase II. One hundred fifty-five, 6-week-old pigs were used to compare starters from 6 to 10 weeks of age. The pigs were assigned to three replicates of the five starters. The pigs were housed in a controlled environment nursery. The 5' x 11' pens are totally slatted with a circulating oxidation ditch below. Each pen was fed ad libitum from a two-hole feeder and an automatic waterer. The pigs were lotted to balance litters and sex in each replicate.

Results and Discussion

Pigs receiving 101C consumed significantly more creep during phase 1 than pigs on any other starter ration (Table 13). Average intake per pig from 10 to 28 days was approximately 1-1½ lbs. Average weaning weights favored 101C and 101E starter rations. They were the medium and most complex rations. Weight gains and feed intake for pigs receiving the simple starter ration were comparable to those on the more expensive starters.

Although starter 101C showed a significant advantage in feed consumption during phase 1, the results reversed during phase 2. Average daily gains on starter 101C and 101A were significantly less than from the other three rations because daily feed intake was significantly less.

Summary

Ingredients that enhance palatability of creep starters may not be of significant value. Weight gains from five starters tested were comparable. Cost of gain during suckling and early post-weaning favor the simple ration. Acceptability of starter rations may be the result of freshness or pellet hardness instead of various appetite stimulators.

Table 12. Simple and complex starter ration formulas used.

Ration complexity ^A	+	++	+++	++++	+++++
	101A	101B	101C	101D	101E
A. Ingredient: lbs.					
Ground corn	665.0	541.0	425.0	---	477.0
Ground sorghum grain	665.0	541.0	425.0	965.5	477.0
Ground oats	---	---	---	200.0	200.0
Rolled oats	---	300.0	300.0	---	---
Alfalfa meal (17%)	---	---	---	40.0	60.0
Sugar	---	---	176.0	80.0	60.0
Soybean meal (50%)	600.0	300.0	260.0	300.0	300.0
Fishmeal	---	---	100.0	60.0	60.0
Meat & bone meal	---	---	---	100.0	100.0
Dried buttermilk	---	---	140.0	---	---
Dried skim milk	---	260.0	140.0	100.0	100.0
Dried whey	---	---	---	100.0	100.0
Edible fat	---	---	---	20.0	20.0
Limestone	20.0	16.0	8.0	---	---
DiCalcium phosphate	28.0	20.0	4.0	10.0	10.0
Salt	10.0	10.0	10.0	10.0	10.0
Trace mineral (5%Zn)	2.0	2.0	2.0	4.0	2.0
B. Premix, gms					
Vit A (10,000 IU/gm)	400.0	400.0	400.0	400.0	600.0
Vit D (15,000 IU/gm)	60.0	60.0	60.0	60.0	80.0
Vit E (44 IU/gm)	---	---	---	---	908.0
Vit K (Klotgen F, 16 Gm/lb)	---	---	---	120.0	120.0
B-Complex (Merck 1233) ^B	908.0	908.0	908.0	908.0	1360.0
Vit B ₁₂ (proferm 20)	908.0	---	908.0	908.0	908.0
Choline chloride (25%)	---	---	---	---	908.0
Aureo SP 250	2272.0	2272.0	2272.0	2272.0	2272.0
Enzyme (zymo-Best)	---	---	---	---	112.0
Monosodium glutamate	---	---	---	---	2724.0
Cost per 100 lbs. ^{D, \$}	4.68	6.91	6.89	6.76	7.18
%protein ^C	20.1	20.7	20.6	20.5	21.7

A--Appraisal of complexity based on number of ingredients used. Simplest ration is + , etc.

B--Contains 80 gms of choline chloride, 24 gms of niacin, 8 gms of riboflavin and 16 gms of D-pantothenic acid per lb.

C--Courtesy of Kansas State University grain science and industry department laboratory.

D--Cost of ration as furnished by Manhattan Milling Company.

Table 13. Phase I: Gains from simple and complex starter rations fed pigs from 10 days of age to weaning.

Starter complexity	+	++	+++	++++	+++++
	101A	101B	101C	101D	101E
No. of litters	9	16	13	16	11
Av. no pigs/litter	9.9	8.6	8.8	8.4	8.5
Av. birth wt./pig, lbs.	3.0	2.9	3.0	2.9	3.0
Av. birth wt./litter, lbs.	29.2	24.7	26.4	24.2	25.7
Av. wean wt/pig, lbs.	14.6	13.8	17.0	15.7	17.2
Av. wean wt./litter, lbs.	144.6	118.4	149.2	132.2	147.0
Av. feed intake/pig, lbs.	.9	.9	1.7	.9	1.1
Av. weaning age, days	30	30	32	31	32

Table 14. Phase II: Gains from simple or complex starters fed 28 days after weaning

Starter complexity	+	++	+++	++++	+++++
	101A	101B	101C	101D	101E
<u>Average initial weight, lbs.</u>					
Rep 1	25.6	27.3	21.9	25.4	26.8
Rep 2	26.2	35.4	32.4	37.8	36.6
Rep 3	24.4	25.4	27.5	26.7	28.3
Av.	28.7	29.3	27.3	29.9	30.6
<u>Average final weight, lbs.</u>					
Rep 1	54.9	64.9	53.6	58.4	62.2
Rep 2	67.8	67.5	58.6	74.1	69.8
Rep 3	51.0	51.1	52.6	53.0	57.9
Av.	57.9	61.2	54.9	61.8	63.3
<u>Average daily gain, lbs.</u>					
Rep 1	1.05	1.34	1.13	1.18	1.26
Rep 2	1.13	1.15	.94	1.30	1.19
Rep 3	.95	.92	.90	.94	1.06
Av.	1.04	1.13	.99	1.14	1.17
<u>Average daily feed intake, lbs.</u>					
Rep 1	1.69	2.29	1.88	2.18	2.33
Rep 2	1.99	2.05	1.52	2.40	2.19
Rep 3	1.63	1.54	1.57	1.68	1.81
Av.	1.77	1.96	1.66	2.08	2.11
<u>Average lbs. feed per lb. of gain</u>					
Rep 1	1.61	1.71	1.66	1.85	1.85
Rep 2	1.76	1.87	1.62	1.85	1.84
Rep 3	1.72	1.67	1.74	1.79	1.71
Av.	1.70	1.75	1.67	1.83	1.80