

Kansas Agricultural Experiment Station Research Reports

Volume 0
Issue 10 *Swine Day (1968-2014)*

Article 156

1975

Yeast culture as an additive in swine grower rations

B A. Koch

G L. Allee

Robert H. Hines

Follow this and additional works at: <https://newprairiepress.org/kaesrr>



Part of the [Other Animal Sciences Commons](#)

Recommended Citation

Koch, B A.; Allee, G L.; and Hines, Robert H. (1975) "Yeast culture as an additive in swine grower rations," *Kansas Agricultural Experiment Station Research Reports*: Vol. 0: Iss. 10. <https://doi.org/10.4148/2378-5977.5996>

This report is brought to you for free and open access by New Prairie Press. It has been accepted for inclusion in Kansas Agricultural Experiment Station Research Reports by an authorized administrator of New Prairie Press. Copyright 1975 Kansas State University Agricultural Experiment Station and Cooperative Extension Service. Contents of this publication may be freely reproduced for educational purposes. All other rights reserved. Brand names appearing in this publication are for product identification purposes only. No endorsement is intended, nor is criticism implied of similar products not mentioned. K-State Research and Extension is an equal opportunity provider and employer.



Yeast culture as an additive in swine grower rations

Abstract

Adding yeast culture at manufacturer's recommended levels to a complete, pelleted, 16% crude protein ration fed to nursery-age pigs produced no measurable improvement in rate of gain, feed efficiency, survival, activity, or appearance.; Swine Day, Manhattan, KS, November 13, 1975

Keywords

Swine day, 1975; Kansas Agricultural Experiment Station contribution; no. 505; Report of progress (Kansas State University. Agricultural Experiment Station and Cooperative Extension Service); 283; Swine; Yeast; Grower rations; Nursery pigs

Creative Commons License



This work is licensed under a [Creative Commons Attribution 4.0 License](https://creativecommons.org/licenses/by/4.0/).

Summary

Adding yeast culture at manufacturer's recommended levels to a complete, pelleted, 16% crude protein ration fed to nursery-age pigs produced no measurable improvement in rate of gain, feed efficiency, survival, activity, or appearance.

Introduction

Commercial products resulting from fermentation of bacteria, molds or yeasts are being widely promoted as supplements in pig diets. Yeast Culture, one such product, is derived from a *Saccharomyces cerevisiae* yeast.

Procedure

Fifty-eight nursery-age pigs averaging 22 pounds were randomly allotted (on the basis of sex, breed, and weight) to 6 pens and fed 28 days on a complete, 16% crude protein, pelleted ration. Forty pounds of Yeast Culture was added to each ton of ration fed to pigs in three pens. Pigs were housed in the controlled-environment, slatted floor nursery at the Swine Research Farm.

Results and Discussion

Results of the feeding trial are summarized in table 29. Adding Yeast Culture to the ration had no measurable effect on average daily gain,

feed efficiency, survival, activity, or appearance of the animals in this 28 day study. The pelleting process may have affected Yeast Culture activity. However, nothing in the manufacturer's recommendations suggested that as a factor.

Table 29. Results from using Yeast Culture as an additive in swine grower rations.

	Control	Fed Yeast Culture
Number of pigs	29	29
Initial wt., lbs.	22	22
Final wt., lbs.	51	51
Avg. daily gain, lbs.	1.01	1.02
Feed/gain ratio	2.12	2.17
Avg. daily intake, lbs.	2.19	2.25