Kansas Agricultural Experiment Station Research Reports

Volume 0
Issue 10 *Swine Day (1968-2014)*

Article 157

1975

Microlite offered free-choice to growing-finishing pigs

B A. Koch

G L. Allee

Robert H. Hines

Follow this and additional works at: <https://newprairiepress.org/kaesrr>

Part of the [Other Animal Sciences Commons](#)

Recommended Citation

Koch, B A.; Allee, G L.; and Hines, Robert H. (1975) "Microlite offered free-choice to growing-finishing pigs," *Kansas Agricultural Experiment Station Research Reports*: Vol. 0: Iss. 10. <https://doi.org/10.4148/2378-5977.5997>

This report is brought to you for free and open access by New Prairie Press. It has been accepted for inclusion in Kansas Agricultural Experiment Station Research Reports by an authorized administrator of New Prairie Press. Copyright 1975 the Author(s). Contents of this publication may be freely reproduced for educational purposes. All other rights reserved. Brand names appearing in this publication are for product identification purposes only. No endorsement is intended, nor is criticism implied of similar products not mentioned. K-State Research and Extension is an equal opportunity provider and employer.

Microlite Offered Free-choice to Growing-finishing Pigs

B. A. Koch, G. L. Allee, and R. H. Hines

Summary

Growing and finishing pigs consumed from 54 to 64 grams of Microlite per head per day when it was available free choice from a self-feeder. It had no measurable affect on average daily gain, feed consumption, feed efficiency, or behavior of pigs eating a complete ration.

Introduction

Microlite, an end product from a smectite-vermiculite mineral deposit strip-mined in southeastern Kansas, is being offered as a supplemental mineral for farm animals. It is guaranteed to contain 10% Mg, 5% Fe, and 2% K plus other mineral nutrients.

Procedure

We offered Microlite to pigs free-choice in two separate trials. In the first trial 48 pigs averaging 67 pounds were randomly allotted (on the basis of sex, breed, and weight) to 6 pens and were fed to an average weight of 204 pounds on a complete, 16% crude protein, pelleted ration. Pigs in 3 pens had access to separate self-feeders containing Microlite.

In the second trial 60 pigs averaging 22 pounds were randomly allotted (on the basis of sex, breed and weight) to 6 pens and were fed for 28 days on the same complete ration, again with 3 pens having access to Microlite free choice.

All pigs were housed in controlled environment, slatted-floor buildings at the University Swine Research Center.

Results and Discussion

Performance data from trials I and II are summarized in table 28. Pigs apparently like the flavor of Microlite. In trial I average consumption per pig per day was 64.7 gms (0.14 pound or approximately 2 1/4 ounces). As pigs were on slatted floors, some mineral may have been wasted unknowingly. It had no measurable affect on daily gain, feed efficiency, or behavior of the pigs.

Younger and smaller pigs were used in trial II. They consumed an average of 54.7 gms. (0.12 pound or approximately 1.9 ounces) per day of Microlite. Again it had no measurable affect on daily gain, feed efficiency, or behavior.

Table 28. Growing and finishing pigs with or without access to a mineral supplement.

Indicated Item	Microlite not available	Microlite avail- able
Growing - Finishing		
Number of pigs	24	23
Initial wt., lbs.	66	67
Final wt., lbs.	204	205
Avg. daily gain, lbs.	1.88	1.86
Feed/gain ratio	2.86	3.00
Avg. daily intake, lbs.	5.39	5.62
Avg. daily mineral intake, gms.	----	64.7
Growing		
Number of pigs	29	29
Initial wt., lbs.	22	22
Final wt., lbs.	51	53
Avg. daily gain, lbs.	1.01	1.10
Feed/gain ratio	2.12	2.03
Avg. daily intake, lbs.	2.19	2.27
Avg. daily mineral intake, gms.	----	54.7