

Kansas Agricultural Experiment Station Research Reports

Volume 0
Issue 10 *Swine Day (1968-2014)*

Article 173

1978

Teat necrosis in newborn gilts

D A. Schoneweis

Follow this and additional works at: <https://newprairiepress.org/kaesrr>



Part of the [Other Animal Sciences Commons](#)

Recommended Citation

Schoneweis, D A. (1978) "Teat necrosis in newborn gilts," *Kansas Agricultural Experiment Station Research Reports*: Vol. 0: Iss. 10. <https://doi.org/10.4148/2378-5977.6013>

This report is brought to you for free and open access by New Prairie Press. It has been accepted for inclusion in Kansas Agricultural Experiment Station Research Reports by an authorized administrator of New Prairie Press. Copyright 1978 Kansas State University Agricultural Experiment Station and Cooperative Extension Service. Contents of this publication may be freely reproduced for educational purposes. All other rights reserved. Brand names appearing in this publication are for product identification purposes only. No endorsement is intended, nor is criticism implied of similar products not mentioned. K-State Research and Extension is an equal opportunity provider and employer.



Teat necrosis in newborn gilts

Abstract

Marked enlargement and edema of the vulva are often noted in newborn gilts. An accompanying, but less conspicuous, lesion is enlarged, engorged mammary glands. Although not fully understood, the changes are thought to result from increased estrogen in sows before they farrow. The edema of the vulva usually causes no lasting problems but enlarged mammary glands are more susceptible to injury from a rough surface like concrete slats. Such injuries may lead to teat necrosis and fibrosis, which makes nipples appear inverted and they are often nonfunctional when the gilts farrow. The incidence of teat necrosis can be reduced by carpeting or other materials that protect pigs underlines from rough surfaces. Applying a protectant like Knee-Kote® to the underline of the gilts shortly after they are born and again whenever they are handled the first few days of life may further reduce the incidence.; Swine Day, Manhattan, KS, November 9, 1978

Keywords

Swine day, 1978; Kansas Agricultural Experiment Station contribution; no. 79-105-S; Report of progress (Kansas State University. Agricultural Experiment Station and Cooperative Extension Service); 342; Swine; Teat necrosis; Newborn gilts; Farrow

Creative Commons License



This work is licensed under a [Creative Commons Attribution 4.0 License](https://creativecommons.org/licenses/by/4.0/).

Marked enlargement and edema of the vulva are often noted in newborn gilts. An accompanying, but less conspicuous, lesion is enlarged, engorged mammary glands. Although not fully understood, the changes are thought to result from increased estrogen in sows before they farrow.

The edema of the vulva usually causes no lasting problems but enlarged mammary glands are more susceptible to injury from a rough surface like concrete slats. Such injuries may lead to teat necrosis and fibrosis, which makes nipples appear inverted and they are often nonfunctional when the gilts farrow.

The incidence of teat necrosis can be reduced by carpeting or other materials that protect pig underlines from rough surfaces. Applying a protectant like Knee-Kote^R to the underline of the gilts shortly after they are born and again whenever they are handled the first few days of life may further reduce the incidence.