

Kansas Agricultural Experiment Station Research Reports

Volume 0
Issue 10 *Swine Day (1968-2014)*

Article 174

1978

Arthrogryposis in piglets

H W. Leipold

R Ely

D Schoneweis

Follow this and additional works at: <https://newprairiepress.org/kaesrr>



Part of the [Other Animal Sciences Commons](#)

Recommended Citation

Leipold, H W.; Ely, R; and Schoneweis, D (1978) "Arthrogryposis in piglets," *Kansas Agricultural Experiment Station Research Reports*: Vol. 0: Iss. 10. <https://doi.org/10.4148/2378-5977.6014>

This report is brought to you for free and open access by New Prairie Press. It has been accepted for inclusion in Kansas Agricultural Experiment Station Research Reports by an authorized administrator of New Prairie Press. Copyright 1978 the Author(s). Contents of this publication may be freely reproduced for educational purposes. All other rights reserved. Brand names appearing in this publication are for product identification purposes only. No endorsement is intended, nor is criticism implied of similar products not mentioned. K-State Research and Extension is an equal opportunity provider and employer.



This report documents experimental reproduction of tetramelic arthrogryposis in purebred Yorkshire swine by breeding trials. Data from the trials indicated arthrogryposis may be due to homozygosity of a simple autosomal recessive gene.

No affected pig was observed to be alive at parturition, although most had been alive in utero just prior to delivery. Other than dystocia observed in the sows, no other significant clinical findings were noted. The condition was noted in five litters from two sows which were bred to the same boar. Approximately 25% of the pigs exhibited the clinical signs of arthrogryposis which included malformed rigidly extended legs.

Gross pathological findings were tetramelic arthrogryposis of all joint articulations and kyphosis of the thoracic vertebra. In those cases, severing the overlying tendons and muscles would result in normal joint movement. The articular

surfaces of the affected joints appeared flattened and lacked normal remodelling of the cartilaginous surfaces. The subcutaneous fascia and skeletal muscles were edematous and lacked fascial tissue separating muscle tissue into individual bundles. An occasional dislocated flexor involving the hind-legs was noted. The skeletal muscle was hypoplastic in the affected limbs. Examining other body organs revealed no gross abnormalities.

¹Dept. of Pathology.

²Dept. of Surgery & Medicine