

1993

## Comparison of feed-grade antibiotics in starter diets containing spray-dried blood products

K Q. Owen

Steven S. Dritz

Michael D. Tokach

*See next page for additional authors*

Follow this and additional works at: <https://newprairiepress.org/kaesrr>



Part of the [Other Animal Sciences Commons](#)

---

### Recommended Citation

Owen, K Q.; Dritz, Steven S.; Tokach, Michael D.; Goodband, Robert D.; and Nelssen, Jim L. (1993) "Comparison of feed-grade antibiotics in starter diets containing spray-dried blood products," *Kansas Agricultural Experiment Station Research Reports*: Vol. 0: Iss. 10. <https://doi.org/10.4148/2378-5977.6369>

This report is brought to you for free and open access by New Prairie Press. It has been accepted for inclusion in Kansas Agricultural Experiment Station Research Reports by an authorized administrator of New Prairie Press. Copyright 1993 Kansas State University Agricultural Experiment Station and Cooperative Extension Service. Contents of this publication may be freely reproduced for educational purposes. All other rights reserved. Brand names appearing in this publication are for product identification purposes only. No endorsement is intended, nor is criticism implied of similar products not mentioned. K-State Research and Extension is an equal opportunity provider and employer.



---

## Comparison of feed-grade antibiotics in starter diets containing spray-dried blood products

### Abstract

A total of 240 pigs (initially 11.6 lb and 19 d of age) was used to compare four different feed-grade antibiotics or antibiotic combinations in phase I (d 0 to 14 postweaning) and phase II (d 14 to 28 postweaning) starter pig diets. Apramycin and carbadox were compared in the phase I diet. Combinations of tylosin/sulfamethazine and neomycin sulfate/oxytetracycline were compared in the phase II diet. No differences occurred in pig performance for the phase I and phase II periods between the feed-grade antibiotics compared in this growth assay. Therefore, determination of appropriate feed-grade antibiotic inclusion will depend on 1) economics, 2) disease profile of the herd, and 3) growth response within a particular producer's herd.; Swine Day, Manhattan, KS, November 18,1993

### Keywords

Swine day, 1993; Kansas Agricultural Experiment Station contribution; no. 94-194-S; Report of progress (Kansas State University. Agricultural Experiment Station and Cooperative Extension Service); 695; Swine; Starter; Performance; Antibiotic

### Creative Commons License



This work is licensed under a [Creative Commons Attribution 4.0 License](https://creativecommons.org/licenses/by/4.0/).

### Authors

K Q. Owen, Steven S. Dritz, Michael D. Tokach, Robert D. Goodband, and Jim L. Nelssen

## COMPARISON OF FEED-GRADE ANTIBIOTICS IN STARTER DIETS CONTAINING SPRAY-DRIED BLOOD PRODUCTS<sup>1</sup>

*S. S. Dritz, M. D. Tokach, R. D. Goodband,  
J. L. Nelssen, and K. Q. Owen*

### Summary

A total of 240 pigs (initially 11.6 lb and 19 d of age) was used to compare four different feed-grade antibiotics or antibiotic combinations in phase I (d 0 to 14 postweaning) and phase II (d 14 to 28 postweaning) starter pig diets. Apramycin and carbadox were compared in the phase I diet. Combinations of tylosin/sulfamethazine and neomycin sulfate/oxytetracycline were compared in the phase II diet. No differences occurred in pig performance for the phase I and phase II periods between the feed-grade antibiotics compared in this growth assay. Therefore, determination of appropriate feed-grade antibiotic inclusion will depend on 1) economics, 2) disease profile of the herd, and 3) growth response within a particular producer's herd.

(Key Words: Starter, Performance, Antibiotic.)

### Introduction

Numerous growth-promoting antibiotics are available for addition to starter pig diets. Producers are continually faced with the challenge of deciding which feed-grade antibiotics are appropriate for their swine business. This decision often is based on nontechnical information and testimonials rather than research data. Therefore, this trial had three objectives 1) to compare apramycin and carbadox inclusion in phase

I (d 0 to 14 postweaning), 2) to compare a tylosin/sulfathiazole combination and a neomycin/oxytetracycline combination in phase II (d 14 to 28 postweaning), and 3) to determine the appropriate sequence of feed grade antibiotic usage in phase I and phase II starter pig diets containing spray-dried blood products.

### Procedures

A total of 240 pigs (initially 11.6 lb and 19 d of age) was used in a 28 d growth trial. The trial consisted of two phases. Phase I was from d 0 to 14 postweaning, and phase II was from d 14 to 28 postweaning. Pigs were allotted by weight and placed in 4 ft by 6 ft pens (10 pigs per pen) in an environmentally controlled nursery facility on a commercial farm in northeast Kansas. Each pen contained two nipple waterers, and pigs were allowed ad libitum access to feed and water. Pigs were assigned to one of four treatments based on the sequence of feed-grade antibiotic inclusion. Either 150 g/ton apramycin or 50 g/ton carbadox were included in the phase I diet and a combination of either 100 g/ton tylosin/100 g/ton sulfamethazine (Tylan-sulfa) or 100 g/ton neomycin sulfate/100 g/ton oxytetracycline (Neo-terra) in the phase II diet. Phase I diets were formulated to contain 1.5% lysine, .42% methionine, .9% calcium, and .8% phosphorus. The phase I diets contained 7.5% spray-dried porcine plasma, 1.75% spray-dried blood

---

<sup>1</sup>Appreciation is expressed to Elanco for financial support and Dr. T. A. Marsteller for technical assistance. The authors also wish to thank Steve and Brent Eichman, St. George, KS for use of facilities and animals.

meal, and 20% dried whey. Phase II diets were formulated to contain 1.25% lysine, .34% methionine, .9% calcium, and .8% phosphorus. The phase II diets contained 10% dried whey and 2.5% spray-dried blood meal. Phase I diets were fed in pellet form, and phase II diets were fed in meal form. Pigs and feeders were weighed on d 7, 14, 21, and 28 to evaluate ADG, ADFI, and F/G.

### Results and Discussion

No mortality occurred during the 28 d growth period. During phase I, no differences occurred in ADG, ADFI, and F/G between pigs fed apramycin or carbadox

(Table 3). Phase I feed-grade antibiotic did not influence phase II performance, and no interactions occurred between feed-grade antibiotic sources fed during phases I and II (Table 5). In addition, no differences occurred in performances between pigs fed tylosin/sulfamethazine or neomycin/oxytetracycline in phase II (Table 4). Because of the similar growth performance, results of this trial indicate that the choice between the feed-grade antibiotics used in this study will depend on economics and risk or perceived risk of tissue residue. In conclusion, this growth assay provides information to swine producers to aid in the selection of feed-grade antibiotics in starter pig diets.

**Table 1. Diet Composition**

Item	Phase I	Phase II
Corn	45.29	59.26
Soybean meal, (48% CP)	16.13	21.26
Dried whey, edible grade	20.00	10.00
Spray-dried porcine plasma	7.50	—
Spray-dried blood meal	1.75	2.50
Soybean oil	5.00	3.00
Monocalcium phosphate	1.91	1.97
Limestone	.69	.83
Antibiotic	1.00	.50
Trace mineral premix	.15	.15
Vitamin premix	.25	.25
DL-methionine	.15	.05
L-lysine HCl	.10	.15
Copper sulfate	.075	.075
Total	100	100

**Table 2. Feed-Grade Antibiotic Inclusion Levels**

Item	Inclusion level, g/ton
<b>Phase I</b>	
Apramycin	150
Carbadox	50
<b>Phase II</b>	
Tylosin/sulfamethazine	100/100
Neomycin sulfate/oxytetracycline	100/100

**Table 3. Phase I Performance (d 0 to 14 Postweaning)<sup>ab</sup>**

Item	Apramycin	Carbadox	CV
ADG, lb	.57	.57	10.4
ADFI, lb	.64	.63	6.7
F/G	1.13	1.12	6.9

<sup>a</sup>Each value is the mean of 12 pens containing 10 pigs per pen. Pigs were initially 19 d of age and 11.6 lb.

<sup>b</sup>No significant treatment effects.

**Table 4. Phase II Performance (d 14 to 28 Postweaning)<sup>ab</sup>**

Item	Tylosin/ sulfamethazine	Neomycin sulfate/ oxytetracycline	CV
ADG, lb	.92	.93	8.7
ADFI, lb	1.45	1.44	4.9
F/G	1.58	1.57	6.4

<sup>a</sup>Each value is the mean of twelve pens containing 10 pigs per pen.

<sup>b</sup>No significant treatment effects.

**Table 5. Growth Performance d 0 to 28 Postweaning<sup>ab</sup>**

Item	Apramycin		Carbadox		CV
	Tylosin/ Sulfa <sup>c</sup>	Neo/ oxy <sup>d</sup>	Tylosin/ Sulfa	Neo/ oxy	
ADG, lb	.91	.90	.94	.96	8.7
ADFI, lb	1.47	1.41	1.43	1.46	4.9
F/G	1.63	1.59	1.53	1.54	6.4

<sup>a</sup>Diets were fed in two phases: Phase I (d 0 to 14 postweaning) and Phase II (d 14 to 28 postweaning). Each value is the mean of six pens containing 10 pigs per pen. Pigs were initially 19 d of age and 11.6 lb.

<sup>b</sup>No significant treatment effects.

<sup>c</sup>Tylosin/sulfamethazine.

<sup>d</sup>Neomycin sulfate/oxytetracycline.