

Kansas Agricultural Experiment Station Research Reports

Volume 0
Issue 10 *Swine Day (1968-2014)*

Article 725

1998

Effects of sorghum starch type, endosperm hardness, and processing on digestibility and growth performance in finishing pigs and chicks

H Cao

Robert H. Hines

B W. Senne

See next page for additional authors

Follow this and additional works at: <https://newprairiepress.org/kaesrr>



Part of the [Other Animal Sciences Commons](#)

Recommended Citation

Cao, H; Hines, Robert H.; Senne, B W.; Froetschner, J R.; Jiang, J M.; Johnston, S L.; Hancock, Joe D.; and Behnke, Keith C. (1998) "Effects of sorghum starch type, endosperm hardness, and processing on digestibility and growth performance in finishing pigs and chicks," *Kansas Agricultural Experiment Station Research Reports*: Vol. 0: Iss. 10. <https://doi.org/10.4148/2378-5977.6565>

This report is brought to you for free and open access by New Prairie Press. It has been accepted for inclusion in Kansas Agricultural Experiment Station Research Reports by an authorized administrator of New Prairie Press. Copyright 1998 Kansas State University Agricultural Experiment Station and Cooperative Extension Service. Contents of this publication may be freely reproduced for educational purposes. All other rights reserved. Brand names appearing in this publication are for product identification purposes only. No endorsement is intended, nor is criticism implied of similar products not mentioned. K-State Research and Extension is an equal opportunity provider and employer.



Effects of sorghum starch type, endosperm hardness, and processing on digestibility and growth performance in finishing pigs and chicks

Abstract

In finishing pigs, waxy sorghum had lower digestibilities for DM and GE than the other genotypes. Also, the hard sorghums were more digestible ($P < .06$) than the medium hardness sorghum. In broiler chicks, the waxy sorghum was similar to the other genotypes for growth performance and nutrient digestibility. The soft sorghum was superior to the medium and hard genotypes for nutrient retention but not for growth performance. Fine grinding improved F/G and increased retention of nutrients and steam-flaked sorghum supported greater growth performance than extruded sorghum.; Swine Day, Manhattan, KS, November 19, 1998

Keywords

Swine day, 1998; Kansas Agricultural Experiment Station contribution; no. 99-120-S; Report of progress (Kansas State University. Agricultural Experiment Station and Cooperative Extension Service); 819; Swine; Sorghum; Processing; Chicks; Finishing pigs

Creative Commons License



This work is licensed under a [Creative Commons Attribution 4.0 License](https://creativecommons.org/licenses/by/4.0/).

Authors

H Cao, Robert H. Hines, B W. Senne, J R. Froetschner, J M. Jiang, S L. Johnston, Joe D. Hancock, and Keith C. Behnke

K

S

**EFFECTS OF SORGHUM STARCH TYPE, ENDOSPERM
HARDNESS, AND PROCESSING ON DIGESTIBILITY
AND GROWTH PERFORMANCE IN FINISHING
PIGS AND CHICKS¹**

U

*H. Cao, J. D. Hancock, R. H. Hines,
K. C. Behnke², B. W. Senne, J. R. Froetschner²,
J. M. Jiang, and S. L. Johnston*

Summary

In finishing pigs, waxy sorghum had lower digestibilities for DM and GE than the other genotypes. Also, the hard sorghums were more digestible ($P < .06$) than the medium hardness sorghum. In broiler chicks, the waxy sorghum was similar to the other genotypes for growth performance and nutrient digestibility. The soft sorghum was superior to the medium and hard genotypes for nutrient retention but not for growth performance. Fine grinding improved F/G and increased retention of nutrients and steam-flaked sorghum supported greater growth performance than extruded sorghum.

(Key Words: Sorghum, Processing, Chicks, Finishing Pigs.)

Introduction

Previous research from our laboratory indicated that processing technologies and (or) endosperm hardness impact growth performance of chicks and pigs fed sorghums. Additionally, waxy sorghum is more susceptible to enzymes during digestion. However, little is known about potential interactions among endosperm type and processing technologies that might optimize nutrient digestibility and growth performance in swine and poultry. Therefore, experiments were designed to determine the effect of endosperm hardness and starch type on nutrient digestibility in pigs and to investi-

gate the interactions among genotype and processing technology in broiler chicks.

Procedures

Six sorghum genotypes were grouped into soft (851111), medium (279 & PL-1), hard (Segolane & 475), and waxy (739) categories. For the pig experiment, the sorghums were ground to a particle size of 550 μm . For the chick assay, the sorghums were (coarsely ground to 1,000 μm & finely ground to 450 μm) and thermally processed (steam-flaked at 150°F and extruded at 235°F).

For the pig experiment, five barrows (average initial BW of 142 lb) were used in a 5 \times 5 Latin square design. The pigs were housed in metabolism cages placed in an environmentally controlled building (70°F). The pigs were fed three times per day (7:00 a.m., 1:00 p.m., and 7:00 p.m.). After a 3-d adjustment period, fresh feces were collected twice each day dried; ground; and analyzed for dry matter, crude protein, gross energy, and chromium. All diets were formulated to .6% lysine, .55% of Ca, and .50% P (Table 1), with the ground sorghums as the only source of energy and protein (Table 1).

In Exp.2, 600 broiler chicks (avg initial BW of 90 g) were used in a 21-d growth assay with 24 treatments (6 genotypes \times 4 processing, and 5 pens/trt and 5 chicks/pen). The chicks were housed in brooder batteries and allowed ad libitum access to feed and

¹Appreciation is extended to the Kansas State Board of Agriculture and the Kansas State Sorghum Commission for funding this project.

²Department of Grain Science and Industry.

water. Feces (with urine) were collected during the last 2 d of the growth assay; dried; ground; and analyzed for DM, CP, GE, and Cr to determine nutrient retention. All diets were sorghum-soybean meal-based with 1.0% tallow and formulated to 1.32% lysine, 1.1% Ca and 0.9% P (Table 1).

Results and Discussion

Chemical analyses of the grains (Table 2) suggested that they were "normal" for moisture (8 to 12%) and crude protein (9 to 11%). However, notable differences occurred among the sorghums for texture and starch type, which allowed us to categorize the genotypes. Sorghum genotypes had less effect on energy consumption during processing than the processing technologies themselves, with thermal processing using more energy than grinding, and extrusion using more energy than steam flaking (Table 3). Energy consumption was similar among the genotypes, except for the low kwh/ton needed to coarsely and finely grind and steam flake Segolane. Also, waxy sorghum required the least energy to steam flake, suggesting a strong genotype \times processing procedure interaction.

In finishing pigs, digestibilities of DM ($P < .01$) and GE ($P < .07$) were less for waxy sorghums than the other genotypes (Table 4). Also, the hard sorghums were more digestible ($P < .06$) than the medium hardness sorghums. Both of these effects were contrary to "conventional wisdom", which would support greater nutrient utilization from waxy and soft endosperm types.

In the broiler experiment (Table 5), no genotype effects on growth performance were observed. Chicks fed soft sorghum had greater ($P < .01$) retentions of DM, N, and GE compared to chicks fed medium and hard endosperm genotypes, whereas those fed

hard sorghums had greater nutrient retention than those fed medium hardness sorghums ($P < .02$).

Grinding improved ADG ($P < .01$), ADFI ($P < .001$), and F/G ($P < .04$) when compared to thermal processing. However, thermal processing improved retention of DM, N, and GE ($P < .03$) compared to grinding. Retentions of DM, CP, and GE were improved with fine grinding versus coarse grinding ($P < .05$) and steam flaking versus extrusion ($P < .06$).

Interactions among genotypes and processing technologies were observed in feed efficiency. Soft sorghum responded more favorably ($P < .05$) to coarse grinding than to fine grinding compared to medium and hard types, and hard sorghum responded more favorably ($P < .01$) to flaking than to extrusion. As for nutrient retention, soft sorghum responded more favorably ($P < .01$) to thermal processing than to grinding compared to harder sorghums. Harder sorghum responded more favorably ($P < .01$) to flaking than to extrusion compared to medium sorghums. Waxy sorghums responded more positively to fine grinding ($P < .04$) and flaking ($P < .06$) than to coarse grinding and extrusion compared to nonwaxy sorghum.

In conclusion, retention of DM and GE for finishing pigs was greater with hard endosperm than soft and waxy sorghums. Feed efficiency was greatest in chicks fed the hard sorghum and steam-flaked treatments. Retentions of DM, N, and ME were greatest for chicks fed soft genotypes and thermally processed sorghums. However, within thermal processing, waxy endosperm sorghum responded more favorably to steam flaking and less favorably to extrusion, whereas soft and medium sorghums showed the opposite response.

Table 1. Compositions of Basal Diets with Exotic Sorghum Genotypes (Exps. 1 and 2)^a

Ingredient, %	Finisher (Exp. 1)	Chick (Exp. 2)
Sorghum	96.30	53.30
Soybean meal (46.5 % CP)	--	39.80
Tallow	--	1.00
Monocalcium phosphate	1.10	2.30
Limestone	.85	1.50
Salt	.30	.50
Lysine HCl	.40	--
Threonine	.16	.10
D, L-methionine	.11	.26
Tryptophan	.12	--
Vitamin premix	.15	.23
Mineral premix	.10	.28
Sow add pack	.05	.26
Chromium	.25	.20
Copper sulfate	--	.06
Antibiotics ^b	.125	.25

^aFinisher diets were formulated to .6% lysine, .55% Ca, and .5% P. Chick diets were formulated to 1.32% lysine, 1.1% Ca, and .9% P.

^bFinisher diets had 40 g/ton tylosin; chick diets had 100 g/ton chlortetracycline and .0125% amprolium.

Table 2. Characteristics of Exotic Sorghum Genotypes

Item	Soft	Medium		Hard		Waxy
	851171	279	PL-1	475	Segolane	739
Physical traits						
Pericap color	white	red	yellow	cream	cream	cream
Endosperm color	white	white	yellow	white	white	white
Texture ^a	soft	medium	medium	hard	hard	soft
Starch type ^b	normal	normal	normal	normal	normal	waxy
Chemical analyses						
Unprocessed						
Moisture, %	10.8	8.5	10.4	9.5	10.6	11.8
CP, %	11.2	11.3	10.8	10.1	10.0	8.8
Extruded						
Moisture, %	4.4	4.5	4.8	5.5	4.7	5.3
CP, %	12.6	12.2	12.7	10.7	11.5	10.8
Steam-flaked						
Moisture, %	8.9	8.6	11.1	9.9	8.8	11.9
CP, %	10.5	11.1	10.5	10.1	10.0	10.3

^aTexture was determined using the Single Kernel Characterization (SKC) method.

^bStarch type was determined by visual appraisal.

Table 3. Processing Energy Consumption of Sorghum Genotypes, kwh/t

Item	Soft	Medium		Hard		Waxy	Processing Mean
	851171	279	PL-1	475	Segolane	739	
Coarse ^a	11.9	13.9	10.8	10.8	8.3	13.6	11.5
Fine ^b	10.3	12.1	12.3	8.2	8.2	15.4	11.1
Flaked ^c	60.4	57.3	50.6	50.6	40.8	36.1	49.3
Extruded ^d	76.0	74.2	72.5	72.1	76.4	66.6	70.2
Genotype mean	39.7	39.4	36.5	35.4	33.4	32.9	

^aGround in a roller mill to a mean particle size of 1,039 μm .

^bGround in a roller mill to a mean particle size of 440 μm .

^cSteam-flaked at 150°F.

^dExtruded at 235°F.

Table 4. Effect of Sorghum Genotype on Nutrient Digestibility in Finishing Pigs^a

Digestibility, %	Soft	Medium	Hard		Waxy	SE	Contrast ^b			
	851171	PL-1	Segolane	475	739		1	2	3	4
DM	83.8	84.3	86.5	87.2	82.2	.9	-- ^c	.06	--	.01
N	69.7	65.6	67.0	68.3	69.5	2.3	--	--	--	--
GE	83.3	83.4	85.8	85.6	82.5	.9	--	.06	--	.07

^aFive finishers were used (avg initial BW of 160 lb).

^bContrast were: 1) soft vs medium & hard; 2) medium vs hard; 3) hard vs hard; and 4) waxy vs others.

^cDashes indicated $P > .1$.

Table 5. Effects of Sorghum Genotypes and Processing on Growth Performance in Chicks^a

Item	Soft	Medium		Hard		Waxy	Processing Mean
	851171	279	PL-1	475	Segolane	739	
ADG, g ^b							
Coarse	39.1	35.7	40.2	37.7	39.4	38.8	38.5
Fine	37.7	38.4	37.9	39.5	39.0	39.1	38.6
Flaked	37.8	37.7	33.1	39.1	39.7	37.9	37.4
Extrude	33.1	34.2	36.0	35.6	29.0	34.3	33.9
Genotype mean	36.9	36.5	36.8	38.0	36.9	37.5	SE 1.1
ADFI, g ^c							
Coarse	56.3	57.2	63.0	57.3	60.6	56.8	58.5
Fine	57.0	56.1	53.5	59.2	55.4	58.2	56.6
Flaked	58.5	57.8	51.1	58.3	55.0	58.0	56.4
Extruded	52.3	50.1	51.4	57.5	49.0	56.6	53.0
Genotype mean	56.1	55.2	54.7	58.1	55.3	57.4	SE 2.1
F/G ^d							
Coarse	1.43	1.46	1.56	1.52	1.54	1.47	1.52
Fine	1.51	1.53	1.41	1.50	1.42	1.49	1.46
Flaked	1.55	1.46	1.54	1.49	1.40	1.54	1.50
Extruded	1.58	1.56	1.42	1.62	1.70	1.65	1.56
Genotype mean	1.51	1.51	1.48	1.53	1.50	1.53	SE .02

^aA total of 600 chicks was used (five chicks/pen and five pen/trt) with an avg initial BW of 90 g. ^bHard vs hard ($P < .06$); ground vs thermally processed ($P < .01$); flake vs extruded ($P < .01$).

^cGround vs thermally processed ($P < .001$); flake vs extruded ($P < .01$); 475 vs Segolane \times ground vs thermally processed ($P < .08$); medium vs hard \times flake vs extrusion ($P < .01$). ^dGround vs thermally processed ($P < .04$); coarse vs fine ($P < .07$); flaked vs extruded ($P < .07$); soft vs medium & hard \times coarse vs fine ground ($P < .05$); medium vs hard \times flaked vs extruded ($P < .001$).

Table 6. Effects of Sorghum Genotypes and Processing on Nutrient Retention in Chicks^a

Item ^c	Soft	Medium		Hard		Waxy	Processing Mean
	851171	279	PL-1	475	Segolane	739	
DM, % ^b							
Coarse	73.7	73.9	73.0	75.3	76.9	71.4	74.0
Fine	75.5	78.7	77.2	75.0	76.9	77.5	77.0
Flaked	80.2	67.2	75.6	81.1	71.3	79.5	76.1
Extrude	80.3	72.7	78.1	75.8	80.1	77.8	77.4
Genotype mean	77.7	73.5	76.0	76.8	76.1	76.6	SE 1.1
N, % ^c							
Coarse	58.4	57.79	56.2	59.9	62.5	53.73	58.1
Fine	61.7	63.2	62.4	57.4	59.79	63.1	61.2
Flaked	67.4	42.4	58.5	69.1	46.7	64.7	58.7
Extruded	71.4	54.6	60.8	57.1	67.1	62.3	62.0
Genotype mean	64.7	55.1	59.5	60.9	58.6	61.0	SE 2.3
GE, % ^d							
Coarse	77.2	78.4	76.2	78.5	80.9	74.8	77.7
Fine	80.1	82.3	81.3	79.0	80.9	81.0	80.8
Flaked	83.8	73.2	80.0	84.8	76.4	83.3	80.5
Extruded	84.4	77.6	82.0	79.6	83.6	81.7	81.4
Genotype mean	81.4	78.1	79.9	80.5	80.3	80.2	SE 1.0

^aA total of 600 chicks was used (five chicks/pen and five pen/trt) with an avg initial BW of 90 g.

^bSoft vs medium & hard (P<.01); medium vs hard (P<.01); ground vs thermally processed (P<.03), coarse vs fine (P<.001); flaked vs extruded (P<.02); soft vs medium & hard × thermal processed (P<.001); ; medium vs hard × ground vs thermally processing (P <.01); medium vs hard × coarse vs fine ground (P<.01).

^cSoft vs medium & hard (P<.001); coarse vs fine (P<.05); flaked vs extruded (P<.06); waxy vs others × ground vs thermally processing (P<.05); waxy vs others × coarse vs fine ground (P<.09); waxy vs others × flaked and extruded (P<.04).

^dSoft vs medium & hard (P<.01); medium vs hard (P<.01); ground vs thermally processed (P<.001), coarse vs fine (P<.001); flaked vs extruded (P<.08), soft vs medium & hard × ground vs thermal processed (P<.001); medium vs hard × ground vs thermally processing (P<.001); medium vs hard × coarse vs fine ground (P<.03); waxy vs others × ground vs thermally processing (P<.06); waxy vs others × coarse vs fine ground (P<.04); waxy vs others × flaked and extruded (P<.01).