

Kansas Agricultural Experiment Station Research Reports

Volume 0
Issue 10 *Swine Day (1968-2014)*

Article 747

2000

Effects of free fatty acid concentrations in choice white grease on growth performance and nutrient digestibility in weanling pigs

Robert H. Hines

H Cao

C A. Maloney

See next page for additional authors

Follow this and additional works at: <https://newprairiepress.org/kaesrr>



Part of the [Other Animal Sciences Commons](#)

Recommended Citation

Hines, Robert H.; Cao, H; Maloney, C A.; Dean, D W.; Lee, D J.; Park, J S.; DeRouchey, Joel M.; and Hancock, Joe D. (2000) "Effects of free fatty acid concentrations in choice white grease on growth performance and nutrient digestibility in weanling pigs," *Kansas Agricultural Experiment Station Research Reports*: Vol. 0: Iss. 10. <https://doi.org/10.4148/2378-5977.6587>

This report is brought to you for free and open access by New Prairie Press. It has been accepted for inclusion in Kansas Agricultural Experiment Station Research Reports by an authorized administrator of New Prairie Press. Copyright 2000 Kansas State University Agricultural Experiment Station and Cooperative Extension Service. Contents of this publication may be freely reproduced for educational purposes. All other rights reserved. Brand names appearing in this publication are for product identification purposes only. No endorsement is intended, nor is criticism implied of similar products not mentioned. K-State Research and Extension is an equal opportunity provider and employer.



Effects of free fatty acid concentrations in choice white grease on growth performance and nutrient digestibility in weanling pigs

Abstract

As much as 53% free fatty acids in choice white grease did not adversely affect piglet performance. Thus, concentration of free fatty acids, if they are not otherwise damaged or rancid, is not an acceptable measure of fat quality.; Swine Day, Manhattan, KS, November 16, 2000

Keywords

Swine day, 2000; Kansas Agricultural Experiment Station contribution; no. 01-138-S; Report of progress (Kansas State University. Agricultural Experiment Station and Cooperative Extension Service); 858; Swine; Weanling pigs; Fat quality; Free fatty acids

Creative Commons License



This work is licensed under a [Creative Commons Attribution 4.0 License](https://creativecommons.org/licenses/by/4.0/).

Authors

Robert H. Hines, H Cao, C A. Maloney, D W. Dean, D J. Lee, J S. Park, Joel M. DeRouchey, and Joe D. Hancock

EFFECTS OF FREE FATTY ACID CONCENTRATIONS IN CHOICE WHITE GREASE ON GROWTH PERFORMANCE AND NUTRIENT DIGESTIBILITY IN WEANLING PIGS

*J. M. DeRouchey, J. D. Hancock, R. H. Hines,
H. Cao, C. A. Maloney, D. W. Dean,
D. J. Lee, and J. S. Park*

Summary

As much as 53% free fatty acids in choice white grease did not adversely affect piglet performance. Thus, concentration of free fatty acids, if they are not otherwise damaged or rancid, is not an acceptable measure of fat quality.

(Key Words: Weanling Pigs, Fat Quality, Free Fatty Acids)

Introduction

Using fat to improve growth performance in nursery pigs is a common practice. However, research about fat utilization in young pigs has been focused on the effects of unsaturated:saturated fatty acid ratios, chain length of the fatty acids, and age of pigs when fat is added to the diet. Thus, few data are available about the effects of "fat quality" (e.g., free fatty acids) and growth performance in pigs. Our objective was to determine the effects of free fatty acids in choice white grease on growth performance and nutrient digestibility in weanling pigs.

Procedures

One hundred twenty five weanling pigs (lines C 22 × 326, PIC, Frankl'n KY) with an average initial BW of 13.7 lb were used in a 33-d growth assay. The pigs were weaned and allotted by BW, sex, and ancestry to five treatments. Initial BW was used as the blocking criterion. The diets (Table 1) were formulated to 1.70% lysine for d 0 to 5, 1.55% lysine for d 5 to 19, and 1.40% lysine for d 19 to 33. They also met or exceeded all nutrient requirements as defined by NRC (1998). Treatments were a corn-soybean

meal-based diet with no added fat; 6% choice white grease; and 6% choice white grease that had been heated at 95°F and treated with 872, 1,175, and 2,248 lipase units/g of fat. The choice white grease allotted for lipase addition was divided in three barrels (55 gal capacity) with 65 lb per barrel, and heaters were attached to maintain a constant temperature. Fat treated with lipase was mixed with water, the lipase was added, and the mixture was agitated for 12 h. The mixture was allowed to stand for 24 h (for separation of the fat and water), and the water was pumped from the bottom of the barrel.

Pigs for the experiment were housed in an environmentally controlled building with a self-feeder and nipple water in each pen to allow ad libitum consumption of feed and water. Pigs and feeders were weighed on d 5, 19, and 33 to allow calculation of ADG, ADFI, and gain/feed.

Pen was the experimental unit. The GLM procedures of SAS were used for all statistical analyses. Polynomial regression was used to determine the shape of the response to increasing free fatty acids in the choice white grease.

Results and Discussion

Chemical analyses of the fat treatments indicated that percentage of free fatty acids increased (from 2 to 53%) as lipase addition to the fat was increased from none to 2,248 lipase units/g of fat (Table 2). Peroxide and p-anisidine values and percentage insoluble impurities and unsaponifiable matter were unchanged as lipase treatment was increased. Also, fatty acid composition, iodine value, and U/S ratio remained virtually unchanged

as indicated by enzyme hydrolysis of the choice white grease. However, moisture increased as the extent of lipase treatment was increased, indicating that less water separation from the choice white grease was possible as free fatty acids increased.

In the growth assay (Table 3), no differences in ADG, ADFI, or F/G were detected from d 0 to 5 ($P>.30$) or d 5 to 19 ($P>.11$). However, for d 19 and 33, ADG tended to increase ($P<.06$) and F/G decreased ($P<.01$) with the fat-added treatments versus the control.

Overall, ADG was not affected ($P>.15$), but F/G was improved ($P<.04$) with the

addition of fat to the diet. No differences occurred in ADG or F/G as concentration of free fatty acids increased ($P>.16$). Also, no differences were observed in digestibility of DM or GE among the dietary treatments ($P>.14$). Also, the digestibility of long-chain unsaturated fatty acids, long-chain saturated fatty acids, and total fat were greater ($P<.001$) for the fat-added treatments compared to control, but fatty acid digestibility was not affected ($P>.17$) by concentration of free fatty acids in the diet.

In conclusion, as much as 53% free fatty acids in choice white grease did not adversely affect piglet performance. Thus, concentration of free fatty acids, if they are not otherwise damaged or rancid is not an acceptable measure of fat quality.

Table 1. Compositions of Diets^a

Item, %	d 0 to 5		d 5 to 19		d 19 to 33	
	Control	Fat Added	Control	Fat Added	Control	Fat Added
Corn	22.84	22.84	44.10	44.10	47.81	47.81
Soybean meal (46.5% CP)	26.36	26.36	31.52	31.52	41.86	41.86
Dried whey	20.00	20.00	10.00	10.00	—	—
Lactose	10.00	10.00	—	—	—	—
Cornstarch	6.00	—	6.00	—	6.00	—
Choice white grease ^b	—	6.00	—	6.00	—	6.00
Spray-dried wheat gluten	4.00	4.00	—	—	—	—
Spray-dried plasma protein	4.00	4.00	1.00	1.00	—	—
Fish meal (menhaden)	2.00	2.00	3.00	3.00	—	—
Monocalcium phosphate (21%)	1.28	1.28	.85	.85	1.32	1.32
Limestone	.89	.89	.84	.84	1.17	1.17
L-lysine·HCl	.32	.32	.25	.25	.05	.05
D,L-methionine	.19	.19	.14	.14	.05	.05
L-threonine	.08	.08	.10	.10	—	—
Salt	.25	.25	.30	.30	.35	.35
Vitamin premix	.25	.25	.25	.25	.25	.25
Mineral premix	.15	.15	.15	.15	.15	.15
Antibiotic ^c	1.00	1.00	1.00	1.00	1.00	1.00
Zinc oxide	.32	.32	.25	.25	—	—
Copper sulfate	—	—	—	—	.09	.09
Chromic oxide	—	—	.25	.25	—	—

^aDiets were formulated to: 1.7% lysine, .9% Ca, and .8% P for d 0 to 5; 1.55% lysine, .8% Ca, and .7% P for d 5 to 19; and 1.4% lysine, .8% Ca, and .7% P for d 19 to 33.

^bChoice white grease with 0, 872, 1,752, and 2,248 lipase units/gram of fat.

^cProvided 150 g of apramycin per ton of feed for d 0 to 19 and 50 g of carbox per ton of feed for d 19 to 33.

Table 2. Chemical Analysis of Choice White Grease

Item	Lipase Units/G of Choice White Grease			
	0	872	1,752	2,248
Free fatty acids, %	2	18	35	53
Peroxide value, mEq/kg	1	1	1	1
p-anisidine value	5.2	4.9	5.0	5.6
Total M.I.U., % ^a	1.17	2.56	2.85	3.34
Moisture, %	.14	1.58	1.85	2.38
Insoluble impurities, %	.01	.01	.03	.02
Unsaponifiable matter, %	1.02	.97	.97	.94
Iodine value	61.2	61.2	60.5	62.7
Unsaturated:saturated ratio	1.49	1.50	1.44	1.52
Fatty acids, % of sample ^b				
C8:0 ^c	.00	.00	.00	.00
C10:0	.00	.00	.00	.00
C12:0	.26	.24	.24	.23
C14:0	2.07	2.12	2.11	2.03
C16:0	25.48	25.48	25.78	25.12
C16:1	3.23	3.33	3.19	3.40
C18:0	11.80	11.63	12.31	11.73
C18:1	48.06	48.02	47.11	47.28
C18:2	7.33	7.31	7.33	8.14
C18:3	.57	.57	.63	.77

^aMoisture, insoluble impurities, and unsaponifiable matter.

^bFatty acids determined by gas chromatography as derivatized fatty acid methyl esters.

^cNumber of carbon atoms and double bonds designated to the left and right of colon, respectively.

Table 3. Effects of Free Fatty Acid Concentration on Growth Performance and Digestibility of Nutrients in Weanling Pigs^a

Item	No Added Fat	Free Fatty Acid Concentration, %					SE	Probability			
								Free Fatty Acid Effects			
		2	18	35	53	No Added Fat vs Others		Linear	Quadratic	Cubic	
d 0 to 5											
ADG, lb	.58	.59	.62	.61	.65	.04	— ^c	—	—	—	
ADFI, lb	.55	.56	.59	.58	.60	.04	—	—	—	—	
F/G	.95	.95	.96	.96	.92	.05	—	—	—	—	
d 5 to 19											
ADG, lb	.86	.84	.87	.81	.83	.04	—	—	—	—	
ADFI, lb	1.32	1.24	1.29	1.28	1.28	.03	.11	—	—	—	
F/G	1.53	1.47	1.48	1.58	1.54	.05	—	—	—	—	
d 19 to 33											
ADG, lb	1.39	1.49	1.44	1.44	1.50	.03	.06	—	.11	—	
ADFI, lb	2.31	2.13	2.20	2.24	2.23	.06	.11	—	—	—	
F/G	1.66	1.43	1.53	1.56	1.49	.05	.01	—	.09	—	
d 0 to 33											
ADG, lb	1.04	1.08	1.07	1.05	1.09	.02	—	—	—	—	
ADFI, lb	1.62	1.51	1.57	1.58	1.58	.02	.02	.04	—	—	
F/G	1.56	1.40	1.47	1.50	1.45	.04	.04	—	—	—	
Apparent Digestibility, %											
DM	79.9	77.3	78.1	80.0	79.1	1.0	—	.14	—	—	
GE	77.6	75.174.6	76.9	78.6	76.3	1.6	—	—	—	—	
N	76.4	73.3	76.3	78.5	76.0	1.3	—	—	.10	—	
Total fat ^b	66.4		74.1	77.7	74.7	2.2	.001	—	—	—	
Long-chain unsaturated		77.9									
fatty acids	69.7		79.2	81.7	79.4	2.2	.001	—	—	—	
Long-chain saturated	51.0	62.8									
fatty acids	91.8	89.4	63.3	68.9	65.2	3.0	.001	—	—	—	
Medium-chain fatty acids			89.9	91.5	89.4	.9	.11	—	—	—	

^aA total of 125 pigs (five pigs per pen and five pens per treatment) with an average initial BW of 13.7 lb.

^bDetermined by gas chromatography as derivatized fatty acid methyl esters.

^cP>.15.