

1999

Effects of pelleting and pellet conditioning temperatures on weanling pig performance

M U. Steidinger

Michael D. Tokach

P R. O'Quinn

See next page for additional authors

Follow this and additional works at: <https://newprairiepress.org/kaesrr>



Part of the [Other Animal Sciences Commons](#)

Recommended Citation

Steidinger, M U.; Tokach, Michael D.; O'Quinn, P R.; Woodworth, J C.; McKinney, Leland J.; Borg, B S.; Campbell, J M.; Goodband, Robert D.; Nelssen, Jim L.; and Dritz, Steven S. (1999) "Effects of pelleting and pellet conditioning temperatures on weanling pig performance," *Kansas Agricultural Experiment Station Research Reports*: Vol. 0: Iss. 10. <https://doi.org/10.4148/2378-5977.6652>

This report is brought to you for free and open access by New Prairie Press. It has been accepted for inclusion in Kansas Agricultural Experiment Station Research Reports by an authorized administrator of New Prairie Press. Copyright 1999 the Author(s). Contents of this publication may be freely reproduced for educational purposes. All other rights reserved. Brand names appearing in this publication are for product identification purposes only. No endorsement is intended, nor is criticism implied of similar products not mentioned. K-State Research and Extension is an equal opportunity provider and employer.



Effects of pelleting and pellet conditioning temperatures on weanling pig performance

Authors

M U. Steidinger, Michael D. Tokach, P R. O'Quinn, J C. Woodworth, Leland J. McKinney, B S. Borg, J M. Campbell, Robert D. Goodband, Jim L. Nelssen, and Steven S. Dritz

K

EFFECTS OF PELLETING AND PELLET CONDITIONING TEMPERATURES ON WEANLING PIG PERFORMANCE¹

S

*M. U. Steidinger, R. D. Goodband, M. D. Tokach², S. S. Dritz³,
J. L. Nelssen, P. R. O'Quinn, J. C. Woodworth, L. McKinney,*

U

B. S. Borg⁴, and J. M. Campbell⁴

Summary

Two studies were conducted to evaluate the effects of pelleting and pellet conditioning temperature of diets containing 5% spray-dried animal plasma (SDAP) on weanling pig growth performance. In Exp. 1, conditioning temperatures evaluated were 140, 150, 160, and 170°F. In Exp. 2, pellet conditioning temperatures were 140, 155, 170, 185, and 200°F. The results suggest that pellet conditioning temperatures above 170°F decrease weanling pig performance from d 0 to 7 after weaning. Pellet conditioning temperature should not exceed 170°F (exit temperature of 180°F) for nursery diets containing 5% SDAP.

(Key Words: Weanling Pigs, Spray-Dried Animals Plasma, Pellet Conditioning Temperature.)

Introduction

Numerous trials have demonstrated improved ADG and F/G in pigs fed pelleted diets versus meal diets. Because starter diets typically contain specialty ingredients, such as dried whey and spray-dried animal plasma (SDAP), making a high quality pellet can be challenging. Some heat and steam are required in the pelleting process, but too much heat could potentially burn or scorch some of the specialty ingredients and reduce their nutritional value. Therefore, the objective of

this experiment was to determine the effects of pellet conditioning temperature of a starter diet containing SDAP and spray-dried whey.

Procedures

General. Two 28 d growth assays were used to determine the effects of pelleting and pellet conditioning temperatures of starter diets fed to weanling pigs. Pigs (PIC L326 × C22) were housed in an environmentally regulated nursery at the Kansas State University Swine Teaching and Research Center in 5 ft × 5 ft pens and were provided ad libitum access to feed and water. Both experiments used seven pens (replications) per treatment, and each pen contained six pigs. The trials were divided into two phases, with the experimental diets fed from d 0 to 14 after weaning (Table 1). From d 14 to 28, a common corn-soybean meal-based diet containing 10 % dried whey and 2.5 % dried blood cells was fed in a meal form (Table 1).

The pelleted diets were conditioned with a 10-second retention time and then pelleted (5/32 in. diameter) using a Master Model HD 1000 series California Pellet Mill equipped with a die of 1.25 in. effective thickness. Samples of the pelleted diets were collected to determine pellet exit temperatures and pellet durability index (PDI).

Pigs were weighed and feed disappearance was determined on d 0, 7, 14, 21, and

¹The authors thank American Protein Corp. for providing the spray-dried animal plasma and partial financial support.

²Northeast Area Extension Office, Manhattan, KS.

³Food Animal Health and Management Center.

⁴American Protein Corp., Ames, IA.

28 after weaning to calculate ADG, ADFI, and F/G. Data were analyzed in a randomized complete block design with pen as the experimental unit.

Experiment 1. Two hundred fifty-two weanling pigs (13.2 lb and 21 d of age) were used. Experimental treatments consisted of a corn-soybean meal control diet (1.4% lysine and 15% dried whey) with no spray-dried animal plasma (SDAP) fed in a meal form or the control diet with 5% SDAP replacing soybean meal on an equal lysine basis also fed in a meal form. Additional treatments were the 5% SDAP diet pelleted at conditioning temperatures of 140, 150, 160, or 170°F. Corresponding exit temperatures were 155, 166, 171, and 176°F, and corresponding PDI's were 95.7, 92.7, 92.9, and 92.6, respectively.

Experiment 2. Two hundred fifty-two weanling pigs (13.8 lb and 22 d of age) were used in the 28 d growth assay. Experimental treatments were the control diet (containing no SDAP) from Exp. 1 pelleted at a conditioning temperature of 140°F or one of five diets containing 5% SDAP pelleted at conditioning temperatures of 140, 155, 170, 185, or 200°F. Corresponding exit temperatures were 158, 158, 169, 180, 185, and 200°F with PDI's of 94.9, 96.7, 95.2, 93.7, 94.9, and 96.1.

Results and Discussion

Experiment 1. From d 0 to 7 and d 0 to 14, pigs fed the control diet had decreased ($P<.08$) ADG, ADFI, and F/G compared to the mean of pigs fed diets containing 5% SDAP. Although not significant, among pigs fed diets containing SDAP from d 0 to 7, those fed the pelleted diets had an 8% increase in ADG and 12% better feed efficiency than pigs fed the diet in meal form. From day 0 to 14, pellet conditioning temperature had no effect on pig performance.

From d 14 to 28, pigs previously fed the control diet had increased ($P<.02$) ADG and better F/G than animals previously

containing 5% SDAP. From d 14 to 28, feed efficiency improved as pellet conditioning temperature was increased in the diet previously fed from d 0 to 14 (linear, $P<.08$). Overall (d 0 to 28), no effects of feeding SDAP or pellet conditioning temperature used from d 0 to 14 were observed (Table 2).

Experiment 2. From d 0 to 7, pigs fed the 5% SDAP diet conditioned at 140°F tended ($P<.12$) to have greater ADG and greater ($P<.05$) ADFI compared to pigs fed the control diet without SDAP. From d 0 to 7, ADG was similar as pellet conditioning temperature increased to 170°F, but then decreased with conditioning temperatures above 170°F (quadratic, $P<.03$). From d 0 to 7, increasing conditioning temperature also decreased feed intake (linear, $P<.002$), with the most pronounced decrease observed after 170°F. From d 0 to 14, increasing pellet conditioning temperature had no effect on ADG or ADFI, but F/G improved with increasing conditioning temperature (linear, $P<.02$), with pigs fed diets conditioned at 170°F having the best F/G.

From d 14 to 28, pigs previously fed the control diet had improved ($P<.08$) feed efficiency compared to pigs fed the diet containing 5% SDAP conditioned at 140°F. Pellet conditioning temperature of the diet fed from d 0 to 14 had no effect on performance from d 14 to 28 (Table 3).

In conclusion, the results of these experiments suggest that diet preparation and formulation are critical for optimal performance in early-weaned pigs. Results also suggest that adding SDAP to weanling pig starter diets, whether pelleted or prepared in meal form, improves growth performance from d 0 to 7. Although no statistical differences in growth performance occurred between pigs fed meal diets and those fed pelleted diets, pigs fed pelleted diets demonstrated a 12% improvement in feed efficiency from d 0 to 7. Results also indicated that conditioning diets containing SDAP at temperatures above 170°F decreases growth performance of weanling pigs.

Table 1. Compositions of Experimental Diets

| Ingredient, % | Day 0 to 14 ^a | | d 14 to 28 ^b |
|---------------------------|--------------------------|---------|-------------------------|
| | Control | 5% SDAP | |
| Corn | 44.64 | 51.47 | 53.79 |
| Soybean meal (46.5% C.P.) | 34.04 | 22.18 | 25.86 |
| Spray-dried animal plasma | -- | 5.00 | -- |
| Dried whey | 15.00 | 15.00 | 10.00 |
| Soy oil | 3.00 | 3.00 | 3.0 |
| Monocalcium P (21% P) | 1.40 | 1.32 | 1.89 |
| Limestone | .95 | 1.00 | 0.81 |
| Salt | .30 | .30 | 0.25 |
| Vitamin premix | .25 | .25 | 0.25 |
| L-lysine HCl | .15 | .15 | 0.15 |
| Trace mineral | .15 | .15 | 0.15 |
| DL-methionine | .10 | .13 | 0.10 |
| Zinc oxide | -- | -- | 0.25 |
| Medication ^c | -- | -- | 1.00 |
| Spray-dried blood cells | -- | -- | 2.50 |

^aFormulated to contain 1.4% lysine, 0.9% Ca, and 0.8% P.

^bFed in meal form to all pigs and formulated to contain 1.35% lysine, .85% Ca, and .75% P.

^cProvided 50 g/ton carbodox.




Table 2. Effects of Pelleting and Increasing Pelleting Temperature on Weanling Pig Performance (Exp. 1)^a

| Item | Meal Diets | | Pellet Conditioning Temperature (°F) | | | | SEM | Contrast (P<) _c | | | |
|------------------------|------------|---------|--------------------------------------|------|------|------|------|----------------------------|-----|-----|-----|
| | Control | 5% SDAP | 140 | 150 | 160 | 170 | | 1 | 2 | 3 | 4 |
| Treatment ^b | 1 | 2 | 3 | 4 | 5 | 6 | | | | | |
| PDI ^d | - | - | 95.7 | 92.7 | 92.9 | 92.6 | | | | | |
| Day 0 to 7 | | | | | | | | | | | |
| ADG, lb | .31 | .43 | .48 | .47 | .48 | .45 | .028 | .001 | .21 | .53 | .65 |
| ADFI, lb | .52 | .59 | .59 | .58 | .55 | .56 | .023 | .05 | .37 | .25 | .56 |
| F/G | 1.86 | 1.42 | 1.28 | 1.24 | 1.17 | 1.27 | .12 | .001 | .19 | .87 | .55 |
| Day 0 to 14 | | | | | | | | | | | |
| ADG, lb | .42 | .56 | .55 | .60 | .55 | .53 | .026 | .001 | .94 | .39 | .24 |
| ADFI, lb | .73 | .82 | .78 | .80 | .76 | .77 | .027 | .08 | .15 | .55 | .80 |
| F/G | 1.75 | 1.49 | 1.44 | 1.34 | 1.39 | 1.45 | .053 | .001 | .15 | .67 | .16 |
| Day 14 to 28 | | | | | | | | | | | |
| ADG, lb | 1.42 | 1.36 | 1.36 | 1.32 | 1.33 | 1.37 | .025 | .02 | .56 | .62 | .13 |
| ADFI, lb | 1.81 | 1.82 | 1.84 | 1.81 | 1.79 | 1.80 | .041 | .98 | .82 | .41 | .69 |
| F/G | 1.28 | 1.34 | 1.36 | 1.38 | 1.35 | 1.31 | .021 | .01 | .71 | .08 | .21 |
| Day 0 to 28 | | | | | | | | | | | |
| ADG, lb | .92 | .96 | .95 | .96 | .94 | .95 | .022 | .42 | .38 | .31 | .97 |
| ADFI, lb | 1.27 | 1.32 | 1.31 | 1.30 | 1.28 | 1.27 | .029 | .42 | .38 | .31 | .96 |
| F/G | 1.39 | 1.38 | 1.38 | 1.36 | 1.36 | 1.35 | .020 | .34 | .44 | .30 | .95 |

^aTwo hundred fifty-two weanling pigs were used (initially 13.2 lb) with six pigs per pen and seven pens per treatment.

^bTreatment 1 contained no spray-dried animal plasma (SDAP). Treatments 2, 3, 4, 5, and 6 contained 5% SDAP.

^cContrasts: 1=Control (1) vs Plasma (Mean of 2, 3, 4, 5, and 6). 2=Meal (2) vs Pellet (Mean of 3, 4, 5, and 6).

3=Linear effect of conditioning temperature (3, 4, 5, and 6). 4=Quadratic effect of conditioning temperature (3, 4, 5, and 6).

^dPellet Durability Index (Standard: ASAE S269.3).

Table 3. Effects of Increasing Pellet Conditioning Temperature on Weanling Pig Performance (Exp. 2)^a

| Item | Control ^b | Pellet Conditioning Temperature (°F) | | | | | SEM | Contrast (P<) ^c | | |
|------------------|----------------------|--------------------------------------|------|------|------|------|------|----------------------------|------|-----|
| | | 140 | 155 | 170 | 185 | 200 | | 1 | 2 | 3 |
| PDI ^d | 94.9 | 96.7 | 95.2 | 93.7 | 94.9 | 96.1 | | | | |
| Day 0 to 7 | | | | | | | | | | |
| ADG, lb | .51 | .57 | .57 | .60 | .52 | .47 | .025 | .12 | .004 | .03 |
| ADFI, lb | .53 | .59 | .58 | .59 | .55 | .49 | .021 | .05 | .002 | .12 |
| F/G | 1.05 | 1.05 | 1.01 | .99 | 1.05 | 1.08 | .040 | .99 | .41 | .21 |
| Day 0 to 14 | | | | | | | | | | |
| ADG, lb | .52 | .52 | .51 | .57 | .55 | .54 | .021 | .88 | .20 | .29 |
| ADFI, lb | .65 | .69 | .68 | .69 | .69 | .67 | .023 | .23 | .64 | .60 |
| F/G | 1.25 | 1.35 | 1.35 | 1.22 | 1.24 | 1.24 | .042 | .23 | .02 | .36 |
| Day 14 to 28 | | | | | | | | | | |
| ADG, lb | 1.38 | 1.38 | 1.33 | 1.37 | 1.38 | 1.38 | .038 | .86 | .58 | .63 |
| ADFI, lb | 1.82 | 1.88 | 1.80 | 1.87 | 1.87 | 1.94 | .058 | .39 | .34 | .31 |
| F/G | 1.32 | 1.37 | 1.36 | 1.36 | 1.35 | 1.40 | .026 | .08 | .44 | .30 |
| Day 0 to 28 | | | | | | | | | | |
| ADG, lb | .95 | .95 | .92 | .97 | .97 | .96 | .025 | .88 | .32 | .92 |
| ADFI, lb | 1.23 | 1.28 | 1.24 | 1.28 | 1.28 | 1.30 | .038 | .28 | .54 | .53 |
| F/G | 1.30 | 1.36 | 1.35 | 1.32 | 1.32 | 1.36 | .023 | .10 | .64 | .19 |

^aTwo hundred fifty weanling pigs were used (initially 13.8 lb) with six pigs per pen and seven pens per treatment.

^bControl contained no spray-dried animal plasma and was pelleted at 140°F. All other diets contained 5% SDAP and were pelleted at their respective temperatures.

^cContrasts: 1=Control vs 140°F. 2=Linear effect of conditioning temperature. 3=Quadratic effect of conditioning temperature.

^dPellet Durability Index (Standard:ASAE S269.3).