

Kansas Agricultural Experiment Station Research Reports

Volume 0
Issue 10 *Swine Day (1968-2014)*

Article 1020

2005

Effects of increasing dietary lysine on growth performance of pigs fed ractopamine HCl (paylean®)

C R. Neill

C W. Hastad

J L. Usry

See next page for additional authors

Follow this and additional works at: <https://newprairiepress.org/kaesrr>



Part of the [Other Animal Sciences Commons](#)

Recommended Citation

Neill, C R.; Hastad, C W.; Usry, J L.; Tokach, Michael D.; Goodband, Robert D.; Nelssen, Jim L.; DeRouchey, Joel M.; and Dritz, Steven S. (2005) "Effects of increasing dietary lysine on growth performance of pigs fed ractopamine HCl (paylean®)," *Kansas Agricultural Experiment Station Research Reports*: Vol. 0: Iss. 10. <https://doi.org/10.4148/2378-5977.6860>

This report is brought to you for free and open access by New Prairie Press. It has been accepted for inclusion in Kansas Agricultural Experiment Station Research Reports by an authorized administrator of New Prairie Press. Copyright 2005 Kansas State University Agricultural Experiment Station and Cooperative Extension Service. Contents of this publication may be freely reproduced for educational purposes. All other rights reserved. Brand names appearing in this publication are for product identification purposes only. No endorsement is intended, nor is criticism implied of similar products not mentioned. K-State Research and Extension is an equal opportunity provider and employer.



Effects of increasing dietary lysine on growth performance of pigs fed ractopamine HCl (paylean®)

Abstract

A total of 919 gilts (PIC L337 Ã– C22) were used in a 21-d trial conducted in a commercial research facility to determine growth and carcass effects of ractopamine HCl (Paylean®; 6.5 g/ton) and different levels of lysine. The diets were formulated to contain 0.75, 0.85, 0.95, 1.05, 1.15, and 1.25% true ileal digestible (TID) lysine to determine the lysine requirement for pigs fed ractopamine. These TID lysine levels correspond to 0.86, 0.97, 1.08, 1.19, 1.29, and 1.40% total lysine. From d 0 to 14, pigs fed 1.05% TID lysine had the greatest improvement in ADG and F/G. In the overall (d 0 to 21) data, however, pigs fed 1.15% TID lysine had the greatest improvement in ADG and F/G. Average daily gain increased (linear, $P<0.005$), whereas there were no differences in ADFI ($P>0.05$). Feed efficiency also improved (linear, $P<0.005$; quadratic $P<0.07$). Although the ADG and F/G responses were linear, there was very little improvement observed beyond 1.15% TID lysine. Percentage lean increased (linear, $P<0.03$) and FFLI tended to increase (linear, $P<0.07$) with increasing levels of TID lysine. The lean premium increased (linear, $P<0.05$) and backfat tended (linear, $P<0.08$) to improve, but other carcass criteria were not affected. Therefore, pigs fed ractopamine required between 1.05 and 1.15% TID lysine (1.19 to 1.29% total lysine) to maximize growth performance.; Swine Day, 2005, Kansas State University, Manhattan, KS, 2005

Keywords

Swine day, 2005; Summary Publication of Report of Progress (Kansas State University. Agricultural Experiment Station and Cooperative Extension Service); 964; Kansas Agricultural Experiment Station contribution; no. 06-63-S; Carcass parameters; Lysine; Paylean; Swine

Creative Commons License



This work is licensed under a [Creative Commons Attribution 4.0 License](https://creativecommons.org/licenses/by/4.0/).

Authors

C R. Neill, C W. Hastad, J L. Usry, Michael D. Tokach, Robert D. Goodband, Jim L. Nelssen, Joel M. DeRouchey, and Steven S. Dritz

EFFECTS OF INCREASING DIETARY LYSINE ON GROWTH PERFORMANCE OF PIGS FED RACTOPAMINE HCl (PAYLEAN®)¹

C. R. Neill, S. S. Dritz², M. D. Tokach, R. D. Goodband, J. L. Nelssen, J. M. DeRouchey, C. W. Hastad, and J. L. Usry³

Summary

A total of 919 gilts (PIC L337 × C22) were used in a 21-d trial conducted in a commercial research facility to determine growth and carcass effects of ractopamine HCl (Paylean®; 6.5 g/ton) and different levels of lysine. The diets were formulated to contain 0.75, 0.85, 0.95, 1.05, 1.15, and 1.25% true ileal digestible (TID) lysine to determine the lysine requirement for pigs fed ractopamine. These TID lysine levels correspond to 0.86, 0.97, 1.08, 1.19, 1.29, and 1.40% total lysine. From d 0 to 14, pigs fed 1.05% TID lysine had the greatest improvement in ADG and F/G. In the overall (d 0 to 21) data, however, pigs fed 1.15% TID lysine had the greatest improvement in ADG and F/G. Average daily gain increased (linear, $P < 0.005$), whereas there were no differences in ADFI ($P > 0.05$). Feed efficiency also improved (linear, $P < 0.005$; quadratic $P < 0.07$). Although the ADG and F/G responses were linear, there was very little improvement observed beyond 1.15% TID lysine. Percentage lean increased (linear, $P < 0.03$) and FFLI tended to increase (linear, $P < 0.07$) with increasing levels of TID lysine. The lean premium increased (linear, $P < 0.05$) and backfat tended (linear, $P < 0.08$) to improve, but other carcass criteria were not affected. Therefore, pigs fed ractopamine re-

quired between 1.05 and 1.15% TID lysine (1.19 to 1.29% total lysine) to maximize growth performance.

(Key Words: Carcass Parameters, Lysine, Paylean.)

Introduction

In December 1999, ractopamine HCl (Paylean®; Elanco, Indianapolis, IN) was approved for use in finishing diets for swine. Ractopamine has been shown to increase ADG, improve F/G, and increase pig weights. Feed intake tends to decrease when ractopamine is added to the diet, which makes it necessary to increase amino acids in the diet to ensure that the pig's requirements are met. With pigs growing faster, increasing lean accretion, and decreasing fat deposition, our objective was to determine the lysine requirement for pigs fed ractopamine the last 21 d before market.

Procedures

This experiment was conducted in a commercial finishing facility in southwestern Minnesota. There were 42 pens, with seven pens per treatment and 21 or 22 pigs per pen. A total of 919 gilts were allotted by weight,

¹Paylean is a registered trademark of Elanco Animal Health, Indianapolis, IN.

²Food Animal Health & Management Center, College of Veterinary Medicine.

³Anjinomoto Heartland Lysine LLC, Chicago, IL.

with an initial weight of 219 lb. Pens were totally slatted, with a self feeder and waterer. The facility was double curtain sided, with natural ventilation and a deep pit for manure storage. The six dietary treatments were based on the amount of true ileal digestible (TID) lysine in the diet (0.75, 0.85, 0.95, 1.05, 1.15, and 1.25%), and each treatment also contained 6.5 g/ton of Paylean (Table 1). These TID lysine levels correspond to 0.86, 0.97, 1.08, 1.19, 1.29, and 1.40% total lysine. All diets were based on corn-soybean meal, with 2% added fat and 1.5 lb/ton of crystalline lysine.

Pen weights and feed intake were determined on d 0, 7, 14, and 21 to calculate ADG, ADFI, and F/G. At the conclusion of the trial, pigs in each pen were identified with a unique tattoo to obtain carcass information. At the slaughtering facility (Swift Inc, Worthington, MN), carcass parameters were measured. Data were analyzed by using the PROC mixed procedure in SAS v. 8.1 as a randomized design.

Results and Discussion

From d 0 to 14, pigs fed 1.05% TID lysine had the greatest ADG and second-lowest F/G (2.59 and 2.25, respectively). Increasing TID

lysine improved ADG (linear, $P < 0.05$) and F/G (linear, $P < 0.01$; quadratic $P < 0.05$) from d 0 to 14. As the trial progressed (overall d 0 to 21), however, ADG and F/G had the greatest improvement when pigs were fed the 1.15% TID lysine treatment. Average daily gain increased (linear, $P < 0.005$) and F/G improved (linear, $P < 0.0004$; quadratic $P < 0.07$) for the overall data, whereas there were no differences in ADFI ($P > 0.75$). Although the ADG and F/G responses were linear, there was no improvement in growth observed for treatments above 1.15% TID lysine.

Percentage lean increased (linear, $P < 0.03$) and FFLI tended to increase (linear, $P < 0.07$ and quadratic, $P < 0.09$). There was a linear trend ($P < 0.08$) for decreased backfat thickness with increasing TID lysine. The lean premium also improved (linear, $P < 0.05$), but other carcass criteria were not affected by TID lysine level. In conclusion, pigs fed ractopamine in this commercial research barn required 1.05% to 1.15% TID lysine (1.19% to 1.29% total lysine) to optimize growth performance. These lysine concentrations are higher than previously observed in pigs fed ractopamine.

Table 1. Diet Composition (As-fed Basis)^a

| Item | TID ^b Lysine Content % | | | | | |
|---------------------------|-----------------------------------|-------|-------|-------|-------|-------|
| | 0.75 | 0.85 | 0.95 | 1.05 | 1.15 | 1.25 |
| Corn | 78.84 | 71.84 | 67.85 | 63.85 | 59.85 | 55.86 |
| Soybean meal (46.5% CP) | 19.85 | 23.83 | 31.20 | 31.79 | 35.77 | 39.75 |
| Choice white grease | 2.00 | 2.00 | 2.00 | 2.00 | 2.00 | 2.00 |
| Monocalcium P (21% P) | 0.60 | 0.58 | 0.56 | 0.54 | 0.52 | 0.50 |
| Limestone | 1.10 | 1.10 | 1.10 | 1.10 | 1.10 | 1.10 |
| Salt | 0.40 | 0.40 | 0.40 | 0.40 | 0.40 | 0.40 |
| L-Lysine HCl | 0.075 | 0.075 | 0.075 | 0.075 | 0.075 | 0.075 |
| DL-methionine | 0.00 | 0.015 | 0.030 | 0.045 | 0.060 | 0.075 |
| L-threonine | 0.00 | 0.022 | 0.043 | 0.065 | 0.086 | 0.108 |
| Vitamin premix | 0.050 | 0.050 | 0.050 | 0.050 | 0.050 | 0.050 |
| Trace mineral premix | 0.050 | 0.050 | 0.050 | 0.050 | 0.050 | 0.050 |
| Ractopamine HCl | 0.036 | 0.036 | 0.036 | 0.036 | 0.036 | 0.036 |
| Total | 100 | 100 | 100 | 100 | 100 | 100 |
| ME (Mcal/lb) | 1.55 | 1.55 | 1.55 | 1.55 | 1.55 | 1.55 |
| Total lysine, % | 0.86 | 0.97 | 1.08 | 1.19 | 1.29 | 1.40 |
| TID methionine, % | 0.24 | 0.27 | 0.30 | 0.34 | 0.37 | 0.40 |
| TID methionine+cystine, % | 0.49 | 0.54 | 0.59 | 0.65 | 0.70 | 0.75 |
| TID threonine, % | 0.50 | 0.58 | 0.65 | 0.73 | 0.80 | 0.88 |
| TID tryptophan, % | 0.15 | 0.18 | 0.20 | 0.22 | 0.24 | 0.26 |
| TID isoleucine, % | 0.57 | 0.63 | 0.70 | 0.77 | 0.83 | 0.90 |
| TID valine, % | 0.65 | 0.72 | 0.79 | 0.85 | 0.92 | 0.98 |

^aAll diets were based on corn and soybean meal.

^bTrue ileal digestible.

Table 2. Effects of Dietary Lysine Level on Performance of Pigs Fed Ractopamine^a

| Item | TID ^b Lysine, % | | | | | | SE | Probability, <i>P</i> < | | |
|-----------------------------------|----------------------------|--------|--------|--------|--------|--------|-------|-------------------------|--------|-----------|
| | 0.75 | 0.85 | 0.95 | 1.05 | 1.15 | 1.25 | | Trt | Linear | Quadratic |
| Initial weight, lb | 219.0 | 219.0 | 219.1 | 219.1 | 219.0 | 219.0 | 3.09 | 1.00 | 0.99 | 0.98 |
| d 0 to 14 | | | | | | | | | | |
| ADG, lb | 2.36 | 2.36 | 2.49 | 2.59 | 2.55 | 2.50 | 0.08 | 0.25 | 0.05 | 0.22 |
| ADFI, lb | 5.85 | 5.71 | 5.78 | 5.83 | 5.69 | 5.78 | 0.11 | 0.89 | 0.69 | 0.72 |
| F/G | 2.49 | 2.44 | 2.32 | 2.25 | 2.23 | 2.34 | 0.07 | 0.06 | 0.01 | 0.05 |
| d 0 to 21 | | | | | | | | | | |
| ADG, lb | 2.12 | 2.16 | 2.20 | 2.28 | 2.34 | 2.27 | 0.06 | 0.07 | 0.005 | 0.34 |
| ADFI, lb | 5.83 | 5.71 | 5.75 | 5.80 | 5.82 | 5.77 | 0.09 | 0.93 | 0.88 | 0.75 |
| F/G | 2.76 | 2.64 | 2.62 | 2.54 | 2.49 | 2.56 | 0.05 | 0.005 | 0.0004 | 0.07 |
| Final weight, lb | 263.4 | 264.5 | 265.4 | 267.4 | 268.2 | 265.8 | 3.22 | 0.91 | 0.37 | 0.55 |
| Carcass weight, lb | 198.2 | 198.3 | 199.2 | 199.9 | 200.5 | 197.6 | 2.50 | 0.96 | 0.83 | 0.47 |
| Yield % | 76.13 | 75.75 | 76.25 | 75.66 | 75.66 | 75.77 | 0.004 | 0.78 | 0.39 | 0.88 |
| Backfat, in | 0.62 | 0.61 | 0.59 | 0.57 | 0.58 | 0.59 | 0.02 | 0.33 | 0.08 | 0.20 |
| Loin depth, in | 2.25 | 2.31 | 2.28 | 2.27 | 2.32 | 2.37 | 0.06 | 0.77 | 0.21 | 0.63 |
| Lean % | 56.16 | 56.39 | 56.88 | 56.87 | 57.00 | 56.91 | 0.30 | 0.28 | 0.03 | 0.27 |
| FFLI | 50.98 | 51.06 | 51.32 | 51.51 | 51.53 | 51.22 | 0.18 | 0.18 | 0.07 | 0.09 |
| Carcass criteria, \$ ^c | | | | | | | | | | |
| Sort loss/cwt | -1.86 | -2.04 | -1.94 | -2.02 | -2.05 | -1.74 | 0.31 | 0.98 | 0.85 | 0.50 |
| Value/cwt carcass ^d | 72.96 | 73.11 | 73.38 | 73.26 | 73.24 | 73.57 | 0.38 | 0.91 | 0.31 | 0.94 |
| Income/pig | 144.05 | 144.49 | 145.41 | 146.43 | 146.51 | 146.17 | 1.77 | 0.88 | 0.24 | 0.66 |
| Lean premium | 4.25 | 4.57 | 4.74 | 4.70 | 4.71 | 4.73 | 0.17 | 0.29 | 0.05 | 0.17 |
| Value live | 54.71 | 54.42 | 55.15 | 55.26 | 54.80 | 55.72 | 0.52 | 0.58 | 0.16 | 0.79 |

^aA total of 919 gilts (PIC L337 × C22; initially 219 lb) were used in a 21-d feeding trial. Pigs were fed ractopamine (6.5 g/ton), with 6 dietary treatments formulated to titrate the lysine requirement. A total of 42 pens were used, with 7 pens per treatment and 21 or 22 pigs per pen.

^bTrue ileal digestible.

^cCarcass base price was \$70.58/cwt.

^dValue/cwt for carcass was calculated as: carcass base price – sort + lean premium.