

# Kansas Agricultural Experiment Station Research Reports

---

Volume 0  
Issue 10 *Swine Day (1968-2014)*

Article 1156

---

2007

## Digestible energy content of corn and triticale when fed to finishing pigs

C Feoli

C R. Monge

Terry L. Gugle

*See next page for additional authors*

Follow this and additional works at: <https://newprairiepress.org/kaesrr>



Part of the [Other Animal Sciences Commons](#)

---

### Recommended Citation

Feoli, C; Monge, C R.; Gugle, Terry L.; and Hancock, Joe D. (2007) "Digestible energy content of corn and triticale when fed to finishing pigs," *Kansas Agricultural Experiment Station Research Reports*: Vol. 0: Iss. 10. <https://doi.org/10.4148/2378-5977.6996>

This report is brought to you for free and open access by New Prairie Press. It has been accepted for inclusion in Kansas Agricultural Experiment Station Research Reports by an authorized administrator of New Prairie Press. Copyright 2007 the Author(s). Contents of this publication may be freely reproduced for educational purposes. All other rights reserved. Brand names appearing in this publication are for product identification purposes only. No endorsement is intended, nor is criticism implied of similar products not mentioned. K-State Research and Extension is an equal opportunity provider and employer.



---

## Digestible energy content of corn and triticale when fed to finishing pigs

### Authors

C Feoli, C R. Monge, Terry L. Gugle, and Joe D. Hancock

## DIGESTIBLE ENERGY CONTENT OF CORN AND TRITICALE WHEN FED TO FINISHING PIGS<sup>1</sup>

*C. Feoli, J. D. Hancock, C. R. Monge, and T. L. Gugle*

### Summary

A total of 96 pigs (average initial wt of 201 lb) was used to determine the DE content of corn and triticale. The pigs were sorted by sex and ancestry, blocked by weight, and assigned with 12 pigs/pen and four pens/treatment. The diets were corn (97.5% of the formulation) and triticale (97.8% of the formulation) with added vitamins, minerals, and amino acids. Feed (meal form) and water were consumed on an *ad libitum* basis. The pigs were allowed to adjust to the experimental diets for 4 d. On the afternoon of d 4 and morning of d 5, feces were collected from no less than six pigs/pen (via rectal massage). The feed and fecal samples were dried, ground, and analyzed for concentrations of DM, N, and GE with chromic oxide used as an indigestible marker. Digestibility of DM was greater ( $P<0.03$ ) for pigs fed corn vs triticale (82.8 vs 81.2%, respectively). However, the opposite was true for digestibility of N ( $P<0.002$ ) with values of 67.8% for corn and 74.7% for triticale. Digestibility of GE was not different ( $P>0.26$ ) among the cereal grains. However, when the gross energy for the cereals was multiplied by their respective digestibility coefficients, triticale grain had greater ( $P<0.02$ ) DE with a value of 1,531 kcal/lb vs 1,479 kcal/lb for corn. The DE of the corn used in this experiment was low compared to NRC values, but nonetheless our

results indicated that this particular triticale was utilized well by finishing pigs supporting greater digestibility of N and having greater DE than corn.

(Key words: feed ingredients, triticale.)

### Introduction

Triticale is a hybrid resulted from a cross between wheat and rye. This “small grain” crop was developed as a forage source for cattle and for its improved protein quality (e.g., greater lysine content) for nonruminants. Development of new varieties has renewed interest in this cereal as a feedstuff in Kansas but also has led to questions about its feeding value for pigs. Therefore, the objective of the experiment reported herein was to determine the nutrient digestibility of corn and triticale when fed to finishing pigs.

### Procedures

A total of 96 pigs (average initial weight of 201 lb) was used. The pigs were sorted by sex and ancestry, blocked by weight, and assigned to pens. There were 12 pigs/pen and four pens/treatment. Diets (Table 1) were yellow corn (97.5% of the formulation) and triticale (ThunderCale as 97.8% of the formulation) with added vitamins, minerals, and amino acids. The diets had 0.52% lysine,

---

<sup>1</sup>Appreciation is given to Ehmke Seed for supplying the triticale (ThunderCale) used in this experiment.

0.45% Ca, and 0.40% total P with 0.25% chromic oxide added as an indigestible marker.

**Table 1. Composition of Diets, %<sup>a</sup>**

Ingredient	Corn	Triticale
Cereal grain	97.53	97.83
Limestone	0.89	0.95
Monocalcium phosphate (21% P)	0.60	0.37
Salt	0.20	0.18
L-lysine HCl	0.34	0.18
L-threonine	0.03	-
L-isoleucine	0.02	-
Vitamin premix	0.03	0.05
Sow add pack	0.02	0.06
Trace mineral premix	0.09	0.13
Chromic oxide <sup>b</sup>	0.25	0.25
Total	100.00	100.00

<sup>a</sup>Diets were formulated to 0.52% lysine, 0.45% Ca, and 0.40% total P.

<sup>b</sup>Used as an indigestible marker.

Feed (meal form) and water were consumed on an *ad libitum* basis with pigs and feeders weighed on d 0 and 5. The pigs were allowed to adjust to the experimental diets for 4 d. Feces were collected from no less than six pigs/pen (via rectal massage) on the after-

noon of d 4 and morning of d 5. The feed and fecal samples were dried, ground, and analyzed for concentrations of DM, N, GE, and Cr to allow calculation of apparent digestibility of nutrients using the indirect ratio method. All data were analyzed as a randomized complete block design using the MIXED procedure of SAS with a significant F-test indicating treatment differences.

## Results and Discussion

The pigs gained weight (an average of 1.75 lb/d) and ate the diets well (greater than 7 lb/d) during this brief (5-d) experiment. Comparison of nutrient utilization among the cereals showed digestibility of DM was greater ( $P<0.03$ ) for pigs fed corn vs triticale (82.8 vs 81.2%, respectively). However, the opposite was true for digestibility of N ( $P<0.002$ ) with values of 67.8% for corn and 74.7% for triticale. Digestibility of GE was not different ( $P>0.26$ ) among the cereal grains. However, when the gross energy for the cereals was multiplied by their respective digestibility coefficients, triticale grain had greater ( $P<0.02$ ) DE with a value of 1,531 kcal/lb vs 1,479 kcal/lb for corn. The DE of the corn used in this experiment was low compared to NRC values, but nonetheless our results indicated that this particular triticale was utilized well by finishing pigs supporting greater digestibility of N and having greater DE than corn.

**Table 2. Digestibility of Nutrients in Corn and Triticale for Finishing Pigs<sup>a</sup>**

Item	Corn	Triticale	SE	P value
Digestibility, %				
Dry matter	82.8	81.2	0.8	0.03
Nitrogen (protein)	67.8	74.7	1.6	0.002
Gross energy	81.1	80.5	0.9	- <sup>b</sup>
DE content of the grain, kcal/lb	1,479	1,531	16	0.02

<sup>a</sup>A total of 96 pigs (12 pigs/pen and four pens/treatment) with an average initial weight of 201 lb.

<sup>b</sup>Dash indicates  $P>0.15$ .