

1999

Protect Streambanks with Trees

Wayne A. Geyer

T. Neopl

K. Brooks

Follow this and additional works at: <https://newprairiepress.org/kaesrr>

Recommended Citation

Geyer, Wayne A.; Neopl, T.; and Brooks, K. (1999) "Protect Streambanks with Trees," *Kansas Agricultural Experiment Station Research Reports*: Vol. 0: Iss. 12. <https://doi.org/10.4148/2378-5977.7340>

This report is brought to you for free and open access by New Prairie Press. It has been accepted for inclusion in Kansas Agricultural Experiment Station Research Reports by an authorized administrator of New Prairie Press. Copyright 1999 Kansas State University Agricultural Experiment Station and Cooperative Extension Service. Contents of this publication may be freely reproduced for educational purposes. All other rights reserved. Brand names appearing in this publication are for product identification purposes only. No endorsement is intended, nor is criticism implied of similar products not mentioned. K-State Research and Extension is an equal opportunity provider and employer.



Protect Streambanks with Trees

Keywords

Keeping up with research; SRL 124 (Nov. 1999); Kansas Agricultural Experiment Station contribution; no. 00-123-S; Trees; Streambank erosion; Natural woodlands; Streambanks

Creative Commons License

This work is licensed under a [Creative Commons Attribution 4.0 License](https://creativecommons.org/licenses/by/4.0/).



PROTECT STREAMBANKS WITH TREES

W.A. Geyer, T. Neppl, and K. Brooks*

Acknowledgement

Support for the evaluation of trees and streambank erosion from the USDA Forest Service National Agroforestry Center is appreciated.

*Professor, Department of Horticulture, Forestry and Recreation Resources and former graduate student and Associate Professor, Department of Landscape Architecture/Regional and Community Planning, Kansas State University.

Contribution No. 99-112-S from the Kansas Agricultural Experiment Station.

Contents of this publication may be freely reproduced for educational purposes. All other rights reserved. In each case, give credit to the author(s), name of work, Kansas State University, and the date of the work was published.

Kansas State University Agricultural Experiment Station and Cooperative Extension Service, Manhattan, Kansas 66506

SRL 122

October 1998

It is the policy of Kansas State University Agricultural Experiment Station and Cooperative Extension Service that all persons shall have equal opportunity and access to its educational programs, services, activities, and materials without regard to race, color, religion, national origin, sex, age, or disability. Kansas State University is an Affirmative Action employer. These materials may be available in alternative formats. 1.8M

Kansas State University
Agricultural Experiment Station and
Cooperative Extension Service




Figure 1. Historical mean discharge of the Kansas River at the beginning point of the study area.

Procedures

Study Area. The study area was in the Kansas River Basin, which covers about 60,000 sq miles in three states. The specific zone of study was a 40-mile portion near Manhattan, Kansas (Figure 2). The woody vegetation along the river consists of many tree species, including cottonwood (*Populus deltoides*), silver maple (*Acer saccharinum*), willow species (*Salix* sp.), hackberry (*Celtis occidentalis*), sycamore (*Platanus occidentalis*), and elm species (*Ulmus* sp.)




Figure 2. Diagram of the river study area.

Tools of Investigation. Aerial photographic images were chosen to compare pre- and postflood streambanks (slides were taken in December 1992 and December 1993), using easily identified features such as riparian vegetation and streambank edge. Local offices of the Consolidated Farm Service Agency provided slide images of the study area.

Data Collection Process. Photocopies of the slides were transferred with a CalComp digitizing tablet into a LandCADDI R12 computer-aided drawing (CAD) program file. The photocopies were calibrated to previously digitized, USGS, 7.5 minute, topographical, quadrangle maps to enable accurate scaling and analysis of the images. Only features useful in calibrating slide images, such as roads, railroad tracks, and mile section lines, were digitized from the quadrangle maps.

Once the base maps were digitized, streambank edges from the 1992 and 1993 slide images of the Kansas River were digitized. Notes as to land-cover condition were made within the drawing as needed. Then a river centerline was interpolated within the CAD drawing using the 1992 streambanks as a guide. The centerline served as a reference for river position and demarcation and land-cover classification.

Data collection points were established at regular 500-foot intervals along the centerline in the study area. At these points, land cover was classified using the 1992 aerial photographs as references. Erosion and deposition amounts were estimated by measuring the perpendicular distance from the 1992 streambank to the 1993 streambank at data collection points.

Land-cover vegetative types in the 100-ft zone next to the 1992 streambank were categorized as: forest land, cropland, grassland, or single tree-row. A forested streambank is one on which the dominant land-cover type (>51%) is woody vegetation. A cropland or grassland streambank is one on which the dominant land-cover type is either agricultural crops or grass. Finally, a single tree-row streambank is one on which a single row of trees exists adjacent to a nonforest land-cover type.

Vegetation data were collected for both the left and right side streambanks at every data collection point and inserted into the CAD program. Data from each side of the river were pooled.

Results and Discussion

Data were collected from a total of 204 (103 right side and 101 left side) streambank points. These data points were classified as follows: 96 points as forest land, 37 as cropland, 47 as grassland, and 24 as single-tree row. Thus, 59% of the banks were tree lined.

About 62% of the banks had sandy soils, and 38% had loamy soils. The streams generally were of two types, meandering (63%) and relatively shallow and braided (37%). Many of the streambanks were entrenched.

Land-cover vegetation significantly (1% level) affected the amount of erosion of the lateral streambank (Figure 3). Both forest and the single tree-vegetation types collected soil, with mean depositions of 10 ft and 4 ft, respectively, which were not significantly different, whereas grassland lost an average of 78 ft, and cropland lost an average of 150 ft. The surface acreage of the lost land was considered for each mile of stream erosion: 9.4 acres for grassland and 18.2 acres for cultivated land. The latter is equivalent to about 1/4 section of land for every 10 miles of stream distance in this study.

These results showed that woody vegetation is highly effective in protecting streambanks. Water movement is slowed by the standing trees, thus reducing the energy available for erosion and allowing deposition of suspended materials. Greater rooting depth, larger and stronger roots, and perhaps greater rooting density also stabilize the soil mantle.




Figure 3. Lateral movement of streambank by land-cover type.

Conclusions

Trees have a highly beneficial effect in protecting streambanks against erosion during large floods. Tree cover reduced the extent of streambank erosion caused by the extreme flood of 1993 on the Kansas River. Forested areas on both sides of the river sustained less erosion than those reaches that had no woody vegetation cover. Natural stands of timber are best left standing to protect the riparian ecosystem, affording greater bank stability and water quality and reducing downstream sedimentation (Figure 4). Cultivating land next to the stream bank should be avoided because of the large acreage loss during high water events, which occur frequently in central Kansas.



Figure 4. Natural woody vegetation should be left to protect streambanks.