

1992

Symptom Response of Sorghum Hybrids Infected by Sugarcane Mosaic Virus Strain MDMV-B (1992)

Dallas Lynn Seifers

Kenneth D. Kofoid


Follow this and additional works at: <https://newprairiepress.org/kaesrr>

Recommended Citation

Seifers, Dallas Lynn and Kofoid, Kenneth D. (1992) "Symptom Response of Sorghum Hybrids Infected by Sugarcane Mosaic Virus Strain MDMV-B (1992)," *Kansas Agricultural Experiment Station Research Reports*: Vol. 0: Iss. 12. <https://doi.org/10.4148/2378-5977.7366>

This report is brought to you for free and open access by New Prairie Press. It has been accepted for inclusion in Kansas Agricultural Experiment Station Research Reports by an authorized administrator of New Prairie Press. Copyright 1992 the Author(s). Contents of this publication may be freely reproduced for educational purposes. All other rights reserved. Brand names appearing in this publication are for product identification purposes only. No endorsement is intended, nor is criticism implied of similar products not mentioned. K-State Research and Extension is an equal opportunity provider and employer.





Keeping
Up With
Research

104

December 1992

**SYMPTOM RESPONSE OF
SORGHUM HYBRIDS INFECTED
BY SUGARCANE MOSAIC VIRUS
STRAIN MDMV-B**

**D. L. Seifers and K. D. Kofoid
Fort Hays Branch Experiment Station**

Sorghum [*Sorghum bicolor* (L.) Moench] infected by sugarcane mosaic virus strain MDMV-B (SCMV-MDMV-B) (formerly maize dwarf mosaic virus strain B) expresses variable symptoms depending on plant genotype and temperature. Infection causes a yellow to dark green pattern (mosaic), and a necrotic reaction (red-leaf) develops on certain genotypes when the temperature drops below 21°C (69°F). Hybrids expressing the red-leaf symptom show greater reduction in yield. Immunity to SCMV-MDMV-B is not present in commercially available hybrid sorghums.

Procedure

Studies were conducted at the Fort Hays Branch Experiment Station to determine symptom response of sorghum hybrids infected by SCMV-MDMV-B both under field conditions (2 years of testing) and in growth chambers. Sorghum seed was planted in 6.1 m rows. When at the two- to three-leaf stage, plants in half of each row were inoculated, and plants in the remaining half of the

AGRICULTURAL EXPERIMENT STATION

Kansas State University, Manhattan
Marc A. Johnson, Director

row served as healthy controls. Plants were inoculated with a Devilbiss EGA-502 spray gun attached to a 3.5 L (1 gal) inoculum reservoir. Compressor line pressure was maintained at 7.0 kg/cm² (100 lbs/in²). Symptom notes were recorded at biweekly intervals during the growing season beginning 2 weeks after inoculation. Planting was delayed until late June to assure cool temperatures for induction of red-leaf necrosis.

In growth chamber tests, sorghum hybrids were planted in metal flats with 15 seeds/entry and 33 entries/flat. Seedlings were inoculated at the two-leaf stage (on second leaf) with a Devilbiss atomizer at the same pressure used for field inoculations. Inoculated plants were maintained in growth chambers at 27 ± 2°C (80 ± 2°F) with 14 hr of fluorescent light. Control plants, inoculated with buffer and abrasive only, were maintained in the same environment. The presence of mosaic symptoms was recorded 14 days following inoculation. The temperature of the growth chambers then was decreased to 15°C (59°F). Four days later, plants were rated for the presence of red-leaf necrosis. Growth chamber tests were conducted once.

Results

Reactions of hybrids are shown in Table 1. SS = inoculated leaf and two to three leaves above the inoculated leaf have necrotic streaks and spots that often coalesce and cover 1–25 percent of a leaf. New leaves above these show few or no symptoms, so that at maturity, no symptoms of any type occur in the upper leaves. RL1 = seedling and juvenile plants have a necrotic red-leaf reaction consisting of various widths of streaks that often coalesce and cover 25–50 percent of a leaf. At maturity, the flag leaf and the subtending 3–4 leaves have a few necrotic streaks covering 1–10 percent of a leaf. RL2 = seedling and juvenile plants have a severe necrotic red-leaf reaction, not in streaks or spots, but continuous necrosis covering 80–100 percent of a leaf. In plants that survive to maturity, any new leaf will develop a like necrosis with onset of cool temperatures. M = plants having only a mosaic reaction. Current data indicate that plants with the M reaction suffer less yield loss than plants developing necrosis (RL types).

**Table 1. Reactions of Sorghum Hybrids to Infection by
Sugarcane Mosaic Virus Strain MDMV-B**

| Brand | Hybrid | Symptom Response ^a |
|------------|----------|-------------------------------|
| Agrigene | AG2470W | SS |
| Agrigene | AG2725 | RL2 |
| Agripro | AP 9250 | SS |
| Agripro | AP 9830 | SS |
| Agripro | AP 9850 | M |
| Agripro | ST 686 | M |
| Agripro | ST D701G | RL2 |
| Asgrow | A504 | SS |
| Asgrow | Madera | M |
| Asgrow | Osage | M |
| Asgrow | Seneca | M |
| Asgrow | Topaz | RL2 |
| Cargill | 1099 | RL2 |
| Cargill | 575 | SS |
| Cargill | 607E | SS |
| Cargill | 618Y | M |
| Cargill | 630 | SS |
| Cargill | 837 | SS |
| Cargill | 847 | M |
| Casterline | SR 305 | M |
| Casterline | SR 314EC | RL2 |
| Casterline | SR 315E | RL2 |
| Casterline | SR 319E | RL2 |
| Casterline | SR 324E | RL1 |
| Dekalb | DK-39Y | RL2 |
| Dekalb | DK-40Y | SS |
| Dekalb | DK-41Y | SS |
| Dekalb | DK-48 | SS |
| Dekalb | DK-56 | M |
| Dekalb | DK-66 | RL2 |
| Dekalb | M-572 | RL2 |
| Delange | DSA 117 | M |
| Delange | DSA 121 | SS |
| Delange | DSA 131 | RL2 |
| Delange | DSA 141C | SS |
| Delange | DSA 922 | RL2 |
| Deltapine | 1506 | RL2 |
| Deltapine | G-1492 | SS |
| Deltapine | G-1616 | SS |
| Deltapine | G-550E | RL1 |
| DyNA | DN 31 | RL1 |
| Fontanelle | 5550 | RL2 |
| Fontanelle | 5590 | RL2 |
| Fontanelle | W 5000 | SS |
| Garrison | SG-651 | SS |
| Garrison | SG-821 | SS |
| Garrison | SG-942 | RL2 |
| Garst | 5319 | RL2 |
| Garst | 5392 | M |
| Garst | 5503 | M |
| Garst | 5511 | RL2 |

| Brand | Hybrid | Symptom Response ^a |
|----------------|--------------|-------------------------------|
| Garst | 5522Y | SS |
| Garst | 5681 | SS |
| Golden Acres | T-E 76 | M |
| Golden Acres | T-E 77-E | RL2 |
| Golden Acres | T-E Dinero | RL2 |
| Golden Acres | T-E Eden | M |
| Golden Acres | T-E Rio | RL2 |
| Golden Acres | T-E Wahoo | SS |
| Golden Acres | T-E Y-60 | SS |
| Golden Acres | T-E Y-75 | M |
| Golden Acres | T-E Y-101G | RL1 |
| Golden Harvest | H-388W | SS |
| Golden Harvest | H-444W | SS |
| Golden Harvest | H-514B | RL2 |
| Golden Harvest | H-515E | SS |
| GRI | 890111 | RL2 |
| Groagri | GSC 3146 | M |
| Groagri | GSC 3148 | M |
| Groagri | GSC 3150 | RL2 |
| Groagri | GSC 3605 | RL1 |
| Hoegemeyer | 6636 | RL1 |
| Hoegemeyer | 671 | SS |
| Hoegemeyer | 6744 | SS |
| Hoegemeyer | 6878 | RL2 |
| Hoegemeyer | 688 | RL2 |
| Horizon | 200Y | SS |
| Horizon | 213Y | SS |
| Horizon | 216G | M |
| Horizon | 76G | M |
| Hyperformer | 1255 DR | RL2 |
| Hyperformer | Honcho | RL2 |
| Hyperformer | HSC 1289C | M |
| Hyperformer | HSC Cherokee | M |
| Jacques | 211 | M |
| Jacques | 377-W | SS |
| Jacques | 411 | M |
| Jacques | 606E | RL1 |
| NC+ | 270 | RL1 |
| NC+ | 371 | SS |
| NC+ | 472 | SS |
| NC+ | 7B81 | SS |
| NC+ | 7C49 | SS |
| NC+ | Y363 | SS |
| Northrup-King | 2656 | RL2 |
| Northrup-King | KS-383Y | SS |
| Northrup-King | KS-555Y | SS |
| Northrup-King | KS-710 | RL2 |
| Northrup-King | KS-714Y | SS |
| Northrup-King | KS-737 | SS |
| Northrup-King | S-9740Y | SS |
| Northrup-King | S 9750 | RL2 |

| Brand | Hybrid | Symptom Response ^a |
|---------|----------|-------------------------------|
| Ohlde | 120C | SS |
| Ohlde | 134 | SS |
| Ohlde | 139A | RL2 |
| Ohlde | 140W | SS |
| Ohlde | 246y | RL1 |
| Oro | Alpha | M |
| Oro | Amigo | RL2 |
| Oro | Baron | M |
| Oro | Edge | M |
| Oro | G XTRA | RL2 |
| Oro | Hombre | M |
| Oro | Ivory | SS |
| Oro | Ultra | RL2 |
| Pioneer | 8231Y | RL1 |
| Pioneer | 8260 | SS |
| Pioneer | 8358 | RL1 |
| Pioneer | 8379 | SS |
| Pioneer | 8500 | SS |
| Pioneer | 8536Y | SS |
| Pioneer | 8557Y | M |
| Pioneer | 8601 | RL1 |
| Pioneer | 8696Y | M |
| Pioneer | 8699 | M |
| Pioneer | 8771 | M |
| SSI | SBP001 | RL1 |
| SSI | SBP005 | RL1 |
| SSI | SBP011 | SS |
| Stine | SM 68 BR | RL2 |
| Stine | SM 69 | M |
| Stine | SM 74 | RL2 |
| Stine | SM 75R | RL2 |
| Terra | TR 4580 | RL1 |
| Terra | TR 4620 | RL2 |
| Terra | TR 4637Y | SS |
| Triumph | TR 46 | M |
| Triumph | TR 50yG | M |
| Triumph | TR 58Y | SS |
| Triumph | TR 60-G | M |
| Triumph | TR 65-G+ | RL1 |
| Triumph | TR 74CR | SS |
| Triumph | TWO 64yG | RL2 |
| Triumph | TWO 80-D | RL2 |
| Warner | W-624-Y | RL2 |
| Warner | W-625-Y | RL2 |
| Warner | W-902-W | M |
| Warner | W-917-E | M |
| Wilson | 515W | SS |
| Wilson | 522W | SS |
| Wilson | 535Y | RL2 |
| Wilson | 622E | M |

^aSymptom responses are the same for both field and growth chamber tests.

Acknowledgements

This research was partially funded by a grant from the Kansas Sorghum Commission.

Contribution no. 93-145-S from the KAES.



Agricultural Experiment Station
Kansas State University
Manhattan 66506-4008

Keeping Up With Research 104

December 1992

Kansas State University is committed to a policy of nondiscrimination on the basis of race, sex, national origin, handicap, religion, age, sexual orientation, or other nonmerit reasons, in admissions, educational programs or activities, and employment, all as required by applicable laws and regulations. Responsibility for coordination of compliance efforts and receipt of inquiries, including those concerning Title IX of the Education Amendments of 1972 and Section 504 of the Rehabilitation Act of 1973, has been delegated to Jane D. Rowlett, Ph.D., Director, Affirmative Action Office, 214 Anderson Hall, Kansas State University, Manhattan, Kansas 66506-0104, (913/532-6220).

1M