

Kansas Agricultural Experiment Station Research Reports

Volume 3

Issue 5 *Southwest Research-Extension Center Reports*

Article 14

2017

Alternative Cropping Systems with Limited Irrigation

A. Schlegel

Kansas State University, schlegel@ksu.edu

Follow this and additional works at: <https://newprairiepress.org/kaesrr>

 Part of the [Agronomy and Crop Sciences Commons](#)

Recommended Citation

Schlegel, A. (2017) "Alternative Cropping Systems with Limited Irrigation," *Kansas Agricultural Experiment Station Research Reports*: Vol. 3: Iss. 5. <https://doi.org/10.4148/2378-5977.7396>

This report is brought to you for free and open access by New Prairie Press. It has been accepted for inclusion in Kansas Agricultural Experiment Station Research Reports by an authorized administrator of New Prairie Press. Copyright 2017 Kansas State University Agricultural Experiment Station and Cooperative Extension Service. Contents of this publication may be freely reproduced for educational purposes. All other rights reserved. Brand names appearing in this publication are for product identification purposes only. No endorsement is intended, nor is criticism implied of similar products not mentioned. K-State Research and Extension is an equal opportunity provider and employer.



Alternative Cropping Systems with Limited Irrigation

Abstract

A limited irrigation study involving four cropping systems and evaluating four crop rotations was initiated at the Southwest Research-Extension Center near Tribune, KS, in 2012. The cropping systems were two annual systems (continuous corn [C-C] and continuous grain sorghum [GS-GS]) and two 2-year systems (corn-grain sorghum [CGS]) and corn-winter wheat [C-W]). In 2016, corn yields were similar in all rotations, as were grain sorghum yields. This tended to agree with the 4-yr average yields, except for average grain sorghum yields being higher following corn than grain sorghum.

Creative Commons License



This work is licensed under a [Creative Commons Attribution 4.0 License](https://creativecommons.org/licenses/by/4.0/).

Cover Page Footnote

The project was funded in part by Western Kansas Groundwater Management District No. 1.

Alternative Cropping Systems with Limited Irrigation

A. Schlegel

Summary

A limited irrigation study involving four cropping systems and evaluating four crop rotations was initiated at the Southwest Research-Extension Center near Tribune, KS, in 2012. The cropping systems were two annual systems (continuous corn [C-C] and continuous grain sorghum [GS-GS]) and two 2-year systems (corn-grain sorghum [C-GS]) and corn-winter wheat [C-W]). In 2016, corn yields were similar in all rotations, as were grain sorghum yields. This tended to agree with the 4-yr average yields, except for average grain sorghum yields being higher following corn than grain sorghum.

Experimental Procedures

A crop rotation study under sprinkler irrigation at the Kansas State University Southwest Research-Extension Center near Tribune was initiated in the spring of 2012. The study evaluates four different crop rotations with a limited irrigation allocation. The rotations include 1- and 2-year rotations. The crop rotations are 1) continuous corn; 2) corn-winter wheat; 3) corn-grain sorghum; and 4) continuous grain sorghum (a total of 6 treatments). All rotations are limited to 10 inches of irrigation water annually. All crops are grown no-till, while other cultural practices (hybrid selection, fertility practices, weed control, etc.) are selected to optimize production. All phases of each rotation are present each year and replicated four times. Irrigations are scheduled to supply water at the most critical stress periods for the specific crops and limited to 1.5 inches/week. Soil water is measured at planting, during the growing season, and at harvest in 1-ft increments to a depth of 8 ft. Grain yields are determined by machine harvest. Nitrogen fertilizer (UAN) was surface applied (stream) in March to all crops (240 lb N/a for corn, 160 lb N/a for grain sorghum, and 120 lb N/a for wheat). Corn was planted on April 28, 2016, and harvested on September 15, 2016. Grain sorghum was planted on June 1, 2016, and harvested on October 20, 2016. Wheat was planted on September 29, 2015, and harvested on July 8, 2016.

Results and Discussion

Weather conditions were good for crop production in 2016. Precipitation was above normal for April, July, August, and September. Open pan evaporation was 13% below normal from April through September. Corn yields in 2016 were similar for all rotations with a range of 174 to 186 bu/a (Table 1). Wheat yields in 2016 (82 bu/a) were greater than the multi-year average yield of 64 bu/a (Table 2). Grain sorghum yields were similar following corn or grain sorghum at about 150 bu/a. Averaged across four years, continuous grain sorghum yields were 10 bu/a less than following corn.

Available soil water at corn planting and harvest was similar for all rotations in 2016 (Table 3). Fallow efficiency was less following wheat than following either corn or grain sorghum. For wheat, available soil water at planting and harvest was greater than the 4-yr average (Table 4). The only difference observed with grain sorghum was more fallow accumulation for grain sorghum following corn than following grain sorghum. This was consistent with the average fallow accumulation for the past 4 years. Average crop water use was similar for all rotations for corn and both rotations with grain sorghum.

Acknowledgment

The project was funded in part by Western Kansas Groundwater Management District No. 1.

Table 1. Grain yield of three crops under limited irrigation as affected by rotation in 2016

Rotation	Corn	Wheat	Grain sorghum
	----- bu/a -----		
Continuous corn	174	---	---
Continuous grain sorghum	---	---	149
Corn-wheat	181	82	---
Corn-grain sorghum	186	---	154
Least significant difference _(0.05)	17	---	25

Table 2. Grain yields of three crops under limited irrigation as affected by rotation across years 2013 - 2016

Rotation	Corn	Wheat	Grain sorghum
	----- bu/a -----		
Continuous corn	170b ¹	---	---
Continuous grain sorghum	---	---	137b
Corn-wheat	184a	64	---
Corn-grain sorghum	183a	---	147a
Least significant difference _(0.05)	12	---	9

¹ Means within a column with the same letter are not statistically different at $P = 0.05$.

Table 3. Profile available soil water, crop water use, and fallow accumulation for crop rotations under limited irrigation, Tribune, KS, 2016

Crop	Rotation	Available water			Crop water use	Fallow accumulation	Fallow efficiency
		Previous harvest	Planting	Harvest			
		----- inches -----					%
Corn	C-C	10.03	16.15	14.85	28.72	6.13	63a ¹
	C-W	10.74	15.27	15.94	26.75	4.53	31b
	C-GS	10.28	15.10	15.27	27.24	4.82	72a
LSD _{0.05}		4.14	2.78	3.95	2.57	2.18	21
ANOVA (P > F)							
System		0.915	0.630	0.800	0.229	0.242	0.008
Wheat	C-W	9.58	9.58	12.69	19.59	0	---
ANOVA (P > F)							
System		---	---	---	---	---	---
Grain sorghum	C-GS	7.69b	15.50	11.37	24.52	7.80a	52
	GS-GS	10.57a	15.54	11.32	24.61	4.97b	41
LSD _{0.05}		1.86	1.38	1.46	1.24	2.70	21
ANOVA (P > F)							
System		0.016	0.923	0.934	0.837	0.044	0.213

Note: All crops received ~10 inches of irrigation.

In season rainfall for corn (4/28/16 – 9/15/16) = 17.91 inches; grain sorghum (6/01/16 – 10/20/16) = 12.61 inches; and wheat (9/29/15 – 7/08/16) = 20.29 inches.

C = corn.

W = wheat.

GS = grain sorghum.

LSD = least significant difference.

ANOVA = analysis of variance.

¹ Means within a column with the same letter are not statistically different at $P = 0.05$.

Table 4. Profile available soil water, crop water use, and fallow accumulation for crop rotations under limited irrigation across years, Tribune, KS, 2013-2016

Crop	Rotation	Available water			Crop water use	Fallow accumulation	Fallow efficiency
		Previous harvest	Planting	Harvest			
		----- inches -----					%
Corn	C-C	10.51a ¹	13.68a	12.02a	25.16	3.17ab	36b
	C-W	10.09ab	13.71a	12.13a	25.08	3.62a	24c
	C-GS	9.21b	11.89b	10.20b	25.19	2.68b	53a
LSD _(0.05)		1.19	1.00	1.14	0.99	0.56	8
ANOVA (P > F)							
System		0.091	0.001	0.002	0.972	0.007	0.001
Year		0.001	0.001	0.001	0.001	0.001	0.001
System × year		0.001	0.004	0.014	0.001	0.001	0.001
Wheat	C-W	10.41	10.41	10.76	20.01	0	---
ANOVA (P > F)							
System		---	---	---	---	---	---
Year		0.001	0.001	0.003	0.001	---	---
System × year		---	---	---	---	---	---
Grain sorghum	C-GS	8.08	12.55	10.64	23.31	4.47a	39
	GS-GS	9.08	12.18	10.60	22.98	3.10b	37
LSD _(0.05)		1.14	1.05	1.07	0.68	0.78	11
ANOVA (P>F)							
System		0.082	0.462	0.937	0.314	0.002	0.818
Year		0.001	0.001	0.001	0.001	0.001	0.001
System × year		0.001	0.009	0.787	0.123	0.001	0.392

Note: All crops received ~10 inches of irrigation each year.

C = corn.

W = wheat.

GS = grain sorghum.

LSD = least significant difference.

ANOVA = analysis of variance.

¹ Means within a column with the same letter are not statistically different at $P = 0.05$.