

2017

Effects of Increasing Copper from Tri-basic Copper Chloride or a Copper-Chelate on Growth Performance of Nursery Pigs

C. Carpenter

Kansas State University, Manhattan, cbcarpenter@k-state.edu

J. C. Woodworth

Kansas State University, Manhattan, jwoodworth@k-state.edu

J. M. DeRouchey

Kansas State University, Manhattan, jderouch@k-state.edu

M. D. Tokach

Department of Animal Science and Industry, Kansas State University, mtokach@ksu.edu

See next page for additional authors

Follow this and additional works at: <https://newprairiepress.org/kaesrr>

 Part of the [Other Animal Sciences Commons](#)

Recommended Citation

Carpenter, C.; Woodworth, J. C.; DeRouchey, J. M.; Tokach, M. D.; Goodband, R. D.; Dritz, S. S.; and Usry, J. L. (2017) "Effects of Increasing Copper from Tri-basic Copper Chloride or a Copper-Chelate on Growth Performance of Nursery Pigs," *Kansas Agricultural Experiment Station Research Reports*: Vol. 3: Iss. 7. <https://doi.org/10.4148/2378-5977.7483>

This report is brought to you for free and open access by New Prairie Press. It has been accepted for inclusion in Kansas Agricultural Experiment Station Research Reports by an authorized administrator of New Prairie Press. Copyright 2017 Kansas State University Agricultural Experiment Station and Cooperative Extension Service. Contents of this publication may be freely reproduced for educational purposes. All other rights reserved. Brand names appearing in this publication are for product identification purposes only. No endorsement is intended, nor is criticism implied of similar products not mentioned. K-State Research and Extension is an equal opportunity provider and employer.



Effects of Increasing Copper from Tri-basic Copper Chloride or a Copper-Chelate on Growth Performance of Nursery Pigs

Abstract

A total of 1,452 pigs [Group 1; 350 barrows (DNA 200 × 400; initially 13.2 lb)] and [Group 2; 1,102 barrows and gilts (PIC 1050 × 280; initially 13.3 lb)] were used to determine the effects of added Cu source and level on nursery pig performance. There were 5 pigs per pen and 10 replications per treatment in group 1, and 24 to 27 pigs per pen and 6 replications per treatment in group 2. Pens of pigs were allotted by BW to 1 of 7 dietary treatments arranged as a 2 × 3 factorial plus a control diet, with main effects of Cu source: IntelliBond-C (TBCC; Micronutrients, Indianapolis, IN), or Mintrex-Cu (Cu-chelate; Novus, St. Charles, MO) and Cu level (50, 100, or 150 ppm). Diets were corn-soybean meal-based and were fed in meal form in 2 phases (d 0 to 14 and 14 to 35). All diets contained a trace mineral premix formulated to contribute 17 ppm of Cu from CuSO₄ to the complete diet.

Overall (d 0 to 35), a Cu source × level interaction was observed (linear, $P = 0.042$) for ADG where the rate of improvement with increasing Cu was greater in pigs fed Cuchelate diets compared to those fed TBCC diets. Increasing added Cu increased (linear, $P = 0.001$) ADG, ADFI, and improved F/G. Although Cu source did not influence F/G, pigs fed Cu from Cu-chelate had greater ($P \leq 0.007$) ADG and ADFI than those fed Cu from TBCC. In summary, these results suggest that increasing TBCC or Cuchelate can improve growth performance of nursery pigs and it appears that pigs provided Cu from Cu-chelate have greater ADG, ADFI, and d 35 BW, than those provided Cu from TBCC.

Keywords

nursery pig, copper, copper chelate, tri-basic copper chloride

Creative Commons License



This work is licensed under a [Creative Commons Attribution 4.0 License](https://creativecommons.org/licenses/by/4.0/).

Cover Page Footnote

The authors would like to express appreciation to Micronutrients, Indianapolis, IN, for partial funding.

Authors

C. Carpenter, J. C. Woodworth, J. M. DeRouche, M. D. Tokach, R. D. Goodband, S. S. Dritz, and J. L. Usry

Effects of Increasing Copper from Tri-basic Copper Chloride or a Copper-Chelate on Growth Performance of Nursery Pigs¹

C.B. Carpenter, J.C. Woodworth, J.M. DeRouchey, M.D. Tokach, R.D. Goodband, S.S. Dritz,² and J.L. Usry³

Summary

A total of 1,452 pigs [Group 1; 350 barrows (DNA 200 × 400; initially 13.2 lb)] and [Group 2; 1,102 barrows and gilts (PIC 1050 × 280; initially 13.3 lb)] were used to determine the effects of added Cu source and level on nursery pig performance. There were 5 pigs per pen and 10 replications per treatment in group 1, and 24 to 27 pigs per pen and 6 replications per treatment in group 2. Pens of pigs were allotted by BW to 1 of 7 dietary treatments arranged as a 2 × 3 factorial plus a control diet, with main effects of Cu source: IntelliBond-C (TBCC; Micronutrients, Indianapolis, IN), or Mintrex-Cu (Cu-chelate; Novus, St. Charles, MO) and Cu level (50, 100, or 150 ppm). Diets were corn-soybean meal-based and were fed in meal form in 2 phases (d 0 to 14 and 14 to 35). All diets contained a trace mineral premix formulated to contribute 17 ppm of Cu from CuSO₄ to the complete diet.

Overall (d 0 to 35), a Cu source × level interaction was observed (linear, $P = 0.042$) for ADG where the rate of improvement with increasing Cu was greater in pigs fed Cu-chelate diets compared to those fed TBCC diets. Increasing added Cu increased (linear, $P = 0.001$) ADG, ADFI, and improved F/G. Although Cu source did not influence F/G, pigs fed Cu from Cu-chelate had greater ($P \leq 0.007$) ADG and ADFI than those fed Cu from TBCC. In summary, these results suggest that increasing TBCC or Cu-chelate can improve growth performance of nursery pigs and it appears that pigs provided Cu from Cu-chelate have greater ADG, ADFI, and d 35 BW, than those provided Cu from TBCC.

¹ The authors would like to express appreciation to Micronutrients, Indianapolis, IN, for partial funding.

² Department of Diagnostic Medicine/Pathobiology, College of Veterinary Medicine, Kansas State University.

³ Micronutrients, Indianapolis, IN.

Introduction

The NRC⁴ reports weanling pigs have a nutritional Cu requirement of 6 ppm. However, according to Flohr et al.,⁵ many US swine nutritionists formulate nursery pig diets to contain as low as 11 ppm and as high as 250 ppm Cu. Feeding high levels of added Cu (125 to 250 ppm) in addition to what is provided by the trace mineral premix has resulted in increased ADFI and ADG. Huang et al.⁶ compared two inorganic sources [tri-basic copper chloride (TBCC) and CuSO₄] of added Cu that ranged from 11 to 327 ppm in nursery pig diets and found growth benefits from feeding added Cu but no difference in growth between sources was observed.

Organic Cu sources are argued to be more bioavailable to the young pig due to their chemical structure compared to inorganic sources. It has also been documented in other species that both TBCC and Cu-chelate are more bioavailable than more typically used sources of Cu.^{7,8} Tri-basic copper chloride and Mintrex-Cu (Cu-chelate; Novus, St. Charles, MO) differ in their chemical characteristics. Tri-basic copper chloride is an inorganic mineral source, which is non-hygroscopic and poorly soluble in water but highly soluble in acidic conditions.⁹ Mintrex-Cu is an organic form of Cu [Cu(HMTBa)₂] that has been shown to be more bioavailable to the pig because of decreased binding activity with other dietary constituents, therefore suggesting less supplementation required in comparison with inorganic minerals in nursery pigs.¹⁰ There is limited research available that directly compares the effects of increasing Cu from TBCC or Cu-chelate on growth performance of nursery pigs. Therefore, this study was designed to investigate the effects of increasing Cu from either TBCC or Cu-chelate on growth performance of nursery pigs.

Procedures

The Kansas State University Institutional Animal Care and Use Committee approved the protocols used for this study. Two groups of pigs were used for the experiment. Group 1 pigs were housed at the K-State Segregated Early Weaning Facility in Manhattan, KS. Group 2 pigs were housed in a commercial research facility in southwestern Minnesota. The research facilities were environmentally controlled. In group 1, each pen (4.0 × 4.0 ft.) had tri-bar flooring and contained one 4-hole dry self-feeder and one cup waterer to provide ad libitum access to feed and water. For group 2, each pen

⁴ NRC. 2012. Nutrient requirements of swine. 11th rev. ed. Natl. Acad. Press, Washington, DC.

⁵ Flohr, J. R.; Tokach, M. D.; Woodworth, J. C.; DeRouchey, J. M.; Dritz, S. S.; and Goodband, R. D. (2015) Vitamin and Trace Minerals: A Survey of Current Feeding Regimens, Kansas Agricultural Experiment Station Research Reports: Vol. 1: Iss. 7. <https://doi.org/10.4148/2378-5977.1127>.

⁶ Y. L. Huang, M. S. Ashwell, R. S. Fry, K. E. Lloyd, W. L. Flowers, and J. W. Spears. 2015. Effect of dietary copper amount and source on copper metabolism and oxidative stress of weanling pigs in short-term feeding. *J. Anim. Sci.* 93:2948–2955. <https://doi.org/10.4148/2378-5977.1127>.

⁷ Spears, J. W., E. B. Kegley, L. A. Mullis, and T. A. Wise, 1997. Bioavailability of copper chloride in cattle. *J. Anim. Sci.* 75 (Suppl. 1):265. (Abstr.).

⁸ Wang, Z., S. Cerrate, C. Coto, F. Yan, and P. W. Waldroup. 2007. Evaluation of Mintrex copper as a source of copper in broiler diets. *Int. J. Poult. Sci.* 6:308-313.

⁹ Miles, R. D., S. F. O'Keefe, P. R. Henry, C. B. Ammerman, and X. G. Luo. 1998. The effect of dietary supplementation with copper sulfate or tribasic copper chloride on broiler performance, relative copper bioavailability, and prooxidant activity. *Poult. Sci.* 77:416–425. doi:10.1093/ps/77.3.416.

¹⁰ J. R. Zhao, J. Harrell, G. Allee, B. Hinson, P. Winkelbauer, C. Atwell, J. D. Richards, and M. Vazquez-Anon. 2009. Effect of an organic copper source on growth performance and tissue copper concentration in nursery pigs. Mid-west Animal Science Meeting, Des Moines, IA.

(12.1 × 7.5 ft.) had plastic slatted flooring and contained one 6-hole dry self-feeder and one pan waterer to provide ad libitum access to feed and water. Dietary treatments for group 1 were manufactured at the Kansas State University O.H. Kruse Feed Technology Innovation Center, Manhattan, KS. Dietary treatments for group 2 were manufactured at a commercial feed mill located in Pipestone, MN.

In group 1, 350 barrows (DNA 200 × 400; initially 13.2 lb) were weaned at approximately 21 d of age and allotted to pen based on initial BW, with 5 pigs per pen and 10 replicate pens per treatment. In group 2, 1,102 pigs (PIC 1050 × 280; initially 13.3 lb) were weaned and randomly placed over 2 consecutive days with 24 to 27 pigs per pen and 3 replicate pens per treatment for each day. Group 1 and 2 pigs were fed a common starter diet for 7 d post-weaning. On d 7 post-weaning, pens were allotted by BW to 1 of 7 dietary treatments arranged as a 2 × 3 factorial plus one control diet, with main effects of Cu source: IntelliBond-C (TBCC; Micronutrients, Indianapolis, IN) or Mintrex-Cu (Cu chelate; Novus, St. Charles, MO) and Cu level (50, 100, or 150 ppm). Diets were corn-soybean meal-based and fed in meal form in 2 phases (d 0 to 14 and 14 to 35; Table 1). The trace mineral premix added to all diets provided complete diets with 17 ppm Cu from CuSO₄. For each group, pigs and feeders were weighed on d 0, 7, 14, 21, 28, and 35 to calculate ADG, ADFI, and F/G.

Complete diet samples were collected from a minimum of 6 feeders and combined to make 1 composite sample per treatment and phase. Each sample was then split, ground, and sent to a commercial lab for analysis of DM, CP, crude fiber, Ca, P, ether extract, ash, and Cu concentrations. Group 1 samples were sent to University of Missouri-Columbia Agriculture Chemical Laboratories (Columbia, MO) for analysis of DM, CP, crude fiber, ether extract, ash, and Cu concentrations, and Ward Laboratories Inc. (Kearney, NE) for analysis of Ca, P, and Cu concentrations. Group 1 Cu concentrations were determined by averaging the analyzed values from each lab. Group 2 samples were sent to Midwest Laboratories (Omaha, NE) for duplicate analysis of DM, CP, crude fiber, ether extract, ash, Ca, P, and Cu concentration. Final nutrient concentrations represent the combined average of the chemical analyses of diets across pig groups 1 and 2.

Data were combined from both groups and analyzed as a randomized complete block design using PROC GLIMMIX (SAS Institute, Inc., Cary, NC) with pen as the experimental unit and dietary treatment as the fixed effect. The random effect of block within group was used in the model. The main effects of source and level, as well as their interaction, were considered significant with $P < 0.05$ and marginally significant when $P < 0.10$ and ≥ 0.05 .

Results and Discussion

The chemical analyses of the complete diets were similar to the intended formulation, and Cu additions increased across dietary treatments (Table 2). However, the chemical analysis for Cu concentration was slightly higher than expected for the control diets. Total Ca and P concentrations were similar among diets across each dietary phase.

From d 0 to 14 (13.2 to 21.5 lb), there were no Cu source × level interactions observed (Table 3). Increasing added Cu increased (linear, $P \leq 0.003$) ADG and ADFI, which

tended to improve (linear, $P = 0.052$) F/G. Although Cu source did not influence F/G, pigs fed added Cu from Cu-chelate tended to have greater ($P = 0.081$) ADG and ($P = 0.053$) ADFI than those fed added Cu from TBCC.

From d 14 to 35 (21.5 to 49.7 lb), there was a source \times level interaction (linear, $P = 0.011$) for ADG reflective of a linear increase in ADG with each increasing level of Cu from Cu-chelate; however, pigs fed increasing levels of Cu from TBCC only had increased ADG at 150 ppm. Increasing Cu increased (linear, $P = 0.001$) ADFI and improved (linear, $P = 0.001$) F/G. Similar to performance from d 0 to 14, Cu source did not influence F/G; however, pigs fed Cu from Cu-chelate had greater ($P \leq 0.009$) ADG and ADFI than those fed Cu from TBCC.

Overall d 0 to 35 (13.2 to 49.7 lb), a Cu source \times level interaction was observed (linear, $P = 0.042$) for ADG where the rate of improvement with increasing Cu was greater in pigs fed Cu-chelate diets compared to those fed TBCC diets. Increasing added Cu increased (linear, $P = 0.001$) ADG, ADFI, F/G, and d 35 BW. Even though Cu source did not influence F/G; pigs fed Cu from Cu-chelate had greater ($P \leq 0.007$) ADG, ADFI, and d 35 BW than those fed added Cu from TBCC.

While research has been conducted comparing Cu sources in swine diets, it is our understanding that there is only one other published study that has directly compared the effects of increasing Cu from TBCC to a Cu-chelate (Mintrex-Cu) on nursery pig performance. Carpenter et al.¹¹ reported that nursery pigs (12.8 to 47.7 lb) fed diets containing increasing added Cu from either TBCC or Mintrex-Cu had improved ADG and ADFI in the initial nursery period and improved ADG in the overall period. In addition, Carpenter et al.¹¹ suggested that nursery pig growth performance is similar for pigs fed either TBCC or Mintrex Cu. The results from Carpenter et al.¹¹ are inconsistent with the study herein. The results from this study suggest that pigs fed diets containing increasing level of added Cu have improved growth performance in the initial, final and overall growth periods and that nursery pigs fed Cu from Cu-chelate have greater growth performance than those fed Cu from TBCC.

In this study, it appears the typical improved ADFI response to increasing added Cu helps to explain the growth response, in that increasing added Cu in the early (13.2 to 21.5 lb), late (21.5 to 49.7 lb), and overall (13.2 to 49.7 lb) nursery periods of the study improved both ADG and F/G. Others have found similar responses in previous studies with nursery pigs.^{12,13}

In summary, these results suggest that increasing TBCC or Cu-chelate can improve growth performance of nursery pigs. From our data, it also appears pigs provided Cu

¹¹ Carpenter, C. B.; Woodworth, J. C.; DeRouche, J. M.; Tokach, M. D.; Goodband, R. D.; Dritz, S. S.; and Ustry, J. (2016) "Effects of Increasing Copper from Tri-basic Copper Chloride or a Copper-Amino Acid Complex on Growth Performance of Nursery Pigs," Kansas Agricultural Experiment Station Research Reports: Vol. 2: Iss. 8. <https://doi.org/10.4148/2378-5977.1297>.

¹² Dove, C. R. 1995. The effect of Cu level on nutrient utilization of weanling pigs. *J. Anim. Sci* 73:166-171. doi:10.2527/1995.731166x.

¹³ Shelton, N. W., M. D. Tokach, J. L. Nelssen, R. D. Goodband, S. S. Dritz, J. M. DeRouche, and G. M. Hill. 2011. Effects of copper sulfate, tri-basic copper chloride, and zinc oxide on weanling pig performance. *J. Anim. Sci.* 89:2440–2451. doi:10.2527/jas.2010-3432.

from Cu-chelate may have greater ADG, ADFI, and d 35 BW than those provided Cu from TBCC. However, the study herein adds to an already conflicting database in the literature on whether nursery pig growth performance is dependent on Cu source.

Table 1. Diet composition (as-fed basis)¹

Item	Phase	
	1	2
Ingredient, %		
Corn	48.47	57.30
Soybean meal (47.7% CP)	27.68	33.73
Dried whey	10.00	---
Liquid fat ²	5.00	5.00
Limestone	0.85	0.85
Monocalcium P (21% P)	1.60	1.70
Salt	0.30	0.35
L-Lys-HCl	0.33	0.33
L-Thr	0.15	0.16
HMB ³	0.22	0.18
HP 300 ⁴	5.00	---
Vitamin premix	0.15	0.15
Trace mineral premix	0.25	0.25
Cu source ^{5,6}	---	---
Total	100.00	100.00
Calculated analysis		
Standardized ileal digestible (SID) AA, %		
Lys	1.30	1.25
Ile:Lys	63	62
Leu:Lys	122	124
Met:Lys	36	35
Met + Cys:Lys	58	58
Thr:Lys	65	65
Trp:Lys	18.4	18.4
Val:Lys	67	67
NE, kcal/lb	1,201	1,188
SID Lys:NE, g/Mcal	4.91	4.77
CP, %	21.7	21.3
Ca, %	0.85	0.80
P, %	0.78	0.75
Available P, %	0.49	0.44

¹ Phase 1 and 2 were fed from d 0 to 14 and 14 to 35, respectively. Dietary treatments were formed by adding 50, 100, or 150 ppm of Cu from either TBCC or Cu-chelate at the expense of corn. The trace mineral premix was formulated to contribute 17 ppm of Cu in the complete diet.

² Group 1 (choice white grease) and group 2 (beef tallow).

³ Hydroxymethylthio-butanoic acid, Novus International (Saint Charles, MO).

⁴ HP 300, Hamlet Protein, Findlay, OH, formulated with 3.25% SID lysine.

⁵ Mintrex Cu, copper methionine hydroxy analogue (St. Charles, MO).

⁶ IntelliBond-C, TBCC (Micronutrients, Indianapolis, IN).

Table 2. Chemical analysis of diets, as-fed basis¹

Item	Phase 1							Phase 2						
	Added Cu, ppm													
	Control	TBCC ²			Cu-chelate ³			Control	TBCC ²			Cu-chelate ³		
	0	50	100	150	50	100	150	0	50	100	150	50	100	150
DM, %	87.5	87.2	86.8	86.7	86.6	86.6	86.8	86.7	86.4	85.8	85.6	86.2	86.1	85.7
CP, %	21.9	21.0	20.6	20.9	20.8	20.8	20.8	19.2	20.4	19.9	21.9	18.9	19.6	21.2
Crude fiber, %	1.9	2.0	1.5	1.6	2.0	2.1	2.1	2.6	2.4	2.1	2.4	1.9	2.0	2.0
Ether extract, %	7.1	6.8	7.2	6.9	6.6	7.6	6.9	6.7	6.5	6.3	5.3	6.3	5.6	5.6
Ash, %	5.7	5.7	5.6	6.0	5.9	5.8	5.9	5.4	5.4	5.3	5.7	5.4	5.7	5.6
Ca, %	0.85	0.78	0.77	0.82	0.75	0.86	0.76	0.75	0.97	1.00	0.76	1.01	0.87	0.91
P, %	0.76	0.74	0.68	0.79	0.73	0.83	0.78	0.62	0.72	0.79	0.77	0.81	0.66	0.88
Cu, ppm	40	88	140	145	104	155	204	40	75	119	174	100	135	230

¹ Multiple samples of each diet were collected, blended and subsampled, and analyzed (Missouri Agricultural Experimentation Lab, Colombia, MO; Ward Laboratories, Kearney, NE; and Midwest Labs, Omaha, NE). All values represent the combined average of the chemical analyses of diets for the experiment.

² IntelliBond-C, tri-basic copper chloride (Micronutrients, Indianapolis, IN).

³ Mintrex Cu, copper methionine hydroxy analogue (St. Charles, MO).

Table 3. Effects of increasing Cu from tri-basic copper chloride (TBCC) or Cu-chelate on growth performance of pigs¹

Item	Added Cu, ppm							SEM	Probability, <i>P</i> <					
	Control	TBCC ²			Cu-chelate ³				Cu	Cu level		Source × level		
		0	50	100	150	50	100			150	Source	Linear	Quadratic	Linear
BW, lb														
d 0	13.2	13.2	13.2	13.2	13.3	13.3	13.2	0.12	0.112	0.998	0.613	0.372	0.402	
d 14	21.1	21.0	21.5	21.7	21.5	21.8	22.0	0.38	0.030	0.001	0.941	0.462	0.318	
d 35	47.4	49.1	49.4	50.3	49.9	50.6	51.5	0.54	0.001	0.001	0.031	0.019	0.384	
d 0 to 14														
ADG, lb	0.55	0.55	0.59	0.60	0.59	0.60	0.62	0.023	0.081	0.001	0.953	0.780	0.377	
ADFI, lb	0.79	0.78	0.82	0.82	0.82	0.82	0.85	0.032	0.053	0.003	0.917	0.511	0.718	
F/G	1.42	1.41	1.40	1.36	1.39	1.38	1.38	0.023	0.620	0.052	0.887	0.720	0.302	
d 14 to 35														
ADG, lb	1.25	1.33	1.33	1.36	1.35	1.37	1.40	0.015	0.001	0.001	0.004	0.011	0.568	
ADFI, lb	1.85	1.92	1.90	1.95	1.96	1.97	1.99	0.030	0.009	0.001	0.159	0.167	0.222	
F/G	1.49	1.45	1.43	1.44	1.45	1.44	1.42	0.018	0.891	0.001	0.206	0.533	0.452	
d 0 to 35														
ADG, lb	0.97	1.02	1.03	1.05	1.05	1.06	1.09	0.013	0.001	0.001	0.027	0.042	0.334	
ADFI, lb	1.43	1.46	1.47	1.49	1.50	1.51	1.53	0.021	0.007	0.001	0.279	0.211	0.277	
F/G	1.47	1.44	1.42	1.42	1.44	1.43	1.41	0.013	0.730	0.001	0.167	0.623	0.710	

¹ A total of 1,452 pigs [Group 1; 350 barrows (DNA 200 × 400; initially 13.2 lb)] and [Group 2; 1,102 pigs (PIC 1050 × 280; initially 13.3 lb)] were used in two 35-d growth studies. Data were combined across the 2 groups with 5 pigs per pen and 10 replications per treatment in group 1; and 24 to 27 pigs per pen and 6 replications per treatment in group 2. The treatment design was the same across both groups of pigs. The trace mineral premix was formulated to contribute 17 ppm of Cu to the complete diet.

² Intellibond-C, tri-basic copper chloride (Micronutrients, Indianapolis, IN).

³ Mintrex Cu, copper methionine hydroxy analogue (St. Charles, MO).